

Brethine/Terbutaline Preterm Labor [1348]

General

Nursing

Vital Signs

| | |
|---|--|
| <input type="checkbox"/> Vital signs - T/P/R/BP | Routine, Every 15 min With each dose of Brethine and then per floor policy, L&D Pre-Delivery |
|---|--|

Activity

| | |
|--|--|
| <input type="checkbox"/> Strict bed rest | Routine, Until discontinued, Starting S, L&D Pre-Delivery |
| <input checked="" type="checkbox"/> Bed rest | Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges L&D Pre-Delivery |

Nursing Care

| | |
|---|---|
| <input checked="" type="checkbox"/> Monitor fetal heart tones | Routine, Once Type: Continuous L&D Pre-Delivery |
| <input checked="" type="checkbox"/> Tocometry | Routine, Until discontinued, Starting S Type: Continuous L&D Pre-Delivery |
| <input checked="" type="checkbox"/> Oral hydration of at least 1000 millileters within 60 minutes | Routine, Once For 1 Occurrences, L&D Pre-Delivery |

Notify

| | |
|--|---|
| <input checked="" type="checkbox"/> Notify Physician | Routine, Until discontinued, Starting S, For maternal pulse greater than 120 bpm or any significant change in maternal status; FHR greater than 180 or any significant change in fetal status; Persistent regular contraction pattern., L&D Pre-Delivery |
|--|---|

IV Fluids

IV Fluids

| | |
|--|--|
| <input checked="" type="checkbox"/> lactated ringer's bolus and infusion | "And" Linked Panel |
| <input checked="" type="checkbox"/> lactated Ringer's bolus | 1,000 mL, intravenous, for 30 Minutes, once, For 1 Doses, L&D Pre-Delivery |
| <input checked="" type="checkbox"/> lactated Ringer's infusion | 125 mL/hr, intravenous, continuous, L&D Pre-Delivery |

Medications

Medications

| | |
|---|--|
| <input type="checkbox"/> Option 1: terbutaline (BRETHINE) SQ Initial Doses and Oral Maintenance Doses | "Followed by" Linked Panel |
| <input type="checkbox"/> Initial Doses - terbutaline (BRETHINE) injection - 1st dose | 0.25 mg, subcutaneous, once, For 1 Doses, L&D Pre-Delivery Stop at first dose if effective. Hold if maternal heart rate is GREATER than 120. |
| <input type="checkbox"/> terbutaline (BRETHINE) injection - 2nd dose | 0.25 mg, subcutaneous, once, Starting H+20 Minutes, For 1 Doses, L&D Pre-Delivery Hold if maternal heart rate is GREATER than 120. |
| <input type="checkbox"/> terbutaline (BRETHINE) injection - 3rd dose | 0.25 mg, subcutaneous, once, Starting H+40 Minutes, For 1 Doses, L&D Pre-Delivery Hold if maternal heart rate is GREATER than 120. |
| <input type="checkbox"/> Maintenance Dose - terbutaline (BRETHINE) tablet | 5 mg, oral, every 4 hours, Starting H+41 Minutes, L&D Pre-Delivery Hold if maternal heart rate is GREATER than 120. |

Option 2: terbutaline (BRETHINE) SQ and Oral Doses

| | |
|---|--|
| <input type="checkbox"/> terbutaline (BRETHINE) injection | 0.25 mg, subcutaneous, every 20 min, For 3 Doses, L&D Pre-Delivery Hold dose if Maternal Pulse is GREATER than 120. |
| <input type="checkbox"/> terbutaline (BRETHINE) tablet | 5 mg, oral, every 4 hours scheduled, For 3 Doses, L&D Pre-Delivery Hold Dose if Maternal Pulse is GREATER than 120. |

VTE

Labs

Microbiology

| | |
|---|--|
| <input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture | Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery |
|---|--|

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For additional Physician Consult orders use sidebar

Physician Consults

| | |
|---|---|
| <input type="checkbox"/> Consult to Neonatology | Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? L&D Pre-Delivery |
|---|---|

Additional Orders