

**General**

**Common Present on Admission Diagnosis**

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

**Elective Outpatient, Observation, or Admission (Single Response)**

<input type="checkbox"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

**Admission or Observation (Single Response)**

Patient has active outpatient status order on file

- |  |  |
|--|--|
| <input type="checkbox"/> Admit to Inpatient  | Diagnosis:<br>Admitting Physician:<br>Level of Care:<br>Patient Condition:<br>Bed request comments:<br>Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.<br>PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Diagnosis:<br>Admitting Physician:<br>Patient Condition:<br>Bed request comments:<br>PACU & Post-op  |
| <input type="checkbox"/> Outpatient in a bed - extended recovery                   | Diagnosis:<br>Admitting Physician:<br>Bed request comments:<br>PACU & Post-op  |
| <input type="checkbox"/> Transfer patient  | Level of Care:<br>Bed request comments:<br>Scheduling/ADT  |
| <input type="checkbox"/> Return to previous bed                                    | Routine, Until discontinued, Starting S, Scheduling/ADT  |

**Admission (Single Response)**

Patient has active status order on file

- |   |  |
|---|--|
| <input type="checkbox"/> Admit to inpatient     | Diagnosis:<br>Admitting Physician:<br>Level of Care:<br>Patient Condition:<br>Bed request comments:<br>Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.<br>PACU & Post-op |
| <input type="checkbox"/> Transfer patient       | Level of Care:<br>Bed request comments:<br>Scheduling/ADT  |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT  |

**Transfer (Single Response)**

Patient has active inpatient status order on file

- |   |   |
|---|---|
| <input type="checkbox"/> Transfer patient       | Level of Care:<br>Bed request comments:<br>Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT   |

**Code Status**

- |   |  |
|---|--|
| <input type="checkbox"/> Full Code                | Code Status decision reached by:<br>Post-op            |
| <input type="checkbox"/> DNR (Do Not Resuscitate) |  |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?<br>Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

### Isolation

<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

### Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

## Nursing

### Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol For 4 Hours Every 1 hour x 4, Post-op
<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol Every 4 hours to begin after initial Q1H x 4 checks, Post-op
<input checked="" type="checkbox"/> Peripheral vascular assessment - bilateral arm	Routine, Every 2 hours Bilateral arm: Until full motor function and sensation regained, if spinal anesthesia administered., Post-op
<input checked="" type="checkbox"/> Peripheral vascular assessment - bilateral leg	Routine, Every 2 hours Bilateral leg; until full motor functions and sensation regained, if spinal anesthesia administered, Post-op

### Activity

<input checked="" type="checkbox"/> Ambulate	Routine, 3 times daily Specify: As tolerated
--	--

### Nursing

<input type="checkbox"/> Saline lock IV	Routine, Continuous When taking PO fluids well, Post-op
<input type="checkbox"/> Foley catheter care	Routine, Until discontinued, Starting S Orders: Maintain Post-op
<input type="checkbox"/> Foley catheter care	Routine, As needed Orders: Maintain PRN for clots, Post-op
<input checked="" type="checkbox"/> Strict intake and output	Routine, Every hour For 4 Occurrences Every 1 hour x 4, Post-op
<input checked="" type="checkbox"/> Strict intake and output	Routine, Every 4 hours Every 4 hours to begin after every 1 hour x 4., Post-op

<input checked="" type="checkbox"/> Turn cough deep breathe	Routine, Every 2 hours, Post-op
<input type="checkbox"/> Incentive spirometry	Routine, Every hour while awake 10 deep puffs every hour, Post-op
<input type="checkbox"/> Apply ice pack	Routine, Until discontinued, Starting S Affected area: Waking hours only? Nurse to schedule? Special Instructions: To perineum, Post-op

### Diet

<input checked="" type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Full Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Encourage fluids, Post-op
--	--

## IV Fluids

### IV Fluids

<input type="checkbox"/> dextrose 5%-0.45% sodium chloride infusion	intravenous, continuous, Post-op
---	----------------------------------

## Medications

### Moderate Pain (Pain Score 4 - 6) (Single Response)

<input type="checkbox"/> morphine injection	2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op May be given IM if no IV access.
<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 4 hours PRN, moderate pain (score 4-6), Post-op

### Severe Pain (Pain Score 7-10) (Single Response)

<input type="checkbox"/> traMADol (ULTRAM) tablet	50 mg, oral, every 8 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> morphine 2 mg/mL injection	2 mg, intravenous, every 4 hours PRN, severe pain (score 7-10), Post-op May be given IM if no IV access.
<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	2 tablet, oral, every 4 hours PRN, severe pain (score 7-10), Post-op

### Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.
--	--

#### Antiemetics - HMSL, HMWB Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Antiemetics - HMSTJ Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Antispasmodics

<input type="checkbox"/> oxybutynin (DITROPAN) tablet	5 mg, oral, 3 times daily PRN, bladder spasms, bladder spasms, Post-op Discontinue when Foley removed.
<input type="checkbox"/> belladonna alkaloids-opium (B&O SUPPRETTES) 16.2-60 MG suppository	60 mg, rectal, every 6 hours PRN, bladder spasms, Post-op Discontinue when Foley removed

#### Bowel Care

<input type="checkbox"/> docusate sodium (COLACE) 100 mg capsule	100 mg, oral, 2 times daily, Post-op
--	--------------------------------------

#### Insomnia: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep, Post-op
---	---

ramelteon (ROZEREM) tablet 8 mg, oral, nightly PRN, sleep, Post-op

**Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)**

ramelteon (ROZEREM) tablet 8 mg, oral, nightly PRN, sleep, Post-op

**VTE**

**DVT Risk and Prophylaxis Tool (Single Response)**

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

PACU & Post-op

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

#### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

<input type="checkbox"/> Low Risk (Single Response)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

## Labs

## Cardiology

## Imaging

## Other Studies

## Respiratory

## Rehab

## Consults

For Physician Consult orders use sidebar

### Consults

<input checked="" type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op, Assess discharge needs
--	--

## Additional Orders