

Nursing

Vital Signs

Vital signs - T/P/R/BP Routine, Every 2 hours For 999 Occurrences
Every 2 hours for 8 hours, then every 4 hours for 48 hours, then per unit standards

Activity

Activity as tolerated Routine, Until discontinued, Starting S
Specify: Activity as tolerated

Activity - Out of bed to chair for all meals daily Routine, 3 times daily
Specify: Other activity (specify)
Other: Out of bed to chair for all meals daily

Ambulate Routine, 4 times daily
Specify:
In hallway as tolerated, with assistance if needed

Patient may shower Routine, Daily
Specify:
Additional modifier:
Shower in am: For patients with pacer wires, cover wires with occlusive waterproof dressing prior to shower. Cover central line

Bed rest Routine, Until discontinued, Starting S
Bathroom Privileges:

Nursing

Telemetry **"And" Linked Panel**

Telemetry monitoring Routine, Continuous
Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box)
Reason for telemetry: Post cardiac surgery
Can be off of Telemetry for tests and baths? Yes

Telemetry Additional Setup Information Routine, Continuous
High Heart Rate (BPM): 120
Low Heart Rate(BPM): 50
High PVC's (per minute): 10
High SBP(mmHg): 175
Low SBP(mmHg): 100
High DBP(mmHg): 95
Low DBP(mmHg): 40
Low Mean BP: 60
High Mean BP: 120
Low SPO2(%): 94

Daily weights Routine, Daily

Intake and output Routine, Every shift

Assess operative site Routine, Every 8 hours
Care of incision site per CV protocol

Site care Routine, Per unit protocol
Site:
Care of epicardial wire site and wire insulation per CV protocol

Saline lock IV Routine, Continuous

Foley catheter - discontinue Routine, Once
If present 1) Remove Foley cath POD 1 or POD 2; 2)
Document reason for not removing foley (Must be documented on POD 1 or POD 2)

Nasogastric tube maintenance Routine, Until discontinued, Starting S
Tube Care Orders:

Chest tube to continuous suction Routine, Until discontinued, Starting S
Level of suction: 20 cm H2O

<input type="checkbox"/> Tube site care	Routine, Every 8 hours Chest tube per CV protocol
<input type="checkbox"/> Discontinue pacemaker generator and insulate pacer wires	Routine, Until discontinued, Starting S
Notify	
<input type="checkbox"/> Notify Physician - Consultants and resident of patient's location	Routine, Once For 1 Occurrences, Consultants and resident of patient location
<input type="checkbox"/> Notify - Physician Assistant and/or nurse practitioner of patient's location	Routine, Once For 1 Occurrences, Physician Assistant and/or nurse practitioner of patient location
<input type="checkbox"/> Notify Physician - NP - PA If potassium level is less than 3.6 milliEquivalents per Liter or greater than 5.0 milliEquivalents per Liter.	Routine, Once For 1 Occurrences, NP - PA If potassium level is less than 3.6 milliEquivalents per Liter or greater than 5.0 milliEquivalents per Liter.

Diet

<input type="checkbox"/> Diet -Clear liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> Diet - Full liquids	Diet effective now, Starting S Diet(s): Full Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> Diet - Heart Healthy	Diet effective now, Starting S Diet(s): Heart Healthy Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> Diet - Renal (80gm Pro, 2-3gm Na, 2-3gm K)	Diet effective now, Starting S Diet(s): Renal (80GM Pro, 2-3GM Na, 2-3GM K) Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> Diet - 1800 Carb Control Diabetic	Diet effective now, Starting S Diet(s): Other Diabetic/Cal Diabetic/Calorie: 1800 Kcal/202 gm Carbohydrate Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:

Medications

ACE/ARB Inhibitors (Single Response)

<input type="checkbox"/> lisinopril (PRINIVIL,ZESTRIL) tablet	10 mg, oral, daily HOLD parameters for this order: Contact Physician if:
<input type="checkbox"/> enalapril (VASOTEC) tablet	5 mg, oral, daily HOLD parameters for this order: Contact Physician if:
<input type="checkbox"/> losartan (COZAAR) tablet	25 mg, oral, daily HOLD parameters for this order: Contact Physician if:

Anti-platelet

<input type="checkbox"/> clopidogrel (PLAVIX) tablet	75 mg, oral, daily
--	--------------------

aspirin (ECOTRIN) enteric coated tablet 81 mg, oral, daily

Beta-Blocker (Single Response)

metoprolol tartrate (LOPRESSOR) tablet 25 mg, oral, 2 times daily at 0600, 1800
DO NOT administer if heart rate is LESS than 60 or systolic blood pressure is LESS than 100;
HOLD parameters for this order: Hold Parameters requested
HOLD for: Other
Other Systolic Blood Pressure (in mmHg): 100 mmHg
HOLD for Heart Rate LESS than: Other
Other Heart Rate (in bpm): 60 bpm
Contact Physician if:

carvedilol (COREG) tablet 3.125 mg, oral, 2 times daily
DO NOT administer if heart rate is LESS than 60; systolic blood pressure is LESS than 100
HOLD parameters for this order: Hold Parameters requested
HOLD for: Other
Contact Physician if:

Diuretics

furosemide (LASIX) Oral or IV (Single Response)

furosemide (LASIX) tablet 40 mg, oral, daily
 furosemide (LASIX) IV 40 mg, intravenous, daily

Statin (Single Response)

simvastatin (ZOCOR) tablet 40 mg, oral, nightly
 atorvastatin (LIPITOR) tablet 40 mg, oral, nightly

Labs

Laboratory Tomorrow AM

Basic metabolic panel AM draw, Starting S+1 For 1 Occurrences
 CBC hemogram AM draw, Starting S+1 For 1 Occurrences
 Prothrombin time with INR AM draw, Starting S+1 For 1 Occurrences

Cardiology

ECG

ECG 12 lead Routine, Once For 1 Occurrences
Clinical Indications:
Interpreting Physician:

Imaging

X-Ray

Chest 1 Vw Portable Routine, 1 time imaging For 1 Occurrences

Respiratory

Respiratory Therapy

Oxygen therapy Routine, Continuous
Device 1: Nasal Cannula
Rate in liters per minute: 2 Lpm
Rate in tenths of a liter per minute:
O2 %:
Device 2:
Device 3:
Titrate to keep O2 Sat Above: 92%
Indications for O2 therapy:
Try weaning patient daily

Incentive spirometry Routine, Every hour while awake For 999 Occurrences

<input checked="" type="checkbox"/> Encourage deep breathing and coughing	Routine, Every hour For 999 Occurrences
<input type="checkbox"/> PEP	Routine, As directed 20 times every hour while awake

Consults

For Physician Consult orders use sidebar

Consults

<input checked="" type="checkbox"/> Consult Cardiac Rehab Phase 1	Routine, Once Clinical Indications: for assistance with mobility
<input type="checkbox"/> Consult to PT eval and treat	Special Instructions: Weight Bearing Status: