

General

Medications

Pharmacy Consult (Selection Required)

Pharmacy consult to monitor dofetilide (TIKOSYN) therapy STAT, Until discontinued, Starting S

Medications prior to initiation of dofetilide (TIKOSYN)

<input type="checkbox"/> Magnesium less than 2 mEq/L: Magnesium sulfate 2 gm IVPB over 2 hours once	2 g, intravenous, once PRN, For Magnesium less than 2 mEq/L prior to initiation of dofetilide (TIKOSYN)
<input type="checkbox"/> Potassium 3.5 mEq/L to 4 mEq/L:potassium chloride (KAYCIEL) 20 mEq solution	20 mEq, oral, once PRN, For Potassium 3.5 mEq/L to 4 mEq/L prior to initiation of dofetilide (TIKOSYN)
<input type="checkbox"/> Potassium less than 3.5 mEq/L:potassium chloride (KAYCIEL) 40 mEq solution	
<input type="checkbox"/> potassium chloride (KAYCIEL) 20 mEq/15 mL solution	40 mEq, oral, once PRN, For Potassium less than 3.5 mEq/L prior to initiation of dofetilide (TIKOSYN)
<input type="checkbox"/> Potassium level	Once For 1 Occurrences Recheck potassium level 2 hours after 40 mEq potassium chloride dose administered prior to initiation of dofetilide (TIKOSYN)
<input type="checkbox"/> Notify Clinical Pharmacist if recheck potassium level is less than 3.5	Routine, Once For 1 Occurrences, Notify Clinical Pharmacist if recheck potassium level is less than 3.5 mEq/L

dofetilide (TIKOSYN) therapy

<input type="checkbox"/> dofetilide (TIKOSYN) 125 mcg capsule	125 mcg, oral, every 12 hours scheduled Initiation: Dofetilide is contraindicated for QTc greater than 440msec in patients without a ventricular conduction abnormality. Dofetilide is contraindicated for QTc greater than 500msec in patients with a ventricular conduction abnormality. Home continuation: Dofetilide in contraindicated for QTc greater than 500msec in patients without a ventricular conduction abnormality. Dofetilide is contraindicated for QTc greater than 550 msec in patients with a ventricular conduction abnormality.
<input type="checkbox"/> dofetilide (TIKOSYN) 250 mcg capsule	250 mcg, oral, every 12 hours scheduled Initiation: Dofetilide is contraindicated for QTc greater than 440msec in patients without a ventricular conduction abnormality. Dofetilide is contraindicated for QTc greater than 500msec in patients with a ventricular conduction abnormality. Home continuation: Dofetilide in contraindicated for QTc greater than 500msec in patients without a ventricular conduction abnormality. Dofetilide is contraindicated for QTc greater than 550 msec in patients with a ventricular conduction abnormality.
<input type="checkbox"/> dofetilide (TIKOSYN) 500 mcg capsule	500 mcg, oral, every 12 hours scheduled Initiation: Dofetilide is contraindicated for QTc greater than 440msec in patients without a ventricular conduction abnormality. Dofetilide is contraindicated for QTc greater than 500msec in patients with a ventricular conduction abnormality. Home continuation: Dofetilide in contraindicated for QTc greater than 500msec in patients without a ventricular conduction abnormality. Dofetilide is contraindicated for QTc greater than 550 msec in patients with a ventricular conduction abnormality.

Nursing

Vital Signs

"And" Linked Panel	
<input checked="" type="checkbox"/> Telemetry	
<input checked="" type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes
<input checked="" type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94

IV Fluids

VTE

Labs

STAT Labs if not drawn in past 24 hours

<input checked="" type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences STAT if not available in the last 24 hours
<input checked="" type="checkbox"/> Magnesium	STAT For 1 Occurrences STAT if not available in the last 24 hours

Repeating Labs

<input checked="" type="checkbox"/> Basic metabolic panel	AM draw repeats For 2 Days
<input checked="" type="checkbox"/> Magnesium	AM draw repeats For 2 Days

Cardiology

ECG

<input type="checkbox"/> ECG 12 lead	STAT, Once For 1 Occurrences Clinical Indications: Cardiac Arrhythmia Interpreting Physician: Baseline ECG, if not already within 24 hours
<input checked="" type="checkbox"/> ECG 12 lead	Routine, Conditional Frequency, Starting S For 6 Occurrences Clinical Indications: Cardiac Arrhythmia Interpreting Physician: Obtain 2 hours after each dose of dofetilide for 3 days

EKG

<input checked="" type="checkbox"/> ECG 12 lead	STAT, Once For 1 Occurrences Clinical Indications: Cardiac Arrhythmia Interpreting Physician: Baseline ECG, if not already within 24 hours
<input checked="" type="checkbox"/> ECG 12 lead	Routine, Conditional Frequency, Starting S For 6 Occurrences Clinical Indications: Cardiac Arrhythmia Interpreting Physician: Obtain 2 hours after each dose of dofetilide for 3 days

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders