

AmIODArone (CORDArone) for ventricular dysrhythmias [1590]

General

Notify Physician (Single Response) (Selection Required)

- | | |
|---|--|
| <input checked="" type="checkbox"/> Notify provider if amlODarone therapy held for heart rate less than 60 or pauses greater than 2.5 seconds | Routine, Until discontinued, Starting S, Notify provider if amlODarone therapy held for heart rate less than 60 or pauses greater than 2.5 seconds |
|---|--|

Nursing

Vital Signs

[X] Telemetry		"And" Linked Panel
<input checked="" type="checkbox"/> Telemetry monitoring		Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes
<input checked="" type="checkbox"/> Telemetry Additional Setup Information		Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94

IV Fluids

Medications

amlODarone (CORDArone) Continous Infusion (Single Response)

Default infusion rate is 1 mg/min. Reduce dose to 0.5 mg/min for infusions longer than 6 hours, if appropriate.

()		"Followed by" Linked Panel
<input type="checkbox"/> amlODarone (CORDArone) 150 mg BOLUS followed by Continuous Infusion		
<input type="checkbox"/> amlODarone (CORDArone) 150 mg BOLUS		150 mg, intravenous, for 10 Minutes, once, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infuson for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours. Administer over 10 minutes.
<input type="checkbox"/> amlODarone 1.8 mg/mL (STANDARD concentration) infusion		1 mg/min, intravenous, continuous, Starting H+10 Minutes Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infuson for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours.
<input type="checkbox"/> NO BOLUS - amlODarone (CORDArone) Continuous Infusion		1 mg/min, intravenous, continuous Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infuson for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours.

Labs

Labs (Baseline checks required if not already completed)

- | | |
|--|--|
| <input checked="" type="checkbox"/> Hepatic function panel | Once
Draw for baseline if not already completed within 24 hours |
|--|--|

[X] TSH

Once
Draw for baseline if not already completed within 24 hours

Cardiology

ECG

[X] ECG 12 lead

STAT, Once For 1 Occurrences
Clinical Indications: Cardiac Arrhythmia
Interpreting Physician:
Baseline ECG if not already completed

Imaging

X-ray

[X] Chest 2 Vw

Routine, 1 time imaging For 1

Other Studies

Respiratory

Consults

For Physician Consult orders use sidebar

Additional Orders