TEE - Transesophageal Echocardiogram Pre Procedure [1671]

Note: Patients on coumadin require an INR within 24 hours of the TEE procedure. Patients on heparin require a PTT or Xa Factor within 12 hours of the TEE procedure.

| General | |
|--|---|
| Nursing | |
| Nursing Care | |
|] Saline lock IV | Routine, Continuous |
| Hold tube feedings for a minimum of 6 hours prior to | Routine, Once |
| exam | Hold tube feedings for a minimum of 6 hours prior to exam. |
| Diet | |
| [X] NPO | Diet effective now, Starting S |
| | NPO: |
| | Pre-Operative fasting options: |
| | Minimum of 6 hours prior to the transesophageal |
| | echocardiogram exam |
| V Fluids | |
| V Fluids | |
|] sodium chloride 0.9 % infusion | 30 mL/hr, intravenous, continuous |
| Medications | |
| VTE | |
| Labs | |
| Labs: Day of TEE Procedure | |
|] Basic metabolic panel | Once |
| CBC hemogram | Once |
| Prothrombin time with INR | Once |
| Partial thromboplastin time | Once |
| | Office |
| Cardiology | |
| ΓΕΕ | |
| X] Echocardiogram transesophageal | Routine, 1 time imaging |
| | NPO 6 hours prior to the exam |
| Anesthesia Consult and Consent | |
| Anesthesia Consult and Consent | |
| [] Consult Anesthesiology | Reason for Consult? Sedation - Transesophageal |
| | Echocardiogram |
| | Patient/Clinical information communicated? |
| | Patient/clinical information communicated? |
| [] Complete consent for | Routine, Once |
| | Procedure: Anesthesia |
| | Diagnosis/Condition: |
| | Physician: |
| | Risks, benefits, and alternatives (as outlined by the Texas |
| | Medical Disclosure Panel, as appears on Houston Methodis |
| | Medical/Surgical Consent forms) were discussed with |
| | patient/surrogate? |
| | |

Additional Orders