

Case Request

Case Request

Case request ep lab Details

Nursing

Consents

Complete consent for left atrial appendage occlusion device Routine, Once
Procedure: left atrial appendage occlusion device
Diagnosis/Condition:
Physician:
Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate?
Pre-Procedure

Complete consent for temporary pacemaker implantation Routine, Once
Procedure: Temporary pacemaker implantation
Diagnosis/Condition:
Physician:
Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate?
Pre-Procedure

Complete consent for Routine, Once
Procedure:
Diagnosis/Condition:
Physician:
Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate?
Pre-Procedure

Vital Signs

Vital signs - T/P/R/BP Routine, Per unit protocol, Pre-Procedure

Telemetry - PACU/PostOp

Telemetry **"And" Linked Panel**

Telemetry monitoring Routine, Continuous
Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box)
Reason for telemetry:
Can be off of Telemetry for tests and baths? Yes
PACU & Post-op

Telemetry Additional Setup Information Routine, Continuous
High Heart Rate (BPM): 120
Low Heart Rate(BPM): 50
High PVC's (per minute): 10
High SBP(mmHg): 175
Low SBP(mmHg): 100
High DBP(mmHg): 95
Low DBP(mmHg): 40
Low Mean BP: 60
High Mean BP: 120
Low SPO2(%): 94
PACU & Post-op

Telemetry - Pre-Procedure

	"And" Linked Panel
<input checked="" type="checkbox"/> Telemetry	
<input checked="" type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes Pre-Procedure
<input checked="" type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94 Pre-Procedure

Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: (Hold diabetic medications morning of procedure), Pre-Procedure
<input type="checkbox"/> Diet-Clear Liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: For breakfast, Pre-Procedure

Pre Procedure Nursing Interventions

<input type="checkbox"/> Insert peripheral IV-left arm	Routine, Once Insert large bore (at least 20 gauge) antecubital on left arm, Pre-Procedure
<input type="checkbox"/> Insert peripheral IV-right arm	Routine, Once Insert large bore (at least 20 gauge) antecubital on right arm, Pre-Procedure
<input type="checkbox"/> Insert and maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Pre-Procedure
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain Pre-Procedure
<input type="checkbox"/> Void on call to Cath Lab	Routine, Once, Pre-Procedure

Medications

PreOp Antibiotics (Single Response)

Ceftriaxone + Vancomycin

<input checked="" type="checkbox"/> cefTRIAxone (ROCEPHIN) IV	2 g, intravenous, for 30 Minutes, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input checked="" type="checkbox"/> vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis Indication: Increased MRSA rate (operation specific)
() Aztreonam + Vancomycin - For severe beta-lactam allergy	
<input type="checkbox"/> aztreonam (AZACTAM) IV	2 g, intravenous, once, For 1 Doses, Pre-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Pre-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis Indication: Increased MRSA rate (operation specific)
() For Vancomycin Allergy - ceFAZolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	
<input type="checkbox"/> ceFAZolin (ANCEF) IV	2 g, intravenous Reason for Therapy: Surgical Prophylaxis
() For Vancomycin Allergy - ceFAZolin (ANCEF) IV - For Patients GREATER than 120 kg	
<input type="checkbox"/> ceFAZolin (ANCEF) IV	3 g, intravenous Reason for Therapy: Surgical Prophylaxis

Labs

Laboratory on Arrival to Unit

<input checked="" type="checkbox"/> Hemoglobin	Once For 1 Occurrences, Pre-Procedure
<input checked="" type="checkbox"/> Prothrombin time with INR	Once For 1 Occurrences, Pre-Procedure
<input checked="" type="checkbox"/> Creatinine level	Once For 1 Occurrences, Pre-Procedure
<input checked="" type="checkbox"/> Albumin level	Once For 1 Occurrences, Pre-Procedure
<input checked="" type="checkbox"/> Platelet count	Once For 1 Occurrences, Pre-Procedure
<input type="checkbox"/> Basic metabolic panel	Once, Pre-Procedure
<input type="checkbox"/> CBC with platelet and differential	Once, Pre-Procedure
<input type="checkbox"/> Prothrombin time with INR	Once, Pre-Procedure

Cardiology

ECG

<input checked="" type="checkbox"/> ECG Pre/Post Op	Routine, Once Clinical Indications: Interpreting Physician: On arrival, Pre-Procedure
---	--