

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="checkbox"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Diagnosis:
Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Diagnosis:
Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

- | | |
|--|--|
| <input type="checkbox"/> Full Code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?
Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum, Post-op
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Notify Physician

<input checked="" type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: Temperature less than: Systolic BP greater than: 150 Systolic BP less than: 90 Diastolic BP greater than: Diastolic BP less than: MAP less than: Heart rate greater than (BPM): 100 Heart rate less than (BPM): 60 Respiratory rate greater than: 26 Respiratory rate less than: 12 SpO2 less than: 95
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Difficulty breathing, Post-op
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Nausea/vomiting not controlled, Post-op
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Inability to keep down oral medications, Post-op

Vitals

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours For 2 Occurrences, Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 8 hours, Starting H+8 Hours, Post-op

Activity

<input type="checkbox"/> Out of bed	Routine, 3 times daily For Until specified Specify: Out of bed Out of bed for 1 hour at a time, or as tolerated, Post-op
-------------------------------------	--

Nursing Assessment

<input type="checkbox"/> Neurological assessment	Routine, Every 4 hours Assessment to Perform: Post-op
<input type="checkbox"/> Strict intake and output	Routine, Every 8 hours Record all oral and IV fluid intake and all output including drains, tubes, nasogastric tube and foley catheter, Post-op
<input type="checkbox"/> Height and weight	Routine, Once, Post-op

Nursing Interventions

<input type="checkbox"/> Head of bed	Routine, Until discontinued, Starting S Head of bed: Unless contraindicated, Post-op
<input type="checkbox"/> Nasogastric tube insertion	Routine, Once Type: Post-op
<input type="checkbox"/> Patient education	Routine, Once Patient/Family: Education for: Post-op
<input type="checkbox"/> Drain care	Routine, Until discontinued, Starting S Drain 1: Drain 2: Drain 3: Drain 4: All Drains: Post-op

Wound Care

<input type="checkbox"/> Provide suture tray to patient bedside	Routine, Once, Post-op
---	------------------------

Diet

<input type="checkbox"/> Diet -	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
---------------------------------	--

IV Fluids

IV Fluids

<input type="checkbox"/> sodium chloride 0.9 % infusion	intravenous, continuous, Post-op
---	----------------------------------

Medications

Medications for Cushing's / Hypercortisolism patient

<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection	50 mg, intravenous, every 6 hours
--	-----------------------------------

Pain Management

<input type="checkbox"/> For Mild Pain - acetaminophen (TYLENOL) tablet	500 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op
<input type="checkbox"/> For Moderate Pain - acetaminophen (TYLENOL) tablet	1,000 mg, oral, every 8 hours PRN, moderate pain (score 4-6), Post-op

For Severe Pain - acetaminophen-codeine (TYLENOL WITH CODEINE #3) 300-30 mg per tablet 1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

URL: "\appt1.pdf"

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op

LOW Risk of DVT (Selection Required)
Low Risk Definition
Age less than 60 years and NO other VTE risk factors

Low Risk (Single Response) (Selection Required)

Low risk of VTE Routine, Once
Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
PACU & Post-op

MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

Moderate risk of VTE Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis **"And" Linked Panel**
BUT order Sequential compression device

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

Contraindications exist for pharmacologic prophylaxis **"And" Linked Panel**
AND mechanical prophylaxis

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel

<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)		
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/>	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/>	patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/>	patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/>	fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/>	heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/>	heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/>	warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/>	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)		
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE		
<input type="checkbox"/> High Risk (Selection Required)		
<input type="checkbox"/>	High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)		
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)		
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/>	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min

<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
[] High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
[] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
[] High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
[] High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min.

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

URL: "\appt1.pdf"

<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	
Low Risk Definition Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/> Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Surgical (Selection Required)	

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

Moderate risk of VTE Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following
contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following
contraindication(s):
PACU & Post-op

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following
contraindication(s):
PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 0600, Starting S+1

patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 0600, Starting S+1
For Patients with CrCL LESS than 30 mL/min

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min 40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.
This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition	
Both pharmacologic AND mechanical prophylaxis must be addressed.	
One or more of the following medical conditions:	
Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)	
Severe fracture of hip, pelvis or leg	
Acute spinal cord injury with paresis	
Multiple major traumas	
Abdominal or pelvic surgery for CANCER	
Acute ischemic stroke	
History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

Labs

Conn's/Hyperaldosteronism Labs - AM

<input type="checkbox"/> Potassium level	AM draw, Starting S+1 For 1 Occurrences, Post-op
<input type="checkbox"/> Renin activity	AM draw, Starting S+1 For 1 Occurrences, Post-op
<input type="checkbox"/> Aldosterone, serum	AM draw, Starting S+1 For 1 Occurrences, Post-op

Cushing's/Hypercortisolism Labs - AM

<input type="checkbox"/> Cortisol level, AM	Once, Starting S+1 at 8:00 AM Draw level at 0800
---	---

Imaging

X-ray

<input type="checkbox"/> XR Chest 1 Vw Portable	Routine, 1 time imaging For 1
---	-------------------------------