

Heparin Induced Thrombocytopenia (HIT) [4118]

IMMEDIATELY DISCONTINUE ALL HEPARIN CONTAINING PRODUCTS IN ANY FORM INCLUDING LOW-MOLECULAR WEIGHT HEPARINS (ENOXAPARIN), HEPARIN FLUSHES, AND HEPARIN COATED CATHETERS

General

HIT 4T's Score and No Anticoagulants including Heparins Order

<input checked="" type="checkbox"/> HIT 4T Score	Thrombocytopenia: Evidence of active thrombosis: Days of heparin treatment when platelet count began to fall: Other causes of thrombocytopenia:
<input checked="" type="checkbox"/> No heparin products	Routine, Until discontinued, Starting S Reason for "No" order: Suspected heparin induced thrombocytopenia

Labs

Heparin PF4 Antibody

<input checked="" type="checkbox"/> Heparin PF4 antibody (IgG)	Once
--	------

Consults

For Physician Consult orders use sidebar

Pharmacy and Hematology Consults

<input type="checkbox"/> Pharmacy consult to manage bivalirudin (ANGIOMAX) for Heparin Induced Thrombocytopenia	Routine, Until discontinued, Starting S
<input type="checkbox"/> Pharmacy consult to manage argatroban for Heparin Induced Thrombocytopenia	Routine, Until discontinued, Starting S
<input type="checkbox"/> Consult Hematology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?