## General Adult Total Parenteral Nutrition [976] ADULT TOTAL PARENTERAL NUTRITION

abs	
abs Once	
Comprehensive metabolic panel	Once For 1 Occurrences
] Magnesium	Once For 1 Occurrences
	prior to starting parenteral nutrition
Phosphorus  Prealbumin level	Once For 1 Occurrences
	prior to starting parenteral nutrition
	Once For 1 Occurrences
	prior to starting parenteral nutrition
Triglycerides	Once For 1 Occurrences
bs Repeat - All Facilities Except HMSJ	
Comprehensive Metabolic Panel	AM draw repeats, Starting S+7 For 2 Occurrences
Phosphorus	AM draw repeats, Starting S+7 For 2 Occurrences
Magnesium level	AM draw repeats, Starting S+7 For 2 Occurrences
Prealbumin	AM draw repeats, Starting S+2 For 2 Occurrences
Triglycerides	AM draw repeats, Starting S+2 For 2 Occurrences
abs Repeat - HMSJ Only	
Comprehensive Metabolic Panel	AM draw repeats, Starting S+1 For 3 Occurrences
Phosphorus	AM draw repeats, Starting S+1 For 3 Occurrences
Magnesium level	AM draw repeats, Starting S+1 For 3 Occurrences
Prealbumin	Every Monday
Triglycerides	Every Monday
Hepatic function panel	Every Monday
riopado iditodori parior	Every Menday
utritional Services Consult	
PN	equired)
PN harmacy Consults (Single Response) (Selection Re	equired) STAT, Until discontinued, Starting S
PN harmacy Consults (Single Response) (Selection Re	
PN narmacy Consults (Single Response) (Selection Re	STAT, Until discontinued, Starting S
PN narmacy Consults (Single Response) (Selection Re	STAT, Until discontinued, Starting S Enteral Nutrition:
PN narmacy Consults (Single Response) (Selection Re Pharmacy consult to manage TPN therapy	STAT, Until discontinued, Starting S Enteral Nutrition: Indication:
PN narmacy Consults (Single Response) (Selection Re Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences
PN narmacy Consults (Single Response) (Selection Re Pharmacy consult to manage TPN therapy	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition:
PN narmacy Consults (Single Response) (Selection Re Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication:
Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access:
PN narmacy Consults (Single Response) (Selection Re Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted:
Pharmacy Consults (Single Response) (Selection Response) (Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy:
Pharmacy Consults (Single Response) (Selection Resp	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number:
Pharmacy Consults (Single Response) (Selection Response) (Selection Response) (Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy:
) Pharmacy consult to monitor and provide	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number: STAT, Once For 1 Occurrences
harmacy Consults (Single Response) (Selection Response)  Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy  No pharmacy TPN consult needed	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number: STAT, Once For 1 Occurrences
harmacy Consults (Single Response) (Selection Response)  Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy  No pharmacy TPN consult needed  harmacy Consults (Single Response) (Selection Response)	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number: STAT, Once For 1 Occurrences  equired)  STAT, Until discontinued, Starting S Enteral Nutrition:
harmacy Consults (Single Response) (Selection Response)  Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy  No pharmacy TPN consult needed  harmacy Consults (Single Response) (Selection Response)	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number: STAT, Once For 1 Occurrences  equired)  STAT, Until discontinued, Starting S Enteral Nutrition: Indication:
PN harmacy Consults (Single Response) (Selection Response) Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy  No pharmacy TPN consult needed harmacy Consults (Single Response) (Selection Response)	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number: STAT, Once For 1 Occurrences  equired)  STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access:
PN harmacy Consults (Single Response) (Selection Response) Pharmacy consult to manage TPN therapy  Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy  No pharmacy TPN consult needed harmacy Consults (Single Response) (Selection Response)	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number: STAT, Once For 1 Occurrences  equired)  STAT, Until discontinued, Starting S Enteral Nutrition: Indication:

( ) Pharmacy consult to monitor and provide	STAT, Once For 1 Occurrences
recommendation ONLY for TPN therapy	Enteral Nutrition:
	Indication:
	Location of venous access:
	Is patient volume restricted:
	Instructions to pharmacy: Physician contact number:
	Physician contact number.
lutrition Services Consult	
[X] Consult to Nutrition Services	Reason For Consult? Other (Specify)
	Specify: TPN
	Purpose/Topic: Nutrition Services to work with Pharmacy on all TPN orders
Premixed TPN solutions (CLINIMIX) - HMSL Only	
X] TPN Solutions - HMSL Only (Single Response)	
( ) amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
() amino acids 4.25%-dextrose 25%- lytes	intravenous, continuous TPN, S at 9:00 PM
(CLINIMIX E) with MVI and trace () amino acids 5%-dextrose 15%- lytes	intravenous, continuous TPN, S at 9:00 PM
(CLINIMIX E) with MVI and trace () amino acids 5%-dextrose 20%- lytes	intravenous, continuous TPN, S at 9:00 PM
(CLINIMIX E) with MVI and trace	
( ) amino acids 5%-dextrose 25%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
() fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
Premixed TPN solutions (CLINIMIX) - HMW Only	
X] TPN Solutions - HMW Only (Single Response)	
( ) amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
( ) amino acids 5%-dextrose 15%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
( ) amino acids 5%-dextrose 20%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
( ) fat emulsion-MCT-olive-soy-fish	250 mL, intravenous, 3 times weekly at 2100
(SMOFlipid) 20% infusion	
remixed TPN solutions (CLINIMIX) HMSJ Only	
Clinimix Premix TPN Solutions (Single Response)	
( ) amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
( ) amino acids 5%-dextrose 20%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM
( ) amino acid 5 %-dextrose 20 % (CLINIMIX) 2,000 mL with multivitamin 10 mL,	intravenous, nightly
multitrace elements-5 concentrate (MTE-5	
( Oncentrate) 1 mi	
Concentrate) 1 mL  Lipids (Single Response)	

() fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
Premixed TPN solutions (CLINIMIX) HMSTC	
[X] Adult Custom Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM
Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
() fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
() soybean oil (INTRAlipid) 20% infusion	intravenous, nightly
TPN (HMSTJ Only)	
[X] TPN Solutions (Single Response)	
( ) Adult Standard Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM hang D10 if TPN is interrupted
( ) Adult Custom Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM hang D10 if TPN is interrupted
( ) Adult Custom Central TPN	intravenous, for 24 Hours, continuous Indication:
() Adult PERIPHERAL TPN	intravenous, for 24 Hours, continuous Indication:
() Adult CYCLIC CENTRAL TPN	intravenous, cyclic TPN - see admin instructions, S at 9:00 PM Indication:
() Adult CYCLIC PERIPHERAL TPN	intravenous, cyclic TPN - see admin instructions, S at 9:00 PM Indication:
[] Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID)  20% infusion	250 mL, intravenous, 3 times weekly at 2100
() fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
TPN (HMW Only)	
[X] TPN Solutions (Single Response)	
() Adult Standard Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM
( ) Adult Custom Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM
( ) Adult Custom Central TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM Indication:
[] Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
( ) fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
Adult TPN - NOT HMSTJ, HMTW, HMSTC	
[X] Adult TPN (Single Response)	
() Adult Central TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM Indication:
() Adult Peripheral TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM Indication:
( ) Adult CYCLIC Peripheral TPN	intravenous, cyclic TPN - see admin instructions, S at 9:00 PM Indication:
( ) Adult CYCLIC Central TPN	intravenous, cyclic TPN - see admin instructions, S at 9:00 PM Indication:
[] Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
() fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100

## **Adult TPN - HMTW Only (Single Response)**

/ \	
( ) Adult Central TPN + Fat Emulsion [ ] Adult CENTRAL TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM
[] Addit OLIVITAL II IV	Indication:
[] Lipids (Single Response)	
<ul><li>( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion</li></ul>	250 mL, intravenous, 3 times weekly at 2100
( ) fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
( ) Adult Peripheral TPN + Fat Emulsion	
[] Adult PERIPHERAL TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM Indication:
[] Lipids (Single Response)	
( ) fat emulsion-olive-soy-lipid (CLINOLIPID) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
( ) fat emulsion-MCT-olive-soy-fish (SMOFlipid) 20% infusion	250 mL, intravenous, 3 times weekly at 2100
( ) Adult Central TPN WITHOUT Fat Emulsion	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM Indication:
( ) Adult Peripheral TPN WITHOUT Fat Emulsion	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM Indication:
dextrose 10% for Disruption of TPN Therapy Start 10% dextrose IV during any interruption in T	FPN at previous TPN rate up to a MAXIMUM rate of 40 mL/hr.
[X] dextrose 10 % infusion	40 mL/hr, intravenous, continuous PRN, other, For disruption of TPN therapy.
	Start 10% dextrose IV prior to initiation or during any