

# EPOprostenol (Flolan) (RESTRICTED)[1173]

A calculator is available at [formweb.com/houstonmethodist](http://formweb.com/houstonmethodist) under Drug Information and Clinical Calculators to assist with epoprostenol or treprostinil calculations. To navigate to this site, please open an internet browser and use the web address noted previously.

## Nursing

### Nursing

<input type="checkbox"/> EPOprostenol (Flolan) instructions for Nursing:	Routine, Until discontinued, Starting S 1. Do NOT infuse any other IV fluids or medications through Flolan line. 2. Do NOT draw blood from a lumen adjacent to Flolan line. 3. Do NOT STOP Flolan for ANY REASON without first contacting MD who ordered Flolan 4. DO NOT STOP Flolan for CODE without first contacting MD who ordered Flolan. Document any infusion rate changes at appropriate times in the MAR
--	--

## IV Fluids

## Medications

### Medications (Single Response) (Selection Required)

#### ( ) Continuation of Therapy

<input type="checkbox"/> Vital Signs	<b>"And" Linked Panel</b>
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol Upon initiation of therapy or dose increase, every 15 minutes x 4, then every 30 minutes x2, then every 4 hours.
<input type="checkbox"/> EPOprostenol (Flolan) straight dose	
<input type="checkbox"/> epoprostenol (FLOLAN) in diluent for epoprostenol (gly) 100 mL infusion (RESTRICTED)	intravenous, for 24 Hours, daily at 1400 DO NOT STOP Flolan for ANY REASON without first contacting MD who ordered Flolan. For CENTRAL LINE administration Infusion Pump Rate (mL per 24 hours): Dosing Weight: (in kg): Dose: ( in ng/kg/min):
<input type="checkbox"/> Pharmacy Consult	
<input type="checkbox"/> Pharmacy consult to review pulmonary arterial hypertension (PAH) infusions	STAT, Until discontinued, Starting S

#### ( ) Therapy Titration

<input type="checkbox"/> Vital Signs	<b>"And" Linked Panel</b>
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol Upon initiation of therapy or dose increase, every 15 minutes x 4, then every 30 minutes x2, then every 4 hours.
<input type="checkbox"/> EPOprostenol (Flolan) straight dose	
<input type="checkbox"/> epoprostenol (FLOLAN) in diluent for epoprostenol (gly) 100 mL infusion (RESTRICTED)	intravenous, for 24 Hours, daily at 1400, Starting S DO NOT STOP Flolan for ANY REASON without first contacting MD who ordered Flolan. For CENTRAL LINE administration  Titration Instructions: Initiate epoprostenol (Flolan) as ordered. {Increase/Decrease:34622} by {.:34620} mL/24hours every {.:34621} hours {Maximum/Minimum:34623} infusion rate of ***mL/24hours Infusion Pump Rate (mL per 24 hours): Dosing Weight: (in kg): Dose: ( in ng/kg/min):
<input type="checkbox"/> Pharmacy Consult	
<input type="checkbox"/> Pharmacy consult to review pulmonary arterial hypertension (PAH) infusions	STAT, Until discontinued, Starting S

