

# Amphotericin B Conventional (Fungizone) Infusion (RESTRICTED) [2390]

!!! CONVENTIONAL formulation !!!

Amphotericin B conventional (FUNGIZONE) infusion is RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists.

## Medications

### Amphotericin B Conventional (FUNGIZONE) (Single Response) (Selection Required)

#### ( ) Amphotericin B Conventional (FUNGIZONE) - Peripheral "And" Linked Panel Line

<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+2 Hours Give 1 hour BEFORE amphotericin
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection	25 mg, intravenous, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection	25 mg, intravenous, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+3 Hours Flush line immediately BEFORE amphotericin.
<input type="checkbox"/> amphotericin B (FUNGIZONE) in dextrose 5% 500 mL IVPB (RESTRICTED)	intravenous, for 4 Hours, every 24 hours, Starting H+3 Hours RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists. Are you an ID, SOT, BMT, or Heme/Onc specialist or ordering on behalf of one? Reason for Therapy:
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+7 Hours Flush line immediately AFTER amphotericin.
<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+7 Hours Give AFTER amphotericin and dextrose 5% flush.

#### ( ) Amphotericin B Conventional (FUNGIZONE) - Central "And" Linked Panel Line

<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+2 Hours Give 1 hour BEFORE amphotericin
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection	25 mg, intravenous, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection	25 mg, intravenous, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+3 Hours Flush line immediately BEFORE amphotericin.
<input type="checkbox"/> amphotericin B (FUNGIZONE) in dextrose 5% 250 mL IVPB (RESTRICTED)	intravenous, for 4 Hours, every 24 hours, Starting H+3 Hours RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists. Are you an ID, SOT, BMT, or Heme/Onc specialist or ordering on behalf of one? Reason for Therapy:
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+7 Hours Flush line immediately AFTER amphotericin.
<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+7 Hours Give AFTER amphotericin and dextrose 5% flush.

## Labs

### Laboratory

<input type="checkbox"/> Basic metabolic panel - STAT	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	Once
<input type="checkbox"/> Magnesium level	Once

[ ] Phosphorus level

Once