

Heparin Infusion Standard Protocol (DVT/PE) WITH Bolus [1768]

General

Nursing

Nursing

[X] Weigh patient	Routine, Once
[X] Notify Pharmacist	Routine, Until discontinued, Starting S For Until specified, NOTIFY pharmacist if patient CANNOT be weighed.
[X] Notify pharmacist	Routine, Until discontinued, Starting S, Contact pharmacist if a physician writes heparin orders. Heparin dose or infusion changes to be made by pharmacist only. Notify Pharmacist immediately if the patient transfers or has an off unit procedure.
[X] Heparin instructions	Routine, Until discontinued, Starting S Do not interrupt heparin infusion unless ordered. Contact pharmacist regarding compatibility with other IV drugs if access is a concern. Contact pharmacist immediately if heparin infusion is stopped for any reason.
[X] Notify Physician and Pharmacist	Routine, Until discontinued, Starting S, Platelet count less than 150,000 microliter or a 50% drop from baseline or hematoma, bleeding or suspected bleeding occurs.
[X] Do not call PTT/Anti Xa UFH results to physician unless ordered.	Routine, Until discontinued, Starting S For Until specified Do not call PTT/Anti Xa UFH results to physician unless ordered.
[X] Do not draw blood from the arm that has heparin infusion or that has been flushed with heparin.	Routine, Until discontinued, Starting S If there is no other access , other than the heparin line for an Anti Xa unfractionated specimen draw, then stop the heparin, flush the line, and aspirate 10 ml of blood to waste prior to drawing a specimen.

IV Fluids

Medications

Medications

[X] HEParin (porcine) injection	80 Units/kg, intravenous, once, For 1 Doses
[X] HEParin 25,000 unit/500 mL (50 unit/mL)	intravenous, titrated Indication: Therapeutic Monitoring Target:

VTE

Labs

Lab - Initiation

[X] Prothrombin time with INR	Once
[X] Partial thromboplastin time	Once
[X] CBC hemogram	Once
[X] Anti Xa, unfractionated	Once Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines. If there is no other access other than the heparin line for an Anti Xa unfractionated specimen draw, stop the heparin, flush the line, and aspirate 10 ml of blood to waste.

Lab - Continuing

<input type="checkbox"/> CBC hemogram	Now then every 24 hours, Starting S+1 at 6:00 AM For 3 Occurrences
<input type="checkbox"/> Anti Xa, unfractionated	Once Six hours after initiation of heparin infusion. Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines. If there is no other access other than the heparin line for an Anti Xa unfractionated specimen draw, stop the heparin, flush the line, and aspirate 10 ml of blood to waste.
<input type="checkbox"/> Partial thromboplastin time	Once For Until specified Six hours after initiation of heparin infusion. Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines. If there is no other access, other than the heparin line for a Partial thromboplastin time (PTT) specimen draw, then stop the heparin, flush the line, and aspirate 10 ml of blood to waste prior to drawing a specimen.
<input checked="" type="checkbox"/> Occult blood, stool	Conditional Frequency, Stool guicac stools when there is a hemoglobin decrease of more than 2 gm/dL in a 24 hour period

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders