

CVIR Post PICC Line Insertion [694]

General

Nursing

Nursing

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 2, PACU & Post-op
<input type="checkbox"/> Ok to use	Routine, Until discontinued, Starting S Device: PICC line PICC trim length to *** centimeters, PACU & Post-op
<input type="checkbox"/> Needleless Connector	Routine, Every Tue, Fri Change every Tuesday and Friday and as needed. Use positive flow needleless valve, caps only (NO EXCEPTIONS), Post-op
<input type="checkbox"/> Change dressing	Routine, Once And as needed., PACU & Post-op
<input type="checkbox"/> Use 10 milliliter syringe or greater (NO EXCEPTIONS)	Routine, Until discontinued, Starting S Use 10 milliliter syringe or greater (NO EXCEPTIONS), PACU & Post-op
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S Instructions: ***, PACU & Post-op
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S Instructions: ***, PACU & Post-op

IV Fluids

Return to Floor

Return to Floor (Single Response)

<input type="checkbox"/> OK to return to previous IP bed	Routine, Until discontinued, Starting S, PACU
--	---

Medications

Post Procedure Medications

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 15 min PRN, severe pain (score 7-10), persistent pain, For 2 Doses, PACU & Post-op Hold for respirations LESS than 12 breath/minute. For persistent pain, call interventional MD.
<input type="checkbox"/> ondansetron (ZOFTRAN) IV	4 mg, intravenous, once PRN, nausea, vomiting, PACU & Post-op

Discharge

Discharge Order (Single Response)

<input type="checkbox"/> Discharge patient when criteria met	Routine, Once Discharge Criteria: Clearing specialty: Scheduling/ADT
<input type="checkbox"/> Discharge to home	Routine, Once Discharge Criteria: Clearing specialty: Scheduling/ADT

VTE

Labs

Cardiology

Imaging

Other Diagnostic Studies

Respiratory

Rehab

Consults

Additional Orders