

Pre-Op Eye Surgery [716]

General

Planned ICU Admission Post-Operatively (Admit to Inpatient Order) (Single Response)

Patients who are having an Inpatient Only Procedure as determined by CMS and patients with prior authorization for Inpatient Care may have an Admit to Inpatient order written pre-operatively.

Admit to Inpatient

Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
Pre-op

Case Request

Case request operating room

Scheduling/ADT, Scheduling/ADT

Nursing

IV Fluids

Medications

Ophthalmic Medications

<input type="checkbox"/> lidocaine (XYLOCAINE) 20 mg/mL (2 %) injection	5 mL, injection, once, For 1 Doses, Pre-op To Operative eye to prepare for Ophthalmic Block
<input type="checkbox"/> flurbiprofen (OCUFEN) 0.03 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution	1 drop, every 5 min, For 4 Doses, Pre-op
<input type="checkbox"/> cyclopentolate (CYCLODRYL) 1 % ophthalmic solution	1 drop, every 5 min, For 4 Doses, Pre-op
<input type="checkbox"/> keTOROLac (ACULAR LS) 0.4 % ophthalmic solution	1 drop, every 5 min, For 4 Doses, Pre-op (NOTE THIS IS A DIFFERENT PERCENTAGE; ALSO CHANGE THE OTHER 0.5% TO 3 DOSES AS WELL)
<input type="checkbox"/> ketorolac (ACULAR) 0.5 % ophthalmic solution	1 drop, every 5 min, For 4 Doses, Pre-op
<input type="checkbox"/> pilocarpine (PILOCAR) 2 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> pilocarpine (PILOCAR) 1 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> tetracaine HCl (PF) 0.5 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op

Ophthalmic Antibiotics

<input type="checkbox"/> besifloxacin (BESIVANCE) 0.6% ophthalmic suspension	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> moxifloxacin (VIGAMOX) 0.5% ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> gentamicin (GARAMYCIN) 0.3% ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op

Pupillary Dilation

<input type="checkbox"/> atropine 1 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> phenylephrine (MYDFRIN) 2.5 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> cyclopentolate (CYCLOGYL) 1 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> homatropine 5 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
<input type="checkbox"/> tropicamide (MYDRIACYL) 1 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op

VTE

Labs

COVID-19 Qualitative PCR

[] COVID-19 qualitative RT-PCR - Nasal Swab

STAT For 1 Occurrences
Specimen Source: Nasal Swab
Is this for pre-procedure or non-PUI assessment? Yes
Pre-op

Cardiology

Diagnostic Imaging

Other Diagnostic Studies

Respiratory

Rehab

Consults

Additional Orders