General

Pre Anesthesia Testing Orders

The ambulatory orders in this section are specifically for Pre Anesthesia Testing. For additional PAT orders, please use 'Future Status' and 'Pre-Admission Testing Phase of Care'

Other Diagnostic Studies	
[] ECG Pre/Post Op	Routine, Ancillary Performed, Status: Future, Expires: S+365, Pre-Admission Testing
[] Pv carotid duplex	Status: Future, Expires: S+365, Routine, Clinic Performed
[] Us vein mapping lower extremity	Status: Future, Expires: S+365, Routine, Clinic Performed
Us duplex venous lower extremity	Status: Future, Expires: S+365, Routine, Clinic Performed
Respiratory	
[] Spirometry pre & post w/ bronchodilator	Routine, Status: Future, Expires: S+365, Pre-Admission Testing Encounter type?
[] Body Plethysmographic lung volumes	Routine, Status: Future, Expires: S+365, Pre-Admission Testing Encounter type?
[] Diffusion capacity	Routine, Status: Future, Expires: S+365, Pre-Admission Testing Encounter type?
[] Spirometry	Routine, Status: Future, Expires: S+365, Pre-Admission Testing Encounter type?
Laboratory: Preoperative Testing Labs - All Fa	
[] COVID-19 qualitative RT-PCR - Nasal	Routine, Status: Future, Expires: S+365, Clinic Collect
Swab	Specimen Source: Nasal Swab
	Is this for pre-procedure or non-PUI assessment? Yes
	Pre-Admission Testing
[] CBC with platelet and differential	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission
	Testing
[] Comprehensive metabolic panel	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Basic metabolic panel	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Prothrombin time with INR	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Partial thromboplastin time	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Hepatic function panel	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Platelet function analysis	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Hemoglobin A1c	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Type and screen	
[] Type and screen	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] ABO and Rh confirmation	Once, Blood Bank Confirmation
[] hCG qualitative, serum screen	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] POC pregnancy, urine	Routine, Point Of Care, Pre-Admission Testing
[] Urinalysis, automated with microscopy	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
Laboratory: Additional Labs - HMWB, HMCL,	

[] Laboratory: Additional Labs - HMWB, HMCL, HMTW

[]	Urinalysis screen and microscopy, with reflex to culture	Routine, Status: Future, Expires: S+365, Clinic Collect Specimen Source: Urine Specimen Site: Pre-Admission Testing
[]	CBC hemogram	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	HIV 1, 2 antibody	Routine, Status: Future, Expires: S+365, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
[]	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Routine, Status: Future, Expires: S+365, Clinic Collect Release to patient (Note: If manual release option is selected, result will
[]	Hepatitis acute panel	auto release 10 days from finalization.): Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Thromboelastograph	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Vitamin D 25 hydroxy level	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	MRSA PCR	Routine, Status: Future, Expires: S+365, Clinic Collect, Nares, Pre-Admission Testing
[]	Т3	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]		Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Thyroid stimulating hormone	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Prostate specific antigen	Routine, Status: Future, Expires: S+365, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
L	aboratory: Additional Labs - HMSL, HMW	
[]	Urinalysis screen and microscopy, with reflex to culture	Routine, Status: Future, Expires: S+365, Clinic Collect Specimen Source: Urine Specimen Site: Pre-Admission Testing
[]	CBC hemogram	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Rapid HIV 1 & 2	Routine, Status: Future, Expires: S+365, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
[]	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Routine, Status: Future, Expires: S+365, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.):
[]	Hepatitis acute panel	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Thromboelastograph	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	Vitamin D 25 hydroxy level	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[]	MRSA PCR	Routine, Status: Future, Expires: S+365, Clinic Collect, Nares, Pre-Admission Testing
	T3	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
ITO		

Collect, Pre-Admission
Collect, Pre-Admission
Collect ion is selected, result will
Collect
Collect, Pre-Admission
Collect ion is selected, result will
Collect
ion is selected, result will
Collect, Pre-Admission
Collect, Pre-Admission
Collect, Pre-Admission
Collect, Nares,
Collect, Pre-Admission
Collect, Pre-Admission
Collect, Pre-Admission
Collect ion is selected, result will
Collect e?
Collect, Pre-Admission

[] Partial thromboplastin time, activated	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Vitamin A level, plasma or serum	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Vitamin B12 level	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Vitamin D 25 hydroxy level	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Copper level, serum	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Folate level	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Vitamin B1 level, whole blood	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
[] Zinc level, serum	Routine, Status: Future, Expires: S+365, Clinic Collect, Pre-Admission Testing
) Admit to Inpatient	Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgmen and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. Pre-op
ase Request	
Case request operating room	Scheduling/ADT, Scheduling/ADT
lursing	
ursing	
(] Vital Signs	Routine, Per unit protocol Temperature; Pulse; Respiration; Blood Pressure, Pre-op
[] Confirm NPO status	Routine, Until discontinued, Starting S Confirm NPO status, Pre-op
<] Height and weight	Routine, Once For 1 Occurrences, Pre-op
 Place/Maintain sequential compression devic continuous 	Routine, Once For 1 Occurrences, Pre-op
K] Complete Consent For	Routine, Once

[X] Notify Surgeon	Routine, Until discontinued, Starting S Temperature greater than: 100.5
	Temperature less than:
	Systolic BP greater than: 160
	Systolic BP less than: 90
	Diastolic BP greater than: 100
	Diastolic BP less than: 50
	MAP less than: 60
	Heart rate greater than (BPM): 100
	Heart rate less than (BPM): 60
	Respiratory rate greater than: 25
	Respiratory rate less than: 8 SpO2 less than: 92
Diet	
[X] NPO	Diet effective now, Starting S
	NPO:
	Pre-Operative fasting options:
[] Diet	Pre-op Dist offective new Storting S
[] Diet	Diet effective now, Starting S Diet(s):
	Other Options:
	Advance Diet as Tolerated?
	IDDSI Liquid Consistency:
	Fluid Restriction:
	Foods to Avoid:
	Foods to Avoid:
	Pre-op
IV Fluids	
IV Fluids	
[] sodium chloride 0.9 % infusion	intravenous, continuous, Pre-op
Medications	
Ophthalmic Medications	
·	
I I KEIUKUIAC (ACULAK LS) U.4 % OPNTNAIMIC SOLUTION	1 drop, every 5 min, For 3 Doses, Pre-op
	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
[] keTOROlac (ACULAR) 0.5 % ophthalmic solution	1 drop, every 5 min, For 3 Doses, Pre-op
[] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution	
 [] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution [] tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic 	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 [] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution [] tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution 	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 [] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution [] tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution [] pilocarpine (PILOCAR) 1 % ophthalmic solution 	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 [] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution [] tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution [] pilocarpine (PILOCAR) 1 % ophthalmic solution [] pilocarpine (PILOCAR) 2 % ophthalmic solution 	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 [] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution [] tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution [] pilocarpine (PILOCAR) 1 % ophthalmic solution [] pilocarpine (PILOCAR) 2 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution 	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 keTOROlac (ACULAR) 0.5 % ophthalmic solution flurbiprofen (OCUFEN) 0.03% ophthalmic solution tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution pilocarpine (PILOCAR) 1 % ophthalmic solution pilocarpine (PILOCAR) 2 % ophthalmic solution Ophthalmic Antibiotics besifloxacin (BESIVANCE) 0.6% ophthalmic solution moxifloxacin (VIGAMOX) 0.5% ophthalmic solution 	 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 keTOROlac (ACULAR) 0.5 % ophthalmic solution flurbiprofen (OCUFEN) 0.03% ophthalmic solution tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution pilocarpine (PILOCAR) 1 % ophthalmic solution pilocarpine (PILOCAR) 2 % ophthalmic solution Ophthalmic Antibiotics besifloxacin (BESIVANCE) 0.6% ophthalmic solution moxifloxacin (VIGAMOX) 0.5% ophthalmic solution 	1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 keTOROlac (ACULAR) 0.5 % ophthalmic solution flurbiprofen (OCUFEN) 0.03% ophthalmic solution tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution pilocarpine (PILOCAR) 1 % ophthalmic solution pilocarpine (PILOCAR) 2 % ophthalmic solution pilocarpine (BESIVANCE) 0.6% ophthalmic suspension moxifloxacin (VIGAMOX) 0.5% ophthalmic solution gentamicin (GARAMYCIN) 0.3% ophthalmic solution 	 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 [] keTOROlac (ACULAR) 0.5 % ophthalmic solution [] flurbiprofen (OCUFEN) 0.03% ophthalmic solution [] tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution [] phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution [] pilocarpine (PILOCAR) 1 % ophthalmic solution [] pilocarpine (PILOCAR) 2 % ophthalmic solution Ophthalmic Antibiotics [] besifloxacin (BESIVANCE) 0.6% ophthalmic solution [] moxifloxacin (VIGAMOX) 0.5% ophthalmic solution [] gentamicin (GARAMYCIN) 0.3% ophthalmic solution [] atropine 1 % ophthalmic solution 	 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 keTOROlac (ACULAR) 0.5 % ophthalmic solution flurbiprofen (OCUFEN) 0.03% ophthalmic solution tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution pilocarpine (PILOCAR) 1 % ophthalmic solution pilocarpine (PILOCAR) 2 % ophthalmic solution Ophthalmic Antibiotics besifloxacin (BESIVANCE) 0.6% ophthalmic solution gentamicin (GARAMYCIN) 0.3% ophthalmic solution Pupillary Dilation atropine 1 % ophthalmic solution phenylephrine (MYDFRIN) 2.5 % ophthalmic solution 	 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 keTOROlac (ACULAR) 0.5 % ophthalmic solution flurbiprofen (OCUFEN) 0.03% ophthalmic solution tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution pilocarpine (PILOCAR) 1 % ophthalmic solution pilocarpine (PILOCAR) 2 % ophthalmic solution pohthalmic Antibiotics besifloxacin (BESIVANCE) 0.6% ophthalmic solution gentamicin (GARAMYCIN) 0.3% ophthalmic solution gentamicin 1 % ophthalmic solution phenylephrine (MYDFRIN) 2.5 % ophthalmic solution cyclopentolate (CYCLODRYL) 1 % ophthalmic solution 	 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op
 keTOROlac (ACULAR) 0.5 % ophthalmic solution flurbiprofen (OCUFEN) 0.03% ophthalmic solution tetracaine HCI (PF) (AKTEN) 0.5 % ophthalmic solution phenylephrine (NEO-SYNEPHRINE) 10 % ophthalmic solution pilocarpine (PILOCAR) 1 % ophthalmic solution pilocarpine (PILOCAR) 2 % ophthalmic solution Ophthalmic Antibiotics besifloxacin (BESIVANCE) 0.6% ophthalmic solution gentamicin (GARAMYCIN) 0.3% ophthalmic solution Pupillary Dilation atropine 1 % ophthalmic solution phenylephrine (MYDFRIN) 2.5 % ophthalmic solution 	 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op 1 drop, every 5 min, For 3 Doses, Pre-op

[] cefazolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy:
[] cefoxitin (MEFOXIN) IV	2 g, intravenous, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy:
IV Antibiotics: For Patients GREATER than 120 kg	
[] cefazolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy:
[] cefoxitin (MEFOXIN) IV	2 g, intravenous, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy:
Antibiotics: if Beta-Lactam Allergy	
[] levofloxacin (LEVAQUIN) IV solution	500 mg, intravenous, on call to O.R., Pre-op On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy:
Antibiotics MRSA Suspected Do not redose post op for patient with CrCI less than 40	
[] vancomycin (VANCOCIN) IV	 15 mg/kg, intravenous, once, For 1 Doses, Pre-op On call to the operating room. Administer one hour prior to the opening incision. Rate should not exceed 1000 mg per hour. Do not exceed MAX dose 2000 mg for surgical prophylaxis. Type of Therapy:
Respiratory	
[] albuterol (VENTOLIN) inhaler	2 puff, inhalation, once, For 1 Doses, Pre-op
] ipratropium (ATROVENT) nebulizer	0.5 mg, nebulization, once, For 1 Doses, Pre-op
	Aerosol Delivery Device:
Blocks	Aerosol Delivery Device:
BIOCKS Houston Methodist Physician Group (Single Response)	Aerosol Delivery Device:
	once, Pre-op Needle: Atkinson
Houston Methodist Physician Group (Single Response)	once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Atkinson 23G
Houston Methodist Physician Group (Single Response) () Dr Joanna Queen	once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op
Houston Methodist Physician Group (Single Response) () Dr Joanna Queen () Dr Rahul Pandit	once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Atkinson 23G Syringe: 10 cc once, Pre-op Needle: Atkinson

() Dr Mattew Benz	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10 cc
) Dr David Brown	once, Pre-op
	Needle: $25G \times 11/2$
	Syringe: 5-6 cc
() Dr Eric Chen	once, Pre-op
	Needle: $25G \times 11/2$
	Syringe: 6 cc
	If doing a Buckle under MAC - Make 2 syringes of the same
() Dr Diskard Fisk Detectively an	above
() Dr Richard Fish - Retrobulbar	once, Pre-op
	Needle: Atkinson
	Syringe: 5/6 cc
() Dr Richard Fish - Van Lente	once, Pre-op
	Needle: 25 x 1 1/2
	Syringe: 10/12 cc
() Dr Chris Henry	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10 cc
) Dr Rosa Kim	once, Pre-op
	Needle: Atkinson
	Syringe: 10/12 cc
) Dr James C Major	once, Pre-op
, ,	Needle: 25G x 1 1/2
	Syringe: 10 cc
) Dr Amy Schefler	once, Pre-op
	Needle: $27G \times 11/4$
	Syringe: 10 cc
() Dr Tien Wong	once, Pre-op
	Needle: $25G \times 11/2$
	Syringe: 10 cc
() Dr Charles Welest	
() Dr Charles Wykoff	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10 cc
Houston Eye Associates (Single Response)	
() Dr Emmanuel Chang	once, Pre-op
	Needle: Retrobulbar Atkinson
	Syringe: 10/12 cc
() Dr Carolyn Chen	once, Pre-op
· · ·	Needle: Atkinson
() Dr Ting Fang-Suarez - Retrobulbar	once, Pre-op
() = · · · · · · · · · · · · · · · · · ·	Needle: Atkinson
	Syringe: 5/6 cc
) Dr Ting Fang-Suarez - Van Lente	once, Pre-op
J Di Ting Tang-Odalez - Vali Leille	Needle: $25G \times 11/2$
	Syringe: 10/12 cc
() Dr Lindsey Harris	once, Pre-op
	Needle: Atkinson
	Syringe: 12 cc
	Synnge. 12 cc
	Make block for all patients, even general
() Dr Eric Holz	Make block for all patients, even general once, Pre-op
	Make block for all patients, even general

) Dr Andrew Kopel	once, Pre-op Needle: 25G x 1 1/2
	Syringe: 10/12 cc
) Dr. Asses Miller	
) Dr Aaron Miller	once, Pre-op
	Needle: Atkinson
	Syringe: 10 cc
) Dr Hassan Rahman	once, Pre-op
	Needle: Atkinson 23G
	Syringe: 10/12 cc
) Dr Mark Vital	once, Pre-op
,	Needle: Atkinson
	Syringe: 10/12 cc
ndependents (Single Response)	
) Dr John Alappatt	once, Pre-op
	Needle: Atkinson
	Syringe: 10cc
) Dr Wael Abdelghani	once, Pre-op
,	Needle: Atkinson
	Syringe: 10/12 cc
) Dr. Asad Abbas - Phacoemulsification with IOL	Gynnigo. 10/12 00
Implantation	
[] flurbiprofen (OCUFEN) 0.03 % ophthalmic	1 drop, see admin instructions, For 4 Doses, Pre-op
solution	Give 1 drop in the operative eye every 2 minutes x 4 doses
[] proparacaine (ALCAINE) 0.5 % ophthalmic	1 drop, see admin instructions, For 4 Doses, Pre-op
solution	Give 1 drop in the operative eye every 2 minutes x 4 doses
[] tropicamide (MYDRIACYL) 1 % ophthalmic	1 drop, see admin instructions, For 4 Doses, Pre-op
solution	
	Give 1 drop in the operative eye every 2 minutes x 4 doses
[] phenylephrine (NEO-SYNEPHRINE) 10 %	1 drop, see admin instructions, For 4 Doses, Pre-op
ophthalmic solution	Give 1 drop in the operative eye every 2 minutes x 4 doses
[] cyclopentolate (CYCLODRYL) 1 %	1 drop, see admin instructions, For 4 Doses, Pre-op
ophthalmic solution	Give 1 drop in the operative eye every 2 minutes x 4 doses
) Dr. Martin J. Arisco - Preoperative Cataract Surg	ery
[] proparacaine (ALCAINE) 0.5 % ophthalmic	1 drop, every 5 min, For 4 Doses, Pre-op
solution	
[] phenylephrine (NEO-SYNEPHRINE) 10 %	1 drop, every 5 min, For 4 Doses, Pre-op
ophthalmic solution	
[] cyclopentolate (CYCLODRYL) 1 %	1 drop, every 5 min, For 4 Doses, Pre-op
ophthalmic solution	
[] flurbiprofen (OCUFEN) 0.03 % ophthalmic	1 drop, every 5 min, For 4 Doses, Pre-op
solution	
[] lactated ringer's infusion	intravenous, continuous, Pre-op
	Start IV lactated ringers to run KVO, if time permits.
) Dr Milton Boniuk - Retrobulbar	
) Dr Milton Boniuk - Retrobulbar	once, Pre-op
	Needle: Atkinson
	Syringe: 5/6 cc
) Dr Milton Boniuk - Van Lente	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10/12 cc
) Dr Christopher Fuller	once, Pre-op
, I	Needle: 25G
	Syringe: 10 cc
) Dr. Judith Egigon Batrahulbar	
) Dr Judith Feigon - Retrobulbar	once, Pre-op
	Needle: Atkinson
	Syringe: 5/6 cc
) Dr Judith Feigon - Van Lente	once, Pre-op
) Dr Judith Feigon - Van Lente	once, Pre-op Needle: 25G x 1 1/2

() Dr Benjamin Frankfort	once, Pre-op
	Needle: Atkinson Syringe: 6 cc
() Dr Burt Ginsburg - Retrobulbar	once, Pre-op
() Di Buit Ginsburg - Netiobulbai	Needle: $25G \times 11/2$
	Syringe: 5/6 cc
() Dr Burt Ginsburg - Van Lente	once, Pre-op
	Needle: 27G x 1 1/4
	Syringe: 10/12 cc
() Dr Rex Hawkins - Retrobulbar	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 6 cc
() Dr Rex Hawkins - Van Lente	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10 cc
() Dr Helen Li	once, Pre-op
	Needle: Atkinson
() Dr. Thomas Marshall, Dreamarsting Cotorest Su	Syringe: 12 cc
() Dr. Thomas Marshall - Preoperative Cataract Sur	
[] proparacaine (ALCAINE) 0.5 % ophthalmic solution	1 drop, once, For 1 Doses, Pre-op
[] tropicamide (MYDRIACYL) 1 % ophthalmic	1 drop, every 10 min, For 3 Doses, Pre-op
solution	Give 1 drop in the operative eye every 10-15 minutes x 3 doses
[] phenylephrine (NEO-SYNEPHRINE) 10 %	1 drop, every 10 min, For 3 Doses, Pre-op
ophthalmic solution	Give 1 drop in the operative eye every 10-15 minutes x 3 doses
[] flurbiprofen (OCUFEN) 0.03 % ophthalmic	1 drop, every 10 min, For 3 Doses, Pre-op
solution	Give 1 drop in the operative eye every 10-15 minutes x 3 doses
[] lactated ringer's infusion	intravenous, continuous, Pre-op
	Start IV of 1000 mL lactated ringers with #22 Catheter to run KVO, if time
	permits.
() Dr. Carvounis Petros	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10/12 cc
() Dr Regina Sun	once, Pre-op
	Needle: Atkinson
() Dr Sanders	once, Pre-op
	Needle: Atkinson
	Syringe: 10 cc
Baylor (Single Response)	
() Dr Peter Chang	once, Pre-op
() Di Peter Chang	Needle: $27G \times 11/4$
	Syringe: 10/12 cc
() Dr Yvonne Chu	
	once. Pre-op
	once, Pre-op Needle: Atkinson
	Needle: Atkinson
() Dr Coffee	
() Dr Coffee	Needle: Atkinson Syringe: 6 cc
() Dr Coffee	Needle: Atkinson Syringe: 6 cc once, Pre-op
() Dr Coffee () Dr Marshall B Hamill - Retrobulbar	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson
	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson Syringe: 10 cc
	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op
	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Atkinson
() Dr Marshall B Hamill - Retrobulbar	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Atkinson Syringe: 5/6 cc
 () Dr Marshall B Hamill - Retrobulbar () Dr Marshall B Hamill - Van Lente 	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Atkinson Syringe: 5/6 cc once, Pre-op
() Dr Marshall B Hamill - Retrobulbar	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Atkinson Syringe: 5/6 cc once, Pre-op Needle: 25G x 1 1/2
 () Dr Marshall B Hamill - Retrobulbar () Dr Marshall B Hamill - Van Lente 	Needle: Atkinson Syringe: 6 cc once, Pre-op Needle: Needle: Atkinson Syringe: 10 cc once, Pre-op Needle: Needle: Atkinson Syringe: 5/6 cc once, Pre-op Needle: Needle: 25G x 1 1/2 Syringe: 10/12 cc

() no name - Van Lente	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10 cc
() Dr Douglas Koch - Retrobulbar Cataract	once, Pre-op
	Needle: Atkinson
	Syringe: 5/6 cc
	Check with Dr Koch if he needs marcaine
() Dr Douglas Koch - Van Lente	once, Pre-op
	Needle: 25G x 1 1/2
	Syringe: 10 cc
	Check with Dr Koch if he needs marcaine
() Dr Alice Matoba - Retrobulbar	once, Pre-op
	Needle: Atkinson
	Syringe: 5/6 cc
() Dr Alice Matoba - Van Lente	once, Pre-op
(Needle: 25G x 1 1/2
	Syringe: 10/12 cc
() Dr Stephen Pflugfelder - PKP and Pterygium	once, Pre-op
	Needle: 30G first, then Atkinson
	Syringe: 10/12 cc
() Dr Stephen Pflugfelder - Retrobulbar	once, Pre-op
	Needle: Atkinson
	Syringe: 10 cc
() Dr Mitchell Weikert - Cataracts & PKP	once, Pre-op
	Needle: Atkinson
	Syringe: 10 cc
	Synnge. 1000

VTE

Labs

COVID-19 Qualitative PCR

[] COVID-19 qualitative RT-PCR - Nasal Swab

STAT For 1 Occurrences Specimen Source: Nasal Swab Is this for pre-procedure or non-PUI assessment? Yes Pre-op

Laboratory

[] Basic metabolic panel	Once For 1 Occurrences, Pre-op
[] Comprehensive metabolic panel	Once For 1 Occurrences, Pre-op
[] CBC and differential	Once For 1 Occurrences, Pre-op
[] Partial thromboplastin time	Once For 1 Occurrences, Pre-op
[] Prothrombin time with INR	Once For 1 Occurrences, Pre-op
[] Calcium	Once For 1 Occurrences, Pre-op
[] hCG, serum, qualitative	Once For 1 Occurrences, Pre-op
[] Pregnancy, urine	Once For 1 Occurrences, Pre-op
[] Urinalysis with microscopic	Once For 1 Occurrences, Pre-op
[] Prothrombin time with INR, I-Stat	Once, Pre-op
[] Bedside glucose	Routine, Once, Pre-op

Cardiology

EKG

[X] ECG 12 lead

Routine, Once Clinical Indications: Pre-Op Clearance Interpreting Physician: Pre-op

Diagnostic Imaging

X-Ray

] Chest 1 Vw Portable	Routine, 1 time imaging, Starting S at	1:00 AM For 1, Pre-op
] Chest 2 Vw	Routine, 1 time imaging, Starting S at	1:00 AM For 1 , Pre-op

СТ

[] CT Orbits Wo Contrast

Routine, 1 time imaging, Starting S at 1:00 AM For 1 2 millimeter cuts with direct coronal if possible., Pre-op

Other Diagnostic Studies

Respiratory

Respiratory

[] Oxygen therapy - nasal cannula, 4 Lpm

Routine, Continuous Device: Nasal Cannula Rate in liters per minute: 4 Lpm Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: Device 2: Device 3: Indications for O2 therapy: Pre-op

Rehab

Consults

Additional Orders