

## General

### Elective Outpatient, Observation (Single Response)

<input type="checkbox"/> Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Outpatient observation services under general supervision	Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/> Outpatient in a bed - extended recovery	Admitting Physician: Bed request comments: PACU & Post-op

## Nursing

### Vital Signs

<input checked="" type="checkbox"/> Vital signs (per unit protocol)	Routine, Per unit protocol, Post-op
---	-------------------------------------

### Activity

<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated Post-op
<input type="checkbox"/> Ambulate with assistance	Routine, 3 times daily Specify: with assistance Post-op
<input type="checkbox"/> Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: Post-op
<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S May resume normal activity 24 hours after PEG placement., Post-op

### Nursing

<input checked="" type="checkbox"/> Observe minimum of 1-2 hours post procedure	Routine, Until discontinued, Starting S, Post-op
<input checked="" type="checkbox"/> Return to floor/unit when stable per Endoscopy protocol	Routine, Until discontinued, Starting S, Post-op
<input checked="" type="checkbox"/> Resume active orders on return to floor/unit	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Positioning instruction	Routine, Until discontinued, Starting S Position: Left Lateral Recumbent Post-op
<input type="checkbox"/> Have patient lie on side until fully awake	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Discontinue IV prior to discharge	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Discharge when meets discharge criteria	Routine, Until discontinued, Starting S, Post-op

### PEG Orders

<input type="checkbox"/> Positioning instruction	Routine, Until discontinued, Starting S For 24 Hours Position: Reverse Trendelenberg Post-op
<input type="checkbox"/> Tube site care	Routine, Every morning Clean PEG site and change PEG dressings with hydrogen peroxide and povidone iodine., Post-op
<input type="checkbox"/> Check residual	Routine, Per unit protocol Check gastric residual volume every 8 hours and hold tube feedings for 2 hours if more than 400 ml. When volume returns below 400 ml, resume tube feeding at previous rate., Post-op

<input type="checkbox"/> Abdominal binder	Routine, Once Waking hours only? Nurse to schedule? Special Instructions: Post-op
<input type="checkbox"/> OK to use PEG for meds now. Ok to start feeds via PEG in 3 hours.	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Keep patients hands away from PEG and prevent patient from pulling PEG out	Routine, Until discontinued, Starting S, Post-op
<b>Notify</b>	
<input type="checkbox"/> Notify Physician for signs of bleeding, abdominal or chest pain, or fever greater than 101.5	Routine, Until discontinued, Starting S, signs of bleeding, abdominal or chest pain, or fever greater than 101.5, Post-op
<b>Diet</b>	
<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: Post-op
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op