

General

Common Present on Admission Diagnosis

<input type="checkbox"/> Present on Admission	Details
<input type="checkbox"/> Acidosis	Details
<input type="checkbox"/> Acute Post-Hemorrhagic Anemia	Details
<input type="checkbox"/> Acute Renal Failure	Details
<input type="checkbox"/> Acute Respiratory Failure	Details
<input type="checkbox"/> Acute Thromboembolism of Deep Veins of Lower Extremities	Details
<input type="checkbox"/> Anemia	Details
<input type="checkbox"/> Bacteremia	Details
<input type="checkbox"/> Bipolar disorder, unspecified	Details
<input type="checkbox"/> Cardiac Arrest	Details
<input type="checkbox"/> Cardiac Dysrhythmia	Details
<input type="checkbox"/> Cardiogenic Shock	Details
<input type="checkbox"/> Decubitus Ulcer	Details
<input type="checkbox"/> Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/> Disorder of the Liver	Details
<input type="checkbox"/> Electrolyte and Fluid Disorder	Details
<input type="checkbox"/> Intestinal Infection due to Clostridium Defficile	Details
<input type="checkbox"/> Methicillin Resistant Staphylococcus Aureus Infection	Details
<input type="checkbox"/> Obstructive Chronic Bronchitis with Exacerbation	Details
<input type="checkbox"/> Other Alteration of Consciousness	Details
<input type="checkbox"/> Other and Unspecified Coagulation Defects	Details
<input type="checkbox"/> Other Pulmonary Embolism and Infarction	Details
<input type="checkbox"/> Phlebitis and Thrombophlebitis	Details
<input type="checkbox"/> Protein-calorie Malnutrition	Details
<input type="checkbox"/> Psychosis, unspecified psychosis type	Details
<input type="checkbox"/> Schizophrenia Disorder	Details
<input type="checkbox"/> Sepsis	Details
<input type="checkbox"/> Septic Shock	Details
<input type="checkbox"/> Septicemia	Details
<input type="checkbox"/> Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not stated as Uncontrolled	Details
<input type="checkbox"/> Urinary Tract Infection, Site Not Specified	Details

Admission or Observation (Single Response) (Selection Required)

<input type="radio"/> Admit to Inpatient	Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="radio"/> Outpatient observation services under general supervision	Admitting Physician: Patient Condition: Bed request comments:
<input type="radio"/> Outpatient in a bed - extended recovery	Admitting Physician: Bed request comments:

Admission or Observation (Single Response)

Patient has active status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. |
| <input type="checkbox"/> Outpatient observation services under general supervision | Admitting Physician:
Patient Condition:
Bed request comments: |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Admitting Physician:
Bed request comments: |

Admission (Single Response)

Patient has active status order on file.

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. |
|---|--|

Code Status

@CERMSG(674511)@

Code Status (Single Response)

DNR and Modified Code orders should be placed by the responsible physician.

- | | |
|--|--|
| <input type="checkbox"/> Full code | Code Status decision reached by: |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Did the patient/surrogate require the use of an interpreter?
Did the patient/surrogate require the use of an interpreter?
Does patient have decision-making capacity? |
| <input type="checkbox"/> Consult to Palliative Care Service | Priority:
Reason for Consult?
Order?
Name of referring provider:
Enter call back number: |
| <input type="checkbox"/> Consult to Social Work | Reason for Consult: |
| <input type="checkbox"/> Modified Code | Did the patient/surrogate require the use of an interpreter?
Did the patient/surrogate require the use of an interpreter?
Does patient have decision-making capacity?
Modified Code restrictions: |

- | | |
|---|---|
| <input type="checkbox"/> Treatment Restrictions ((For use when a patient is NOT in a cardiopulmonary arrest)) | I understand that if the patient is NOT in a cardiopulmonary arrest, the selected treatments will NOT be provided. I understand that all other unselected medically indicated treatments will be provided.
Treatment Restriction decision reached by:
Specify Treatment Restrictions: |
|---|---|

Isolation

- | | |
|---|--------------|
| <input type="checkbox"/> Airborne isolation status | |
| <input type="checkbox"/> Airborne isolation status | Details |
| <input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics. | Once, Sputum |
| <input type="checkbox"/> Contact isolation status | Details |

<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Details
<input type="checkbox"/> Fall precautions	Increased observation level needed:
<input type="checkbox"/> Latex precautions	Details
<input type="checkbox"/> Seizure precautions	Increased observation level needed:

Nursing

Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 6 hours based on acuity status
--	---

Activity

<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S Bathroom Privileges: HOB 30 Degrees
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated
<input checked="" type="checkbox"/> Ambulate patient	Routine, 3 times daily Specify: in hall,with assistance

Nursing

<input checked="" type="checkbox"/> Intake and output	Routine, Every shift Per floor protocol
<input checked="" type="checkbox"/> Weigh patient	Routine, Once For 1 Occurrences Upon arrival
<input checked="" type="checkbox"/> Weigh patient	Routine, Daily, Starting S+1
<input checked="" type="checkbox"/> Head of bed 30 degrees	Routine, Until discontinued, Starting S Head of bed: 30 degrees
<input type="checkbox"/> Nasogastric tube maintenance	Routine, Until discontinued, Starting S Tube Care Orders: To low intermittent wall suction.
<input type="checkbox"/> Foley catheter care	Routine, Until discontinued, Starting S Orders: Maintain To gravity drainage, Foley care per protocol
<input type="checkbox"/> Drain care	Routine, Every 12 hours PRN Drain 1: Drain 2: Drain 3: Drain 4: All Drains: Every 12 hours
<input type="checkbox"/> Measure drainage	Routine, Every 8 hours Type of drain: If abdominal drains in place, empty and record output every 8 hours and as needed
<input type="checkbox"/> Bedside glucose	Routine, 4 times daily before meals and at bedtime Before meals and before bedtime. Notify physician if blood glucose LESS than 70 or GREATER than 180.
<input checked="" type="checkbox"/> Nursing communication	All orders to be cleared through Liver Attending
<input type="checkbox"/> Telemetry	"And" Linked Panel
<input type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes Upon arrival to the unit

<input type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94
---	--

Aspiration Precautions - Nursing

<input type="checkbox"/> Place sign for Aspiration Precautions at HOB	Routine, Until discontinued, Starting S
<input type="checkbox"/> Elevate head of bed 45 degrees	Routine, Until discontinued, Starting S Head of bed: 45 degrees
<input type="checkbox"/> Oral suction to bedside	Routine, As needed Place suction setup at bedside with Yankauer suction tip.
<input type="checkbox"/> If eating/drinking, remain sitting up for 2 hours afterward.	Routine, Until discontinued, Starting S
<input type="checkbox"/> Nasogastric tube insertion	Routine, Once Type: MD to insert if history of gastric bypass.
<input type="checkbox"/> Nasogastric tube to low intermittent suction	Routine, Until discontinued, Starting S Tube Care Orders: To Low Intermittent Suction

Notify

<input type="checkbox"/> Physician communication order	Routine, Once For 1 Occurrences Liver Transplant Service upon patient arrival to Floor
<input type="checkbox"/> Notify Liver Transplant Service	Routine, Until discontinued, Starting S Temperature greater than: 100.5 Temperature less than: Systolic BP greater than: 160 Systolic BP less than: 90 Diastolic BP greater than: Diastolic BP less than: MAP less than: Heart rate greater than (BPM): 110 Heart rate less than (BPM): 60 Respiratory rate greater than: Respiratory rate less than: SpO2 less than:

Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Except meds Pre-Operative fasting options: Give only specifically ordered medications
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Other Options: Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Foods to Avoid:

Oral supplements

Routine
Can/Bottle Supplements:
Can/Bottle Supplements:
Can/Bottle Supplements:
Can/Bottle Supplements:
Can/Bottle Supplements:
Can/Bottle Supplements:
Can/Bottle Supplements:

IV Fluids

IV Fluids (Single Response)

- | | |
|--|--|
| <input type="checkbox"/> sodium chloride 0.45 % infusion | 75 mL/hr, intravenous, continuous, Post-op
Replace urine output with continuous IV 0.45% sodium chloride mL per mL. Replacement fluids not to exceed a maximum of 250 mL per hour and a minimum of 75 mL per hour. |
| <input type="checkbox"/> sodium chloride 0.45 % 1,000 mL with sodium bicarbonate 75 mEq/L infusion | 75 mL/hr, intravenous, continuous, Post-op
Replace urine output with continuous IV 0.45% sodium chloride with 75 mEq sodium bicarbonate mL per mL.
Replacement fluids not to exceed a maximum of 250 mL per hour and a minimum of 75 mL per hour |

Medications

Alternate Route Immunosuppression

- | | |
|---|---|
| <input type="checkbox"/> IV - methylPREDNISolone sodium succinate (Solu-MEDROL) injection | 40 mg, intravenous, daily
If given by IV Push, administer over no less than 3 minutes. |
| <input type="checkbox"/> IV - mycophenolate (CELLCEPT) in dextrose 5% IVPB | intravenous, for 2 Hours, 2 times daily at 0600, 1800 |
| <input type="checkbox"/> Sublingual - tacrolimus (PROGRAF) capsule | sublingual, 2 times daily at 0600, 1800 |
| <input type="checkbox"/> Sublingual - tacrolimus (PROGRAF) capsule - Select this option if patients doses are different for Morning and Evening doses | "And" Linked Panel |
| <input type="checkbox"/> tacrolimus (PROGRAF) capsule | sublingual, daily at 0600 (TIME CRITICAL) |
| <input type="checkbox"/> tacrolimus (PROGRAF) capsule | sublingual, daily at 1800 |

Anti-Viral Prophylaxis (Single Response)

- | | |
|---|---|
| <input type="checkbox"/> ganciclovir (CYTOVENE) Options (Single Response) | |
| <input type="checkbox"/> For CrCL GREATER than 50 mL/min - ganciclovir (CYTOVENE) IVPB | 5 mg/kg, intravenous, nightly, Post-op
Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.
Reason for Therapy: |
| <input type="checkbox"/> For CrCL between 30 - 50 mL/min - ganciclovir (CYTOVENE) IVPB | 2.5 mg/kg, intravenous, nightly, Post-op
Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.
Reason for Therapy: |
| <input type="checkbox"/> For CrCL between 15 - 30 mL/min - ganciclovir (CYTOVENE) IVPB | 0.625 mg/kg, intravenous, nightly, Post-op
Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.
Reason for Therapy: |
| <input type="checkbox"/> For CrCL LESS than 15 mL/min or HD - ganciclovir (CYTOVENE) IVPB | 0.625 mg/kg, intravenous, every 48 hours, Post-op
Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.
Reason for Therapy: |
| <input type="checkbox"/> For CRRT - ganciclovir (CYTOVENE) IVPB | 2.5 mg/kg, intravenous, nightly, Post-op
Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.
Reason for Therapy: |

<input type="checkbox"/> acyclovir (ZOVIRAX)	5 mg/kg, intravenous, every 8 hours Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> acyclovir (ZOVIRAX) oral	200 mg, oral, 2 times daily Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> valACYclovir (VALTREX) tablet	500 mg, oral, 2 times daily Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> valGANciclovir (VALCYTE) tablet	450 mg, oral, 2 times daily Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> valGANciclovir (VALCYTE) tablet	450 mg, oral, user specified, S at 5:00 PM Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> valGANciclovir (VALCYTE) 50 mg/mL oral solution	450 mg, oral, daily Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:

Antifungal Prophylaxis

<input type="checkbox"/> nystatin (MYCOSTATIN) 100,000 unit/mL suspension	5 mL, oral, 4 times daily, Post-op Paint mouth with swab while intubated. Once extubated, swish and swallow. Reason of Therapy:
---	---

Stress Ulcer Prophylaxis

<input type="checkbox"/> pantoprazole (PROTONIX) EC tablet	40 mg, oral, daily at 0600 Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, daily

IV Antibiotics

<input type="checkbox"/> vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, every 6 hours Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> piperacillin-tazobactam (ZOSYN) IV	2.25 g, intravenous, every 6 hours Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> piperacillin-tazobactam (ZOSYN) IV	3.375 g, intravenous, every 6 hours Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> IMipenem/cilastatin IVPB Options (Single Response)	
<input type="checkbox"/> For CrCL GREATER than 60 mL/min - imipenem-cilastatin (PRIMAXIN)	500 mg, intravenous, for 30 Minutes, every 6 hours Reason for Therapy:
<input type="checkbox"/> For CrCL between 30-60 mL/min - imipenem-cilastatin (PRIMAXIN)	500 mg, intravenous, for 30 Minutes, every 8 hours Reason for Therapy:
<input type="checkbox"/> For CrCL between 10-30 mL/min - imipenem-cilastatin (PRIMAXIN)	250 mg, intravenous, for 30 Minutes, every 6 hours Reason for Therapy:
<input type="checkbox"/> For CrCL LESS than 10 mL/min - imipenem-cilastatin (PRIMAXIN)	250 mg, intravenous, for 30 Minutes, every 12 hours Administer while on hemodialysis Reason for Therapy:

[] cefTRIAxone (ROCEPHIN) IV	2 g, intravenous, for 30 Minutes, once, For 1 Doses Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
[] levofloxacin (LEVAQUIN) IV	500 mg, intravenous, every 24 hours Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy:
[] levofloxacin (LEVAQUIN) IV	750 mg, intravenous, once, For 1 Doses Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy:
[] metronidazole (FLAGYL) IV	500 mg, intravenous, for 30 Minutes, once, For 1 Doses Please send cultures prior to starting antibiotics Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy:

Oral Antibiotics

[] ciprofloxacin HCl (CIPRO) tablet	500 mg, oral, 2 times daily at 0600, 1600 Please send cultures prior to starting antibiotics. Take 1 hour before or 2 hours after meals. Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
[] levofloxacin (LEVAQUIN) tablet	500 mg, oral, daily Please send cultures prior to starting antibiotics. Separate by 2 hours from any milk product, antacid or iron. May cause QT interval prolongation Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
[] levofloxacin (LEVAQUIN) tablet	500 mg, oral, daily at 0600 (TIME CRITICAL) Please send cultures prior to starting antibiotics. Separate by 2 hours from any milk product, antacid or iron. May cause QT interval prolongation Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
[] levofloxacin (LEVAQUIN) tablet	750 mg, oral, daily Please send cultures prior to starting antibiotics. Separate by 2 hours from any milk product, antacid or iron. May cause QT interval prolongation Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
[] levofloxacin (LEVAQUIN) tablet	750 mg, oral, daily at 0600 (TIME CRITICAL) Please send cultures prior to starting antibiotics. Separate by 2 hours from any milk product, antacid or iron. May cause QT interval prolongation Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:

<input type="checkbox"/> metroNIDAZOLE (FLAGYL) tablet	500 mg, oral, every 8 hours Please send cultures prior to starting antibiotics. Separate by 2 hours from any milk product, antacid or iron. May cause QT interval prolongation Reason for Therapy:
--	--

Hepatitis B Prophylaxis

<input type="checkbox"/> hepatitis B immune globulin (HEPAGAM B) IVPB 10,000 Units	10,000 Units, intravenous, for 3 Hours, once, For 1 Doses Decrease the rate to 60 mL/hr or LESS if the patient gets uncomfortable, if the patient has infusion related adverse events, or if concern about the infusion speed exists.
--	--

Premedications

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet	25 mg, oral, once, For 1 Doses With sip of water on call to OR. 30 minutes prior to HBIG® infusion
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, once, For 1 Doses With sip of water on call to OR. 30 minutes prior to HBIG® infusion

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) injection	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Restricted Medication

<input checked="" type="checkbox"/> No NSAIDs EXcluding aspirin	STAT, Until discontinued, Starting S Reason for "No" order: transplant patient Pre-op
---	---

Insomnia

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
---	--------------------------------

Other Medications

<input type="checkbox"/> ursodiol (ACTIGALL) capsule	300 mg, oral, 2 times daily
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	500 mg, oral, every 6 hours PRN, fever, Mild pain (score 1-3) PO tab
<input type="checkbox"/> calcium carbonate-vitamin D3 250-125 mg-unit per tablet	2 tablet, oral, 3 times daily
<input type="checkbox"/> magnesium oxide (MAG-OX) tablet	400 mg, oral, 2 times daily
<input type="checkbox"/> multivitamin with minerals tablet	1 tablet, oral, daily
<input type="checkbox"/> insulin GLARGINE (LANTUS) injection	subcutaneous, daily before breakfast Be aware of soundalike/lookalike potential with other insulins.
<input type="checkbox"/> darbepoetin (ARANESP) Non-Oncology Orders	
<input type="checkbox"/> darbepoetin (ARANESP) injection solution	0.45 mcg/kg, subcutaneous, once Aranesp® (darbepoetin alfa) is restricted to the outpatient setting with prior financial approval for non-oncology indications. Do you attest that the patient is being treated in an approved care setting? Diagnosis: Hemoglobin Goal (g/dL): Hematocrit Goal (%):
<input type="checkbox"/> Hemoglobin and hematocrit	STAT For 1 Occurrences
<input type="checkbox"/> Ferritin level	STAT For 1 Occurrences
<input type="checkbox"/> CBC with platelet and differential	STAT For 1 Occurrences
<input type="checkbox"/> Transferrin	STAT For 1 Occurrences
<input type="checkbox"/> Iron level	STAT For 1 Occurrences
<input type="checkbox"/> Creatinine level	STAT For 1 Occurrences
<input type="checkbox"/> Total iron binding capacity	STAT For 1 Occurrences

Aspiration Precautions - Medications

<input type="checkbox"/> chlorhexidine (PERIDEX) 0.12 % solution	15 mL, Swish & Spit, 2 times daily
<input type="checkbox"/> metoclopramide (REGLAN) injection	5 mg, intravenous, every 6 hours Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:
<input type="checkbox"/> metoclopramide (REGLAN) tablet	10 mg, oral, 4 times daily before meals and nightly

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

VTE/DVT Risk Definitions

URL:

"\\appt1\epicappprod\Restricted\OrderSets\VTEDEVTRISK

() Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)	
() Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
() Place/Maintain sequential compression device continuous	Routine, Continuous
() Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
() Place/Maintain sequential compression device continuous	Routine, Continuous
() High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
() Place/Maintain sequential compression device continuous	Routine, Continuous
() High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
() Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> LOW Risk of DVT (Selection Required)		
Low Risk Definition Age less than 60 years and NO other VTE risk factors		
<input type="checkbox"/> Low Risk (Single Response) (Selection Required)		
<input type="checkbox"/>	Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
<input type="checkbox"/> MODERATE Risk of DVT - Surgical (Selection Required)		
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission		
<input type="checkbox"/> Moderate Risk (Selection Required)		
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)		
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)		
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

MODERATE Risk of DVT - Non-Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

<input type="checkbox"/> Moderate risk of VTE	Routine, Once
---	---------------

Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis - **"And" Linked Panel**
Order Sequential compression device

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

Contraindications exist for pharmacologic prophylaxis **"And" Linked Panel**
AND mechanical prophylaxis

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):

<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S+1 For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

HIGH Risk of DVT - Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

<input type="checkbox"/> High risk of VTE	Routine, Once
---	---------------

High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
--	---

enoxaparin for VTE Prophylaxis (Single Response)

<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg daily at 1700	30 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
---	--

<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg every 12 hours	30 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg daily at 1700	40 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg every 12 hours	40 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, Starting S+1 For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1 Indication(s): VTE Prophylaxis

<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

VTE/DVT Risk Definitions

URL:

"\\appt1\epicappprod\Restricted\OrderSets\VTEDEVTRISK DEFINITIONS.pdf"

[Anticoagulation Guide for COVID patients](#)

URL:

"https://formweb.com/files/houstonmethodist/documents/COVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf"

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)

Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)

Moderate risk of VTE

Routine, Once

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	
Low Risk Definition Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/> Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
<input type="checkbox"/> MODERATE Risk of DVT - Surgical (Selection Required)	

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

Moderate risk of VTE Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

Place/Maintain sequential compression device continuous Routine, Continuous

Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 0600, Starting S+1
Indication(s): VTE Prophylaxis

patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 0600, Starting S+1
For Patients with CrCL LESS than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min 40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, Starting S+1
For patients with weight GREATER than 100 kg.

<input type="checkbox"/>	warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/>	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/>	Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/>	MODERATE Risk of DVT - Non-Surgical (Selection Required)	
	Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/>	Moderate Risk (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S+1 For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours Recommended for patients with high risk of bleeding, e.g. weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin for VTE Prophylaxis (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg daily at 1700	30 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg every 12 hours	30 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg daily at 1700	40 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg every 12 hours	40 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, Starting S+1 For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:

<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis

<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
() enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
() Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

DVT Risk and Prophylaxis Tool (Single Response)

VTE/DVT Risk Definitions

URL:

"\\appt1\epicappprod\Restricted\OrderSets\VTEDVTRISK DEFINITIONS.pdf"

[Anticoagulation Guide for COVID patients](#)

URL:

"https://formweb.com/files/houstonmethodist/documents/COVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf"

() Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)

() Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)

Moderate risk of VTE

Routine, Once

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	
Low Risk Definition Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/> Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
<input type="checkbox"/> MODERATE Risk of DVT - Surgical (Selection Required)	

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

Moderate risk of VTE Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

Place/Maintain sequential compression device continuous Routine, Continuous

Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 0600, Starting S+1
Indication(s): VTE Prophylaxis

patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 0600, Starting S+1
For Patients with CrCL LESS than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min 40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
For patients with weight GREATER than 100 kg.

<input type="checkbox"/>	warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/>	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/>	Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/>	MODERATE Risk of DVT - Non-Surgical (Selection Required)	
	Moderate Risk Definition	
	Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	
	One or more of the following medical conditions:	
	CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	
	Age 60 and above	
	Central line	
	History of DVT or family history of VTE	
	Anticipated length of stay GREATER than 48 hours	
	Less than fully and independently ambulatory	
	Estrogen therapy	
	Moderate or major surgery (not for cancer)	
	Major surgery within 3 months of admission	
<input type="checkbox"/>	Moderate Risk (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S+1 For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/>	patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin for VTE Prophylaxis (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg daily at 1700	30 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg every 12 hours	30 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg daily at 1700	40 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg every 12 hours	40 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, Starting S+1 For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min. Indication(s): VTE Prophylaxis

() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min. Indication(s): VTE Prophylaxis
() enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
() Rivaroxaban and Pharmacy Consult (Selection Required)	
[] rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
[] Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
() warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
() Place/Maintain sequential compression device continuous	Routine, Continuous

Labs

COVID-19 Qualitative PCR

[] COVID-19 qualitative RT-PCR - Nasal Swab	STAT For 1 Occurrences Specimen Source: Nasal Swab Is this for pre-procedure or non-PUI assessment? Yes
--	---

Laboratory Every Monday x 3

[X] C-reactive protein	Every Monday For 3 Occurrences, Post-op
[X] Prealbumin level	Every Monday For 3 Occurrences, Post-op
[X] Cytomegalovirus by PCR	Every Monday For 3 Occurrences Specimen Source: Post-op

Laboratory Stat Upon Arrival

[X] Basic metabolic panel	Once, Post-op
[X] Hepatic function panel	Once, Post-op
[X] Magnesium level	Once, Post-op
[X] Phosphorus level	Once, Post-op
[X] Ionized calcium	Once, Post-op
[X] CBC with platelet and differential	Once, Post-op
[X] Prothrombin time with INR	Once, Post-op

<input checked="" type="checkbox"/> Partial thromboplastin time	Once, Post-op
<input checked="" type="checkbox"/> Arterial blood gas	Once, Post-op
<input checked="" type="checkbox"/> LDH	STAT For 1 Occurrences, Post-op
<input checked="" type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, Post-op

Laboratory Daily AM x 3

<input checked="" type="checkbox"/> Basic metabolic panel	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Hepatic function panel	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Magnesium level	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Phosphorus level	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Ionized calcium	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> LDH	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> CBC with platelet and differential	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Prothrombin time with INR	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Partial thromboplastin time	AM draw repeats For 3 Days, Post-op
<input checked="" type="checkbox"/> Arterial blood gas	AM draw repeats For 3 Days While intubated, Post-op
<input checked="" type="checkbox"/> Fibrinogen	AM draw repeats For 3 Occurrences, Post-op

Laboratory Trough Level at 05:30 x 3

<input type="checkbox"/> FK506 Tacrolimus level, random	AM draw repeats, Starting S+1 at 5:30 AM For 3 Days Trough level
<input type="checkbox"/> Cyclosporine level, random	AM draw repeats, Starting S+1 at 5:30 AM For 3 Days Trough level

Microbiology

<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Conditional Frequency For 1 Occurrences Specimen Source: Urine Specimen Site: If temperature greater than 100.5 deg F, Post-op
<input type="checkbox"/> Sputum culture	Conditional Frequency For 1 Occurrences, Sputum, Not otherwise specified If temperature greater than 100.5 deg F, Post-op
<input type="checkbox"/> Blood culture x 2	"And" Linked Panel
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used., Post-op
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used., Post-op
<input type="checkbox"/> Cytomegalovirus by PCR	Conditional Frequency For 1 Occurrences Specimen Source: For temperature GREATER than 100.5 F, Post-op
<input type="checkbox"/> Epstein Barr Virus (EBV) by PCR	STAT For 1 Occurrences Specimen Source: Post-op
<input type="checkbox"/> Respiratory Pathogen Panel with COVID-19 RT-PCR - Suspected (Selection Required)	
<input type="checkbox"/> Respiratory pathogen panel with COVID-19 RT-PCR	Once, Nasopharyngeal, Swab
<input type="checkbox"/> Blood culture, fungus x 2	"And" Linked Panel
<input type="checkbox"/> Blood culture, fungus	Once, Blood
<input type="checkbox"/> Blood culture, fungus	Once, Blood
<input type="checkbox"/> Ova & parasites, direct examination	Once, Stool

Cardiology

Cardiology

<input type="checkbox"/> ECG 12 lead	STAT, Once, Starting S For 1 Occurrences Clinical Indications: Interpreting Physician: STAT upon arrival to unit
<input type="checkbox"/> Transthoracic Echocardiogram Complete, (w contrast, Strain and 3D if needed)	Routine, 1 time imaging, Starting S For 1 Occurrences

Imaging

Diagnostic X-Ray

<input type="checkbox"/> XR Chest 2 Vw W Apical Lordotic	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences Stat upon arrival to unit
<input type="checkbox"/> XR Chest 1 Vw Portable	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences STAT upon arrival to unit
<input checked="" type="checkbox"/> XR Chest 1 Vw Portable	STAT, Conditional Frequency For 1 Occurrences If temperature is greater than 100.5 degrees Farenheit
<input type="checkbox"/> XR Abdomen 1 Vw	Routine, 1 time imaging, Starting S at 1:00 AM For 1

Aspiration Precautions - X-ray

<input type="checkbox"/> XR Chest 1 Vw Portable	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> XR Abdomen Ap And Lateral	Routine, 1 time imaging, Starting S at 1:00 AM For 1

Other Studies

Other Studies

<input type="checkbox"/> CT Abdomen W Contrast (Omnipaque)	"And" Linked Panel For those with iodine allergies, please order the panel with Readi-Cat (barium sulfate).
<input type="checkbox"/> CT Abdomen W Contrast	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> iohexol (OMNIPAQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
<input type="checkbox"/> CT Chest W Wo Contrast	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> CT Angiogram Pe Chest	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> CTA Abdomen W Wo Contrast	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> CTA Abdomen Pelvis W And Or Wo Contrast	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> US Abdominal Doppler	Routine, 1 time imaging, Starting S at 1:00 AM For 1
<input type="checkbox"/> NM Hepatobiliary (HIDA Scan)	Routine, 1 time imaging, Starting S at 1:00 AM For 1

Respiratory

Respiratory Therapy

<input type="checkbox"/> Oxygen therapy	Routine, Continuous Device: Nasal Cannula Rate in liters per minute: Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 90% Indications for O2 therapy: Keep pulse oximetry between 92-95%.
<input checked="" type="checkbox"/> Incentive spirometry	STAT, Every hour while awake, Starting S For 999 Occurrences Every 1 Hour While Awake
<input type="checkbox"/> Encourage deep breathing and coughing	Routine, Every 2 hours For 999 Occurrences Every 2 Hours

Rehab

Consults

For Physician Consult orders use sidebar

Consults

<input checked="" type="checkbox"/> Consult to PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: To evaluate and treat for muscle strengthening and activity
<input checked="" type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Other (Specify) Specify: Post Transplant Diet Education Registered Dietitian
<input checked="" type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Positive Nutrition Screen Nutrition assessment, Registered Dietitian
<input checked="" type="checkbox"/> Consult to Case Management	Consult Reason: Discharge Planning
<input type="checkbox"/> Consult to Transplant Social Work	Reason for Consult? Organ Transplant: Liver Contact Liver Transplant Social Worker at 713-441-5451
<input type="checkbox"/> Consult to Transplant Financial Services	Reason for Consult? Organ Transplant: Liver Discharge Medication Insurance Verification. Contact Liver Transplant Financial Services at 713-441-5451
<input type="checkbox"/> Consult to Diabetes Educator	Reason for Consult: New Onset,Self Care / Meter,Insulin Initiation New onset. Blood sugar checks, insulin sliding scale and diet

Aspiration Precautions - Consults

<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult: Reason for SLP?
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic:

Additional Orders