

Fondaparinux (Arixtra) for Therapeutic Anticoagulation [3957]

This order set is intended for ordering of treatment doses of anticoagulation with fondaparinux (Arixtra) (i.e.: DVT, PE, Acute Coronary Syndrome)

Baseline Labs

Baseline Labs

<input checked="" type="checkbox"/> Basic metabolic panel	Once
<input checked="" type="checkbox"/> CBC hemogram	Once
<input checked="" type="checkbox"/> Prothrombin time with INR	Once
<input checked="" type="checkbox"/> Partial thromboplastin time, activated	Once

Medications

Fondaparinux for Therapeutic Anticoagulation (Single Response) (Selection Required)

Fondaparinux (Arixtra) is not recommended for use in patients with CrCl LESS THAN 30mL/min. Use caution in patients with CrCl 30mL/min to 50mL/min

() For patient weight LESS THAN 50kg - fondaparinux (ARIXTRA) 5 mg Every 24 Hours

fondaparinux (ARIXTRA) injection 5 mg, subcutaneous, every 24 hours
Indication(s):

() For patient weight 50kg to 100kg - fondaparinux (ARIXTRA) 7.5 mg Every 24 Hours

fondaparinux (ARIXTRA) injection 7.5 mg, subcutaneous, every 24 hours
Indication(s):

() For patient weight GREATER THAN 100kg - fondaparinux (ARIXTRA) 10 mg Every 24 Hours

fondaparinux (ARIXTRA) injection 10 mg, subcutaneous, every 24 hours
Indication(s):

Labs Continuing

Labs Continuing

Consider anti-Xa monitoring for fondaparinux (Arixtra) in patients with CrCl less than 50mL/min, extremes in body weight, pregnancy, or elderly patients. Anti-Xa levels can be obtained 3 hours after the 3rd dose.

<input type="checkbox"/> Basic metabolic panel	Daily For 3 Days
<input type="checkbox"/> CBC hemogram	Daily For 3 Days
<input type="checkbox"/> CBC hemogram	Every 72 hours For 3 Occurrences
<input type="checkbox"/> Anti Xa Arixtra	Once