

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="checkbox"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Outpatient observation services under general supervision	Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Outpatient in a bed - extended recovery	Admitting Physician: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Admit to Inpatient	Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

@CERMSG(674511:):@

Code Status (Single Response)

DNR and Modified Code orders should be placed by the responsible physician.

- | | |
|--|--|
| <input type="checkbox"/> Full code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Did the patient/surrogate require the use of an interpreter?
Did the patient/surrogate require the use of an interpreter?
Does patient have decision-making capacity?
Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Did the patient/surrogate require the use of an interpreter? Did the patient/surrogate require the use of an interpreter? Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions ((For use when a patient is NOT in a cardiopulmonary arrest))	I understand that if the patient is NOT in a cardiopulmonary arrest, the selected treatments will NOT be provided. I understand that all other unselected medically indicated treatments will be provided. Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum, Post-op
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol, Post-op
<input checked="" type="checkbox"/> Hemodynamic Monitoring	Routine, Continuous Measure: Arterial Line MAP, Arterial Line BP Post-op

Activity

<input checked="" type="checkbox"/> Dangle at bedside	Routine, Once Begin on POD 0, Post-op
<input checked="" type="checkbox"/> Out of bed	Routine, Until discontinued, Starting S Specify: Out of bed, Up in chair Additional modifier: for meals Chair x 3 daily, Post-op
<input checked="" type="checkbox"/> Ambulate	Routine, 4 times daily Specify: in hall, with assistance If needed, Post-op

Nursing

<input checked="" type="checkbox"/> Daily weights	Routine, Daily Weigh patient at 0800 Daily, Post-op
---	--

[X] Head of bed	Routine, Until discontinued, Starting S Head of bed: other degrees (specify) Specify: 35 Post-op
[X] Neurological assessment	Routine, Every hour, Starting S Assessment to Perform: Cranial Nerves, Glasgow Coma Scale, Level of Consciousness, Level of Sedation, Pupils Post-op
[X] Site care	Routine, Per unit protocol Site: epicardial pacing wire site Post-op
[X] Apply warming blanket (bair hugger)	Routine, Once For 1 Occurrences To achieve body temperature of 98.6 F, Post-op
[X] Foley catheter care	Routine, 2 times daily Orders: Maintain Clean with CHG cloths, Post-op
[X] Chest tube to continuous suction	Routine, Until discontinued, Starting S Level of suction: 20 cm H2O Post-op
[X] Tube site care (chest tube)	Routine, Per unit protocol Chest tube site care daily and prn per protocol, Post-op
[X] Oral care	Routine, 2 times daily Every 12 hours Per CVICU protocol. Toothbrush every 12 hours, Post-op
[X] Bedside glucose	Routine, Every hour For Until specified (Q1 hour x 6) ONLY IF HISTORY OF DIABETES Routine, Every hour For Until specified Monitor every hour for first 6 hours then change to every 4 hours if not started on an insulin drip; Notify physician for blood glucose less than 70 mg/dL OR blood glucose greater than 300 mg / dL, Post-op
[X] Pacemaker settings	Routine, Until discontinued, Starting S Atrial Setting (MA): Ventricular Setting (MA): Sensitivity Setting (millivolts): AV Interval (milliseconds): Options: Post-op
[X] If rhythm changes to arterial fibrillation and hemodynamics are stable, physician call not required	Routine, Once, Post-op
[X] If patient is being externally paced, have monitor set to detect/capture pacing spikes	Routine, Once, Post-op
[X] Apply Bacitracin ointment to chest tube sites and cover with 2x2s	Routine, Once, Post-op
[X] For chest tube removal	Routine, Once Please have the following supplies at bedside: bacitracin ointment, 2x2s gauze, suture removal kit and wide tape, Post-op
Discontinue	
[X] Discontinue arterial line	Routine, Conditional Frequency For 1 Occurrences Before transfer out of ICU; if arterial line not already discontinued, Post-op
[X] Foley catheter - discontinue	Routine, Conditional Frequency For 1 Occurrences 1) Remove Foley cath POD 1 or POD 2; If unable to remove Foley reason for not removing MUST be documented on POD 1 or POD 2. , Post-op
[] Discontinue Pacemaker Generator and Insulate Pacer Wires	Routine, Conditional Frequency For 1 Occurrences Before transfer out of ICU; if not already discontinued. , Post-op

Notify

<input checked="" type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 102.5 Temperature less than: 95 Systolic BP greater than: 180 Systolic BP less than: 80 Diastolic BP greater than: Diastolic BP less than: MAP less than: 55 Heart rate greater than (BPM): 120 Heart rate less than (BPM): 40 Respiratory rate greater than: 30 Respiratory rate less than: SpO2 less than: 90
--	---

<input checked="" type="checkbox"/> Notify Physician - for chest output greater than 200 milliliters/hour	Routine, Until discontinued, Starting S, for chest output greater than 200 milliliters/hour, Post-op
---	--

<input checked="" type="checkbox"/> Notify Physician - for urine output LESS THAN 160 ml/ 8 hour shift or less then 240 mL per 12 hour shift	Routine, Until discontinued, Starting S, for urine output LESS THAN 160 ml/ 8 hour shift or less then 240 mL per 12 hour shift, Post-op
--	---

Diet

<input checked="" type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
--	---

<input type="checkbox"/> Prune Juice or Prunes	Routine, Until discontinued, Starting S Give with breakfast daily starting post op day 2, Post-op
--	--

Medications

PostOp Antibiotics: For Patients LESS than or EQUAL to 120 kg (Single Response)

<input checked="" type="checkbox"/> ceFAZolin (ANCEF) IV 1g	1 g, intravenous, every 8 hours, For 2 Doses, Post-op Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> If Beta-Lactam Allergic - vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Administer 12 hours after procedure Reason for Therapy: Surgical Prophylaxis

Post-Op Antibiotics: For Patients GREATER than 120 kg (Single Response)

<input checked="" type="checkbox"/> ceFAZolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, every 8 hours, For 2 Doses, Post-op Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> If Beta-Lactam Allergic - vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Administer 12 hours after procedure Reason for Therapy: Surgical Prophylaxis

ACE Inhibitors (Single Response)

<input type="checkbox"/> captopril (CAPOTEN) tablet	25 mg, oral, 3 times daily, Post-op Consult MD before administering if urine output less than 0.5 mL/kg/hour and creatinine greater than 1.3. BP & HR HOLD parameters for this order: Hold Parameters requested BP & HR HOLD for: Other Please specify: 90 HOLD for Heart Rate LESS than: Contact Physician if:
---	---

enalapril (VASOTEC) tablet 2.5 mg, oral, 2 times daily, Post-op
 Consult MD before administering if urine output less than 0.5 mL/kg/hr and creatinine greater than 1.3.
 BP & HR HOLD parameters for this order: Hold Parameters requested
 BP & HR HOLD for: Other
 Please specify: 90
 HOLD for Heart Rate LESS than:
 Contact Physician if:

lisinopril (PRINIVIL,ZESTRIL) tablet 5 mg, oral, daily, Post-op
 Consult MD before administering if urine output less than 0.5 mL/kg/hr and creatinine greater than 1.3.
 BP & HR HOLD parameters for this order: Hold Parameters requested
 BP & HR HOLD for: Other
 Please specify: 90 mmHg
 HOLD for Heart Rate LESS than:
 Contact Physician if:

amlODarone (CORDARONE) 24-hr Infusions HARD-Stop (Single Response)

Loading Dose and Maintenance Infusion (Single Response)

Select Standard or Double concentration

Standard

CENTRAL Line Administration: amlODarone (CORDArone) 150 mg LOADING Dose followed by STANDARD concentration 24-hour Infusion for Atrial Fibrillation- NOT HMWB **"Followed by" Linked Panel**

amlODarone (CORDArone) 150 mg BOLUS 150 mg, intravenous, once, Starting S, For 1 Doses, Post-op
 Patients should be monitored for QTc prolongation.

amlODarone 1.8 mg/mL (STANDARD concentration) infusion 1 mg/min, intravenous, continuous, Starting H+10 Minutes, Post-op
 Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours.
 HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.

REDUCE rate for amlODarone (CORDArone) 450 mg/ 250 mL NS 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses, Post-op
 HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused.

amlODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag 0.5 mg/min, intravenous, continuous, Starting H+8 Hours, Post-op
 HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours.

CENTRAL Line Administration: amlODarone (CORDArone) 150 mg LOADING Dose followed by STANDARD concentration 24-hour Infusion for Atrial Fibrillation-HMWB ONLY **"Followed by" Linked Panel**

amlODarone (CORDArone) 150 mg BOLUS 150 mg, intravenous, once, Starting S, For 1 Doses
 Patients should be monitored for QTc prolongation.

amlODarone 1.8 mg/mL (STANDARD concentration) infusion 1 mg/min, intravenous, continuous, Starting H+10 Minutes
 Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours.
 HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.

REDUCE rate for amlODarone (CORDArone) infusion 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses
 HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused.

<input type="checkbox"/>	amIODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag	0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours.
() Double		
<input type="checkbox"/>	CENTRAL Line Administration: amIODarone (CORDArone) 150 mg LOADING Dose followed by DOUBLE concentration 24-hour Infusion for Atrial Fibrillation	"Followed by" Linked Panel
<input type="checkbox"/>	amIODarone (CORDArone) 150 mg BOLUS	150 mg, intravenous, once, Starting S, For 1 Doses, Post-op Patients should be monitored for QTc prolongation.
<input type="checkbox"/>	amIODarone (CORDArone) 900 mg/ 250 mL NS	1 mg/min, intravenous, continuous, Starting H+10 Minutes, Post-op Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line.
<input type="checkbox"/>	REDUCE rate for amIODarone (CORDArone) 900 mg/ 250 mL infusion	0.5 mg/min, intravenous, continuous, Starting H+6 Hours, Post-op Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. Do not take down 1st infusion until entire content of bag is infused.
() Maintenance Infusion (Single Response) Select Standard or Double Concentration		
() Standard		
<input type="checkbox"/>	NO LOADING DOSE - Central Line Administration: amIODarone (CORDArone) STANDARD concentration 24-hour Infusion for Atrial Fibrillation - NOT HMWB	"Followed by" Linked Panel
<input type="checkbox"/>	amIODarone 1.8 mg/mL (STANDARD concentration) infusion	1 mg/min, intravenous, continuous, Post-op Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line.
<input type="checkbox"/>	REDUCE rate for amIODarone (CORDArone) 450 mg/ 250 mL NS	0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses, Post-op Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused.
<input type="checkbox"/>	amIODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag	0.5 mg/min, intravenous, continuous, Starting H+8 Hours, Post-op HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.
<input type="checkbox"/>	NO LOADING DOSE - Central Line Administration: amIODarone (CORDArone) STANDARD concentration 24-hour Infusion for Atrial Fibrillation - HMWB Only	"Followed by" Linked Panel
<input type="checkbox"/>	amIODarone 1.8 mg/mL (STANDARD concentration) infusion	1 mg/min, intravenous, continuous Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.
<input type="checkbox"/>	REDUCE rate for amIODarone (CORDArone) 360 mg/ 200 mL NS	0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused.

<input type="checkbox"/> amlODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag	0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.
---	---

() Double (Single Response)

() NO LOADING DOSE - Central Line Administration:
amlODarone (CORDArone) Double Concentration
24-hour Infusion for Atrial Fibrillation

"Followed by" Linked Panel

<input type="checkbox"/> amlODarone (CORDArone) 900 mg/ 250 mL NS	1 mg/min, intravenous, continuous, Post-op Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.
---	---

<input type="checkbox"/> REDUCE rate for amlODarone (CORDArone) 900 mg/ 250 mL NS	0.5 mg/min, intravenous, continuous, Starting H+6 Hours, Post-op Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. Do not take down 1st infusion until entire content of bag is infused.
---	---

amlODarone (PACerone) tablet

You MUST be sure the oral tablet order is set to start TOMORROW with the start time set to 24 hours AFTER the start time of the INITIAL infusion order above.

<input type="checkbox"/> amlODarone (PACERONE) tablet **** You MUST CHANGE the START DATE to TOMORROW and set the Start TIME to be 24 hours after the Start Time of the Infusion	200 mg, oral, every 24 hours, Starting H+24 Hours amiodarone (Pacerone) tablets must start 24 hours after the start of the infusion order.
--	---

Beta Blockers (Single Response)

() metoprolol tartrate (LOPRESSOR) tablet	25 mg, oral, 2 times daily at 0600, 1800, Starting S+1, Post-op DO NOT administer if patient is on inotrope, vasopressor or has epicardial pacing BP & HR HOLD parameters for this order: Hold Parameters requested BP & HR HOLD for: 110 mmHg HOLD for Heart Rate LESS than: Other Other Heart Rate (in bpm): 60 Contact Physician if:
--	---

() carvedilol (COREG) tablet	3.125 mg, oral, 2 times daily at 0600, 1800, Starting S+1, Post-op DO NOT administer if heart rate is less than 60; systolic blood pressure is less than 110; on inotrope, vasopressor or has epicardial pacing BP & HR HOLD parameters for this order: Hold Parameters requested BP & HR HOLD for: 110 mmHg HOLD for: Contact Physician if:
-------------------------------	---

colchicine

<input type="checkbox"/> colchicine tablet FOR DIABETIC ONLY	0.6 mg, oral, daily, Post-op For prevention of atrial fibrillation post cardiac surgery. Call provider for diarrhea.
--	---

furosemide (LASIX) Oral or IV (Single Response)

(X) furosemide (LASIX) tablet	20 mg, oral, daily, Starting S If unable to swallow oral tablets, discontinue and change to IV daily.
-------------------------------	--

() furosemide (LASIX) IV	20 mg, intravenous, daily
---------------------------	---------------------------

predniSONE oral taper

<input type="checkbox"/>	predniSONE oral taper	"Followed by" Linked Panel
<input type="checkbox"/>	predniSONE (DELTASONE) tablet 15 mg BID	15 mg, oral, 2 times daily, For 6 Doses, Post-op
<input type="checkbox"/>	predniSONE (DELTASONE) tablet 10 mg BID	10 mg, oral, 2 times daily, For 6 Doses, Post-op
<input type="checkbox"/>	predniSONE (DELTASONE) tablet 10 mg daily	10 mg, oral, daily, For 3 Doses, Post-op

Respiratory Medications

<input type="checkbox"/>	Scheduled	
<input type="checkbox"/>	Scheduled - albuterol (PROVENTIL) nebulizer solution	2.5 mg, nebulization, Respiratory Therapy - every 6 hours, Post-op Aerosol Delivery Device: Hand-Held Nebulizer
<input type="checkbox"/>	Scheduled - ipratropium (ATROVENT) 0.02 % nebulizer solution	0.5 mg, nebulization, Respiratory Therapy - every 6 hours, Post-op Aerosol Delivery Device: Hand-Held Nebulizer
<input type="checkbox"/>	PRN	
<input type="checkbox"/>	PRN - albuterol (PROVENTIL) nebulizer solution	2.5 mg, nebulization, every 6 hours PRN, wheezing, Post-op Aerosol Delivery Device: Hand-Held Nebulizer
<input type="checkbox"/>	PRN - ipratropium (ATROVENT) 0.02 % nebulizer solution	0.5 mg, nebulization, every 6 hours PRN, wheezing, Post-op Aerosol Delivery Device: Hand-Held Nebulizer

Multimodal Pain Management

<input checked="" type="checkbox"/>	pregabalin (LYRICA) capsule	100 mg, oral, 2 times daily
<input checked="" type="checkbox"/>	dexMEDEtomidine (PREcedex) infusion	
<input checked="" type="checkbox"/>	Richmond agitation sedation scale	Routine, Per unit protocol Hold infusion daily at: Do not hold sedation Reason sedation not held. Patient on: Target RASS: -1 BIS Monitoring (Target BIS: 40-60): Reassess RASS at least Every 4 Hours and PRN. Follow titration instructions in sedative order. During daily sedation hold, restart sedation protocol if any of the following occur MAP less than 50mmHg or greater than 120mmHg Development of acute distress HR greater than 120 bpm RR greater than 38 breaths/min SpO2 less than 88%
<input checked="" type="checkbox"/>	dexMEDEtomidine (PREcedex) infusion	0.2 mcg/kg/hr, intravenous, continuous, Post-op Titrate for postoperative pain in increments of 0.1 mcg/kg/hr up to maximum dose of 0.6 mcg/kg/hr. If needed for sedation, this order will need to be modified to the ICU sedation order to include titration parameters and dose range. Discontinue Dexmedetomidine (Precedex) IV infusion after extubation. Discontinue on postoperative day 1. Reassess RASS within 1 hour. DESIRED sedation effect: Continue the same rate. Reassess sedation within 4 hours GREATER than desired sedation effect: DECREASE rate by 0.1 mcg/kg/hour. Reassess RASS within one hour.
<input checked="" type="checkbox"/>	acetaminophen (OFIRMEV) intravenous solution	1,000 mg, intravenous, for 15 Minutes, every 6 hours, For 1 Doses, Post-op Total Tylenol/ acetaminophen dose (which includes, IV, PO or combination i.e. Norco, APAP etc) IV acetaminophen (Ofirmev) is restricted to use only in OR, PACU, or ICU areas, and for patients that cannot tolerate oral, per tube, or rectal routes of administration. Do you attest that this restriction has been met?
<input checked="" type="checkbox"/>	acetaminophen (TYLENOL) tablet	500 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Starting H+6 Hours, Post-op May alternate with ibuprofen 400 mg oral every 3 hours. Total Tylenol/ acetaminophen dose (which includes, IV, PO or combination i.e. Norco, APAP etc)

<input checked="" type="checkbox"/> ibuprofen (ADVIL) tablet OR ketorolac (TORadol) IV (Single Response)	
<input type="checkbox"/> ibuprofen (ADVIL) tablet	400 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op May alternate with Tylenol 500 mg every 3 hours.
<input type="checkbox"/> keTOROlac (TORadol) IV "Followed by" Linked Panel	
<input type="checkbox"/> ketorolac (TORADOL) injection	30 mg, intravenous, once, For 1 Doses, Post-op Maximum 120mg/day in adults more than 50kg. Maximum 60mg/day in adults less than 50kg.
<input type="checkbox"/> ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours, Starting H+6 Hours, For 3 Doses, Post-op Maximum 120mg/day in adults more than 50kg. Maximum 60mg/day in adults less than 50kg.

Breakthrough Pain

<input checked="" type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 10-325 mg per tablet	1 tablet, oral, every 4 hours PRN, severe pain (score 7-10), Post-op Total Tylenol/ acetaminophen dose (which includes, IV, PO or combination i.e. Norco, APAP etc)
<input checked="" type="checkbox"/> morPHINE injection	2 mg, intravenous, every 1 hour prn, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed.

PUD Prophylaxis (Single Response)

<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, 2 times daily, Post-op
<input checked="" type="checkbox"/> pantoprazole (PROTONIX) EC tablet	40 mg, oral, daily before breakfast, Post-op Indication(s) for Proton Pump Inhibitor (PPI) Therapy:

Bowel Care

<input checked="" type="checkbox"/> Scheduled	
<input checked="" type="checkbox"/> Scheduled: polyethylene glycol (MIRALAX) packet - POD #1	17 g, oral, daily, Starting S+1, Post-op
<input checked="" type="checkbox"/> Docusate - Oral OR Nasogastric "Or" Linked Panel	
<input checked="" type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily, Post-op Give if patient can tolerate oral medication
<input checked="" type="checkbox"/> docusate (COLACE) 50 mg/5 mL liquid	100 mg, oral, 2 times daily, Post-op Give if patient has a nasogastric tube
<input type="checkbox"/> polyethylene glycol (MIRALAX) packet - start today	17 g, oral, daily, Post-op
<input type="checkbox"/> PRN	
<input type="checkbox"/> As Needed: polyethylene glycol (MIRALAX) packet	17 g, oral, daily PRN, constipation, Post-op RN may use second option based on the patient response to the first option attempted.
<input type="checkbox"/> As Needed: Docusate - Oral OR Nasogastric "Or" Linked Panel	
<input type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily PRN, constipation, Post-op RN may use second option based on the patient response to the first option attempted.
<input type="checkbox"/> docusate (COLACE) 50 mg/5 mL liquid	100 mg, oral, 2 times daily PRN, constipation, Post-op RN may use second option based on the patient response to the first option attempted. Use if cannot swallow capsule.
<input type="checkbox"/> sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet	1 tablet, oral, 2 times daily PRN, constipation, Post-op AS NEEDED AFTER FIRST BM
<input type="checkbox"/> bisacodyl (DULCOLAX) suppository	10 mg, rectal, daily PRN, constipation, Post-op FOR RECTAL USE ONLY. AS NEEDED TO MAINTAIN 3 BOWEL MOVEMENTS PER WEEK. DO NOT GIVE IF DIARRHEA NOTED. Administer if patient has not had a BM in 24 hours after oral therapy

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

VTE/DVT Risk Definitions

URL:

"\appt1\epicapprod\Restricted\OrderSets\VTEDEVTRISK

[Anticoagulation Guide for COVID patients](#)

- () Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)
- () Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)
- | | |
|---|--|
| <input type="checkbox"/> Moderate risk of VTE | Routine, Once, PACU & Post-op |
| <input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis | Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op |
| <input type="checkbox"/> Place sequential compression device (Single Response) | |
| () Contraindications exist for mechanical prophylaxis | Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op |
| () Place/Maintain sequential compression device continuous | Routine, Continuous, PACU & Post-op |
- () Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)
- | | |
|---|--|
| <input type="checkbox"/> Moderate risk of VTE | Routine, Once, PACU & Post-op |
| <input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis | Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op |
| <input type="checkbox"/> Place sequential compression device (Single Response) | |
| () Contraindications exist for mechanical prophylaxis | Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op |
| () Place/Maintain sequential compression device continuous | Routine, Continuous, PACU & Post-op |
- () High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)
- | | |
|---|--|
| <input type="checkbox"/> High risk of VTE | Routine, Once, PACU & Post-op |
| <input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis | Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op |
| <input type="checkbox"/> Place sequential compression device (Single Response) | |
| () Contraindications exist for mechanical prophylaxis | Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op |
| () Place/Maintain sequential compression device continuous | Routine, Continuous, PACU & Post-op |
- () High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)
- | | |
|---|-------------------------------|
| <input type="checkbox"/> High risk of VTE | Routine, Once, PACU & Post-op |
|---|-------------------------------|

<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	LOW Risk of DVT (Selection Required)	
	Low Risk Definition Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/>	Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/>	Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/>	MODERATE Risk of DVT - Surgical (Selection Required)	
	Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/>	Moderate Risk (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	

() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
() patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
() patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
() patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
() warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
[] Moderate Risk (Selection Required)	
[] Moderate risk of VTE	Routine, Once, PACU & Post-op

[] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
() Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
[] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
[] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
[] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
[] Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
() patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
() patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
() patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, PACU & Post-op For patients with weight GREATER than 100 kg.
() warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

HIGH Risk of DVT - Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Surgical Patient

(Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin for VTE Prophylaxis (Single Response)

enoxaparin (LOVENOX) 30 mg daily at 1700 30 mg, subcutaneous, daily at 1700
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) 30 mg every 12 hours 30 mg, subcutaneous, every 12 hours
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) 40 mg daily at 1700 40 mg, subcutaneous, daily at 1700
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) 40 mg every 12 hours 40 mg, subcutaneous, every 12 hours
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1
If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, Starting S+1
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet oral, daily at 1700, Starting S+1
Indication:

Pharmacy consult to manage warfarin (COUMADIN) STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

HIGH Risk of DVT - Non-Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op
Indication(s): VTE Prophylaxis

patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op
For Patients with CrCL LESS than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min 40 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, PACU & Post-op
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.
This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, PACU & Post-op

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, PACU & Post-op
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, PACU & Post-op
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet oral, daily at 1700, PACU & Post-op
Indication:

Pharmacy consult to manage warfarin (COUMADIN) STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1, PACU & Post-op Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), PACU & Post-op Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

VTE/DVT Risk Definitions

URL:

"\\appt1\epicappprod\Restricted\OrderSets\VTEDEVTRISK DEFINITIONS.pdf"

[Anticoagulation Guide for COVID patients](#)

URL:

"https://formweb.com/files/houstonmethodist/documents/C OVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf"

<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)	
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	
Low Risk Definition Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/> Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/> Moderate Risk (Selection Required)	

<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
()	Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
()	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
()	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
()	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
()	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
()	patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
()	patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
()	fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
()	heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
()	heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
()	HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
()	warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
()	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/>	Mechanical Prophylaxis (Single Response) (Selection Required)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition	
Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	
One or more of the following medical conditions:	
CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	
Age 60 and above	
Central line	
History of DVT or family history of VTE	
Anticipated length of stay GREATER than 48 hours	
Less than fully and independently ambulatory	
Estrogen therapy	
Moderate or major surgery (not for cancer)	
Major surgery within 3 months of admission	
<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin for VTE Prophylaxis (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg daily at 1700	30 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg every 12 hours	30 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg daily at 1700	40 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg every 12 hours	40 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, Starting S+1 For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op

<input type="checkbox"/>	heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/>	HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/>	warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/>	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)		
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)		
High Risk Definition		
Both pharmacologic AND mechanical prophylaxis must be addressed.		
One or more of the following medical conditions:		
Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)		
Severe fracture of hip, pelvis or leg		
Acute spinal cord injury with paresis		
Multiple major traumas		
Abdominal or pelvic surgery for CANCER		
Acute ischemic stroke		
History of PE		
<input type="checkbox"/> High Risk (Selection Required)		
<input type="checkbox"/>	High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)		
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/>	aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)		
<input type="checkbox"/>	apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1, PACU & Post-op Indications: VTE prophylaxis
<input type="checkbox"/>	Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)		
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min. Indication(s): VTE Prophylaxis

() enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
() Rivaroxaban and Pharmacy Consult (Selection Required)	
[] rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), PACU & Post-op Indications: VTE prophylaxis
[] Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
() warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

Labs

Labs Today

[] Lactic acid level	Once, Post-op
[X] Basic metabolic panel	Once, Post-op
[X] CBC with platelet and differential	Once, Post-op
[] Magnesium level	Once, Post-op
[] Phosphorus level	Once, Post-op
[] Calcium level	Once, Post-op
[] Ionized calcium	Once, Post-op
[] Prothrombin time with INR	Once, Post-op
[] Partial thromboplastin time	Once, Post-op
[] Platelet function P2Y12	Once, Post-op
[] Platelet mapping	Once Anticoagulant Therapy: Diagnosis: Fax Number (For TEG Graph Result): Post-op
[] Troponin T	Once, Post-op
[] B natriuretic peptide	Once, Post-op
[] Anti Xa, unfractionated	Once, Post-op
[] Fibrinogen	Once, Post-op

<input type="checkbox"/>	Cortisol level, random	Once, Post-op
<input type="checkbox"/>	Type and screen	
<input type="checkbox"/>	Type and screen	Once, Post-op
<input type="checkbox"/>	ABO and Rh confirmation	Once, Blood Bank Confirmation
<input type="checkbox"/>	Blood gas, arterial	Once, Post-op

Labs Today

<input type="checkbox"/>	Lactic acid level	Once, Post-op
<input checked="" type="checkbox"/>	Basic metabolic panel	Once, Post-op
<input checked="" type="checkbox"/>	CBC with platelet and differential	Once, Post-op
<input type="checkbox"/>	Magnesium level	Once, Post-op
<input type="checkbox"/>	Phosphorus level	Once, Post-op
<input type="checkbox"/>	Calcium level	Once, Post-op
<input type="checkbox"/>	Ionized calcium	Once, Post-op
<input type="checkbox"/>	Prothrombin time with INR	Once, Post-op
<input type="checkbox"/>	Partial thromboplastin time	Once, Post-op
<input type="checkbox"/>	Platelet function P2Y12	Once, Post-op
<input type="checkbox"/>	Troponin T	Once, Post-op
<input type="checkbox"/>	B natriuretic peptide	Once, Post-op
<input type="checkbox"/>	Anti Xa, unfractionated	Once, Post-op
<input type="checkbox"/>	Fibrinogen	Once, Post-op
<input type="checkbox"/>	Cortisol level, random	Once, Post-op
<input type="checkbox"/>	Type and screen	
<input type="checkbox"/>	Type and screen	Once, Post-op
<input type="checkbox"/>	ABO and Rh confirmation	Once, Blood Bank Confirmation
<input type="checkbox"/>	Blood gas, arterial	Once, Post-op

Cardiology

Cardiology

<input checked="" type="checkbox"/>	ECG 12 lead - Once	Routine, Once Clinical Indications: Post-Op Surgery Interpreting Physician: Post operative, Post-op
<input type="checkbox"/>	ECG 12 lead - Daily starting tomorrow	Routine, Daily, Starting S+1 For 3 Occurrences Clinical Indications: Post-Op Surgery Interpreting Physician: Post-op
<input type="checkbox"/>	Transthoracic Echocardiogram Complete, (w contrast, Strain and 3D if needed)	Routine, 1 time imaging, Starting S at 1:00 AM, Post-op

Imaging

X-Ray

<input checked="" type="checkbox"/>	Chest 1 Vw Portable	Routine, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences, Post-op
<input checked="" type="checkbox"/>	XR Chest 1 Vw Portable	Routine, 1 time imaging, Starting S at 1:00 AM For 1 At 0700 in ICU, Post-op
<input type="checkbox"/>	Chest 1 Vw Portable (Daily)	Routine, Daily imaging For 3 Occurrences, Post-op
<input checked="" type="checkbox"/>	Chest 1 Vw Portable(after chest tube removal)	Routine, Conditional Frequency For 1 Occurrences After chest tube removal, Post-op

Ultrasound

<input type="checkbox"/>	Pv carotid duplex	Routine, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences, Post-op
--------------------------	-------------------	---

Respiratory

Respiratory

<input checked="" type="checkbox"/>	Encourage deep breathing and coughing	Routine, Every hour, Post-op
-------------------------------------	---------------------------------------	------------------------------

<input checked="" type="checkbox"/> Incentive spirometry	Routine, As directed 10 x every hour while awake, Post-op
<input type="checkbox"/> Positive Expiratory (PEP) Device	Routine, Once Twenty (20) times every hour while awake, Post-op
<input type="checkbox"/> Oxygen therapy	Routine, Continuous Device: Device 2: Device 3: Titrate to keep O2 Sat Above: Indications for O2 therapy: Post-op

Consults

For Physician Consult orders use sidebar

Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason: Discharge Planning Post-op
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Consult PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: Post-op
<input type="checkbox"/> Consult PT wound care	Special Instructions: Location of Wound? Post-op
<input type="checkbox"/> Consult OT eval and treat	Reason for referral to Occupational Therapy (mark all that apply): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: Post-op
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic: Post-op
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult? Post-op
<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult: Post-op
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult: Reason for consult: Post-op
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult? Post-op