

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="radio"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="radio"/>	Outpatient observation services under general supervision	Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="radio"/>	Outpatient in a bed - extended recovery	Admitting Physician: Bed request comments: PACU & Post-op
<input type="radio"/>	Admit to inpatient	Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status (Single Response)

- | | |
|--|--|
| <input type="checkbox"/> Full code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Did the patient/surrogate require the use of an interpreter?
Did the patient/surrogate require the use of an interpreter?
Does patient have decision-making capacity?
Post-op |
| <input type="checkbox"/> Consult to Palliative Care Service | Priority:
Reason for Consult?
Order?
Name of referring provider:
Enter call back number: |

<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Did the patient/surrogate require the use of an interpreter? Did the patient/surrogate require the use of an interpreter? Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions ((For use when a patient is NOT in a cardiopulmonary arrest))	I understand that if the patient is NOT in a cardiopulmonary arrest, the selected treatments will NOT be provided. I understand that all other unselected medically indicated treatments will be provided. Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum, Post-op
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input checked="" type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min For 999 Occurrences Times 4, then every 30 minutes times 4, then every hour times 4, then every 4 hours times 24 hours, then every 8 hours if stable., PACU & Post-op
---	---

Activity

<input type="checkbox"/> Ambulate patient	Routine, 3 times daily Specify: with assistive device Device: Day of surgery, Post-op
<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Up with assistance	Routine, As needed Specify: Up with assistance Post-op
<input type="checkbox"/> Weight bearing	
<input type="checkbox"/> Weight bearing LLE	Routine, Until discontinued, Starting S Weight Bearing Status: Extremity: Post-op
<input type="checkbox"/> Weight bearing RLE	Routine, Until discontinued, Starting S Weight Bearing Status: Extremity: Post-op

<input type="checkbox"/> Consult to PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status:
<input type="checkbox"/> Consult to PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status:
<input type="checkbox"/> Elevate Operative Extremity	Routine, Until discontinued, Starting S Position: Additional instructions: elevate the operative extremity Place 2-3 pillows under affected extremity., Post-op

Equipment

<input type="checkbox"/> Obtain supply / device:	Routine, Once Obtain: Post-op
<input type="checkbox"/> Postop shoe	Routine, Once Left/Right: Sizes: Gender Size: Special Instructions: Post-op
<input type="checkbox"/> Cast shoe	Routine, Once Left/Right: Sizes: Gender Size: Special Instructions: Post-op
<input type="checkbox"/> Cam Walker boot	Routine, Once Left/Right: Sizes: Gender Size: Special Instructions: Post-op

Nursing Assessments

<input checked="" type="checkbox"/> Peripheral vascular assessment	Routine, Per unit protocol Until discharge., Post-op
--	---

Nursing Interventions

<input checked="" type="checkbox"/> Pulse oximetry	Routine, With vitals For 24 Hours Current FIO2 or Room Air: And while on PCA., Post-op
<input checked="" type="checkbox"/> Incentive spirometry	Routine, Every hour For 10 Occurrences Respiratory to instruct., Post-op
<input type="checkbox"/> Intake and output	Routine, Every shift For 48 Hours, Post-op
<input type="checkbox"/> Insert and Maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: If unable to void., Post-op
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S For 48 Hours Orders: to gravity Post-op
<input type="checkbox"/> Remove Foley catheter	Routine, Once Post-op day ***, Post-op

<input checked="" type="checkbox"/> Place antiembolic stockings	Routine, Once Calf sleeve to non operative leg while in bed., PACU & Post-op
<input checked="" type="checkbox"/> Apply ice pack	Routine, Until discontinued, Starting S Affected area: Waking hours only? Nurse to schedule? Special Instructions: To affected extremity., Post-op
<input type="checkbox"/> Wound Care Orders + Medication	
<input type="checkbox"/> Wound care orders	Routine, Daily, Starting S+1 Location: Site: Irrigate wound? Apply: Dressing Type: Post-op
<input type="checkbox"/> wound care medications Please select the appropriate medication for wound care	
<input type="checkbox"/> mupirocin (BACTROBAN) 2 % ointment	Topical, 3 times daily
<input type="checkbox"/> collagenase (SANTYL) ointment	Topical, daily
<input type="checkbox"/> silver sulfadiazine (SILVADENE, SSD) 1 % cream	Topical, daily
<input type="checkbox"/> zinc oxide 20 % ointment	Topical
<input type="checkbox"/> white petrolatum-mineral oil (EUCERIN) topical cream	Topical, PRN, dry skin
<input type="checkbox"/> silver gel	topical (top)
<input type="checkbox"/> gel dressing gel	topical (top)
<input type="checkbox"/> Verify patient has DME equipment	Routine, Once For 1 Occurrences, Post-op
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Care Order Instruction	Routine, Until discontinued, Starting S Dispense patient instructions, Post-op
Notify	
<input type="checkbox"/> Notify Physician (Anesthesia)	Routine, Once For 1 Occurrences, Contact Anesthesia for all issues with Nerve Block., Post-op
<input type="checkbox"/> Notify Hospitalist/Internist of patient's location	Routine, Once For 1 Occurrences, Post-op
Diet	
<input type="checkbox"/> Diet - Clear liquids, advance as tolerated to Regular	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Regular Advance target diet criteria: IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Please assess bowel sounds between progressions., Post-op
<input type="checkbox"/> Diet - Clear liquids, advance as tolerated to Diabetic 1800 Carb Control	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Diabetic 1800 Carb Control Advance target diet criteria: IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Please assess bowel sounds between progressions., Post-op

<input type="checkbox"/> Diet - Clear liquids, advance as tolerated to Heart Healthy	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Heart Healthy Advance target diet criteria: IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Please assess bowel sounds between progressions., Post-op
<input type="checkbox"/> Diet - Clear liquids, advance as tolerated to Renal (80GM Pro, 2-3GM Na, 2-3GM K)	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Renal (80GM Pro, 2-3GM Na, 2-3GM K) Advance target diet criteria: IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Please assess bowel sounds between progressions., Post-op

IV

IV Fluids (Single Response)

<input type="checkbox"/> sodium chloride 0.9 % infusion	75 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> lactated Ringer's infusion	75 mL/hr, intravenous, continuous, Post-op

Medications

IV Antibiotics: For Patients LESS than or EQUAL to 120 kg

<input type="checkbox"/> cefazolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, once, For 1 Doses, Post-op For patients LESS than or EQUAL to 120 kg Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> FOR MRSA CONCERN OR SEVERE PENICILLIN ALLERGY - vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Reason for Therapy: Surgical Prophylaxis

IV Antibiotics: For Patients GREATER than 120 kg

<input type="checkbox"/> cefazolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, once, For 1 Doses, Post-op Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> FOR MRSA CONCERN OR SEVERE PENICILLIN ALLERGY - vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Reason for Therapy: Surgical Prophylaxis

Pain Medications

Check Prescription Drug Monitoring Program.

Prior to initiation of opioid therapy, it is recommended to check the prescription monitoring program (PMP) database to assess patient's opioid tolerance status. A summarized version of the PMP report may be accessed by clicking on the NaRx Score on the patient's Storyboard. You may access the full version of the Texas PMP here."

(<https://texas.pmpaware.net/login>)

Texas PMP

Pain Management Guide

Opioid PCA Conversion to Oral Opioid Regimen

Due to risk of accumulation of toxic metabolite, the use of morphine in patients with renal dysfunction is not recommended. An alternative opioid should be utilized, if possible.

Scheduled Pain Medications (Single Response)

Consider scheduled option if pain source is present and patient unable to reliably communicate needs. Do not order both scheduled and PRN NSAIDs/APAP simultaneously.

acetaminophen (TYLENOL) 500 mg tablet or liquid

"Or" Linked Panel

<input type="checkbox"/>	acetaminophen (TYLENOL) tablet	500 mg, oral, every 6 hours scheduled Use if patient can tolerate oral tablet.
<input type="checkbox"/>	acetaminophen (TYLENOL) liquid	500 mg, oral, every 6 hours scheduled
<input type="checkbox"/>	acetaminophen (TYLENOL) 650 mg tablet or liquid	"Or" Linked Panel
<input type="checkbox"/>	acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours scheduled Use if patient can tolerate oral tablet.
<input type="checkbox"/>	acetaminophen (TYLENOL) liquid	650 mg, oral, every 6 hours scheduled
<input type="checkbox"/>	NSAIDs: For Patients LESS than 65 years old (Single Response)	
<input type="checkbox"/>	ibuprofen (ADVIL, MOTRIN) tablet or oral suspension	"Or" Linked Panel
<input type="checkbox"/>	ibuprofen (ADVIL, MOTRIN) tablet	600 mg, oral, every 6 hours scheduled Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury. Give if patient is able to tolerate oral medication.
<input type="checkbox"/>	ibuprofen (MOTRIN) 100 mg/5 mL suspension	600 mg, oral, every 6 hours scheduled Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
<input type="checkbox"/>	naproxen (NAPROSYN) tablet	250 mg, oral, 2 times daily
<input type="checkbox"/>	celecoxib (CeleBREX) capsule	100 mg, oral, 2 times daily
<input type="checkbox"/>	ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours scheduled For patients LESS THAN 65 years old. Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
<input type="checkbox"/>	NSAIDs: For Patients GREATER than or EQUAL to 65 years old (Single Response)	
<input type="checkbox"/>	ibuprofen (ADVIL, MOTRIN) tablet or oral suspension	"Or" Linked Panel
<input type="checkbox"/>	ibuprofen (ADVIL, MOTRIN) tablet	600 mg, oral, every 6 hours scheduled, Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury. Give if patient is able to tolerate oral medication.
<input type="checkbox"/>	ibuprofen (MOTRIN) 100 mg/5 mL suspension	600 mg, oral, every 6 hours scheduled, Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury. Use if patient cannot swallow tablet.
<input type="checkbox"/>	naproxen (NAPROSYN) tablet	250 mg, oral, 2 times daily, Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
<input type="checkbox"/>	celecoxib (CeleBREX) capsule	100 mg, oral, 2 times daily, Post-op For age GREATER than or EQUAL to 65 yo and patients LESS than 50kg. Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
<input type="checkbox"/>	ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours scheduled, Post-op

PRN Pain Medications

Consider scheduled option if pain source is present and patient unable to reliably communicate needs. Monitor closely for response. Adjust dose for renal/liver function and age. Do not order both scheduled and PRN NSAIDs/APAP simultaneously. Order ONLY one short acting PO and short acting IV simultaneously. Oral option and IV options to be ordered simultaneously.

PRN Oral Medications for Mild Pain (Pain Score 1-3):
For Patients LESS than 65 years old (Single Response)

Do not order both scheduled and PRN NSAIDs/APAP simultaneously.

<input type="checkbox"/>	acetaminophen (TYLENOL) tablet OR oral suspension OR rectal suppository	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/>	acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, mild pain (score 1-3) Give if patient able to swallow tablet.
<input type="checkbox"/>	acetaminophen (TYLENOL)suspension	650 mg, oral, every 6 hours PRN, mild pain (score 1-3) Use if patient cannot tolerate oral tablet.
<input type="checkbox"/>	acetaminophen (TYLENOL) suppository	650 mg, rectal, every 6 hours PRN, mild pain (score 1-3) Use if patient cannot tolerate oral tablet OR oral solution.
<input type="checkbox"/>	ibuprofen (ADVIL, MOTRIN) tablet or oral suspension	"Or" Linked Panel

<input type="checkbox"/>	ibuprofen (ADVIL, MOTRIN) tablet	600 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Give if patient is able to tolerate oral medication.
<input type="checkbox"/>	ibuprofen (MOTRIN) 100 mg/5 mL suspension	600 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Use if patient cannot swallow tablet.
<input type="checkbox"/>	naproxen (NAPROSYN) tablet	250 mg, oral, every 8 hours PRN, mild pain (score 1-3), Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
<input type="checkbox"/>	celecoxib (CeleBREX) capsule	100 mg, oral, 2 times daily PRN, mild pain (score 1-3), Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
<input type="checkbox"/>	ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours PRN, mild pain (score 1-3) Give if patient unable to swallow tablet.
<input type="checkbox"/> PRN Oral Medications for Mild Pain (Pain Score 1-3): For Patients GREATER than or EQUAL to 65 years old (Single Response)		
Consider scheduled option if pain source is present and patient unable to reliably communicate needs. Monitor closely for response. Adjust dose for renal/liver function and age. Do not order both scheduled and PRN NSAIDs/APAP simultaneously. Order ONLY one short acting PO and short acting IV simultaneously. Oral option and IV options to be ordered simultaneously.		
<input type="checkbox"/>	acetaminophen (TYLENOL) tablet OR oral suspension	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/>	acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, mild pain (score 1-3) Use if patient can tolerate oral tablet.
<input type="checkbox"/>	acetaminophen (TYLENOL)suspension	650 mg, oral, every 6 hours PRN, mild pain (score 1-3) Use if patient cannot tolerate oral tablet.
<input type="checkbox"/> PRN Oral Medications for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)		
<input type="checkbox"/>	acetaminophen-codeine (TYLENOL #3) tablet OR elixir	"Or" Linked Panel
<input type="checkbox"/>	acetaminophen-codeine (TYLENOL WITH CODEINE #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication. The use of codeine-containing products is contraindicated in patients LESS THAN 12 years of age. Is this patient OVER 12 years of age? Y/N:
<input type="checkbox"/>	acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet. The use of codeine-containing products is contraindicated in patients LESS THAN 12 years of age. Is this patient OVER 12 years of age? Y/N:
<input type="checkbox"/>	HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/>	HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient able to swallow tablet.
<input type="checkbox"/>	HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient unable to swallow tablet.
<input type="checkbox"/>	oxyCODONE (ROXICODONE) immediate release tablet	5 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Tablets may be crushed. Give if patient able to swallow tablet
<input type="checkbox"/>	traMADoL (ULTRAM) tablet	50 mg, oral, every 6 hours PRN, moderate pain (score 4-6) Max daily dose 200 mg/day in patients with CrCl < 30 ml/min. Give if patient able to swallow tablet.

<input type="checkbox"/> PRN Oral Medications for Moderate Pain (Pain Score 4-6): For Patients GREATER than or EQUAL to 65 years old (Single Response)	
<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) tablet OR elixir	"Or" Linked Panel
<input type="checkbox"/> acetaminophen-codeine (TYLENOL WITH CODEINE #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication. The use of codeine-containing products is contraindicated in patients LESS THAN 12 years of age. Is this patient OVER 12 years of age? Y/N:
<input type="checkbox"/> acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet. The use of codeine-containing products is contraindicated in patients LESS THAN 12 years of age. Is this patient OVER 12 years of age? Y/N:
<input type="checkbox"/> HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	"Or" Linked Panel
Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)	
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient able to swallow tablet.
<input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient unable to swallow tablet.
<input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet	2.5 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Tablets may be crushed. Give if patient able to swallow tablet
<input type="checkbox"/> traMADoL (ULTRAM) tablet	25 mg, oral, every 6 hours PRN, moderate pain (score 4-6) Max daily dose 300mg in patients age GREATER THAN 75 years or 200 mg/day in patients with CrCl < 30 ml/min. Give if patient able to swallow tablet.
<input type="checkbox"/> PRN IV Medications for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old if unable to tolerate Oral Pain Medication. (Single Response)	
Due to risk of toxicity, the use of morphine products in patients with renal dysfunction, particularly in ESRD, is not recommended. An alternative opioid should be utilized.	
<input type="checkbox"/> morPHINE injection	2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op Give if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications.
<input type="checkbox"/> hydromorPHONE (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op Give if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications
<input type="checkbox"/> ketorolac (TORADOL) IV (Single Response)	Do NOT use in patients with eGFR LESS than 30 mL/min. WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery.
<input type="checkbox"/> For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), Post-op Use if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications
<input type="checkbox"/> PRN IV Medications for Moderate Pain (Pain Score 4-6): For Patients GREATER than or EQUAL to 65 years old if unable to tolerate Oral Pain Medication. (Single Response)	

Due to risk of toxicity, the use of morphine products in patients with renal dysfunction, particularly in ESRD, is not recommended. An alternative opioid should be utilized. (adjust dose for renal/liver function and age)

- | | |
|--|--|
| () morPHINE injection | 2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications. |
| () hydromorPHONE (DILAUDID) injection | 0.2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications |

[] PRN Oral Medications for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)

Due to risk of toxicity, the use of morphine products in patients with renal dysfunction, particularly in ESRD, is not recommended. An alternative opioid should be utilized.

- | | |
|--|--|
| () HYDROcodone-acetaminophen 10/325 (NORCO) tablet
OR elixir | "Or" Linked Panel
Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) |
| [] HYDROcodone-acetaminophen (NORCO) 10-325 mg per tablet | 1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient able to swallow tablet. |
| [] HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution | 20 mL, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient unable to swallow tablet. |
| () morPHINE immediate-release tablet | 15 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Tablets may be crushed. Give if patient able to swallow tablet |
| () oxyCODONE (ROXICODONE) immediate release tablet | 10 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Tablets may be crushed. Give if patient able to swallow tablet |

[] PRN Oral Medications for Severe Pain (Pain Score 7-10): For Patients GREATER than or EQUAL to 65 years old (Single Response)

Due to risk of toxicity, the use of morphine products in patients with renal dysfunction, particularly in ESRD, is not recommended. An alternative opioid should be utilized.

- | | |
|---|---|
| () oxyCODONE (ROXICODONE) immediate release tablet | 5 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Oral tablets may be crushed. Give if patient able to swallow tablet |
| () morPHINE immediate-release tablet | 7.5 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Oral tablets may be crushed. Give if patient able to swallow tablets. |
| () HYDROcodone-acetaminophen 7.5/325 (NORCO) tablet
OR elixir | "Or" Linked Panel
Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) |
| [] HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet | 1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient able to swallow tablet. |
| [] HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution | 10 mL, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient unable to swallow tablet. |
| () HYDROcodone-acetaminophen 10/325 (NORCO) tablet
OR elixir | "Or" Linked Panel
Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) |
| [] HYDROcodone-acetaminophen (NORCO) 10-325 mg per tablet | 1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient able to swallow tablet. |
| [] HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution | 20 mL, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient unable to swallow tablet. |

traMADoL (ULTRAM) tablet 50 mg, oral, every 6 hours PRN, severe pain (score 7-10)
Max daily dose 300mg in patients age GREATER THAN 75 years or 200 mg/day in patients with CrCl < 30 ml/min. Give if patient able to swallow tablet.

PRN IV Medications for Severe Pain (Pain Score 7-10):
For Patients LESS than 65 years old if unable to tolerate Oral Pain Medication. (Single Response)

Due to risk of toxicity, the use of morphine products in patients with renal dysfunction, particularly in ESRD, is not recommended. An alternative opioid should be utilized.

fentaNYL (SUBLIMAZE) injection 25 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10),
Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain 60 minutes after giving oral pain medications.

morPHINE injection 4 mg, intravenous, every 4 hours PRN, severe pain (score 7-10),
Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications.

hydromorPHONE (DILAUDID) injection 0.8 mg, intravenous, every 3 hours PRN, severe pain (score 7-10),
Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain 60 minutes after giving oral pain medications.

PRN IV Medications for Severe Pain (Pain Score 7-10):
For Patients GREATER than or EQUAL to 65 years old if unable to tolerate Oral Pain Medication. (Single Response)

Due to risk of toxicity, the use of morphine products in patients with renal dysfunction, particularly in ESRD, is not recommended. An alternative opioid should be utilized.

fentaNYL (SUBLIMAZE) injection 12.5 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10),
Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain 60 minutes after giving oral pain medications.

morPHINE injection 2 mg, intravenous, every 4 hours PRN, severe pain (score 7-10),
Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain unrelieved 60 minutes after giving oral pain medications.

hydromorPHONE (DILAUDID) injection 0.5 mg, intravenous, every 4 hours PRN, severe pain (score 7-10),
Post-op
Give if patient is NPO, unable to swallow oral medication, or if pain 60 minutes after giving oral pain medications.

PCA Medications - Not HMSJ (Single Response)

fentaNYL PCA (SUBLIMAZE) 1500 mcg/30 mL +
Nursing PCA Orders

fentaNYL PCA (SUBLIMAZE) 1500 mcg/30 mL (Single Response)

fentaNYL (SUBLIMAZE) 1500 mcg/30 mL
PCA solution for Opioid Naive
Nurse Loading Dose: Not Ordered
PCA Dose: 10 mcg
Lockout: 10 Minutes
Four Hour Dose Limit: 150 mcg intravenous, continuous
For breakthrough pain: Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. RN may bolus *** every *** hours as needed. If pain persists, may increase PCA demand dose by *** mcg ONCE. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber.

Adjust doses for age, renal function or other factors.

Nursing PCA Orders

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change
<input type="checkbox"/> PCA Documentation	Routine, Every 12 hours At the beginning (or end of each shift), prior to clearing PCA pump data, the following must be documented: doses delivered, number of attempts, total amount of medication infused (in mg or mcg), and volume remaining in syringe (residual volume).
<input type="checkbox"/> Patient education Pain pump	Routine, Once, Starting S For 1 Occurrences Patient/Family: Education for: Pain pump Provide patient education on appropriate use of PCA including no PCA by proxy. Only the patient may press the dosing button.
<input type="checkbox"/> Pasero Opioid-induced Sedation Scale	Routine, Every 6 hours, Starting S Assess POSS while patient has an active PCA order. Contact provider if score 3 or 4.
<input type="checkbox"/> Notify Physician	Routine, Until discontinued, Starting S, - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy
<input type="checkbox"/> Stop the PCA pump and call ordering physician and/or CERT team for any of the following:	Routine, Until discontinued, Starting S, - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention
<input type="checkbox"/> IV Fluids for provision of PCA Therapy (Single Response)	
<input type="checkbox"/> sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous
<input type="checkbox"/> dextrose 5% infusion	30 mL/hr, intravenous, continuous
<input type="checkbox"/> hydromorPHONE PCA (DILAUDID) 15 mg/30 mL + Nursing PCA Orders	
<input type="checkbox"/> hydromorPHONE PCA (DILAUDID) 15 mg/30 mL (Single Response)	
<input type="checkbox"/> hydromorPHONE (DILAUDID) 15 mg/30 mL in sodium chloride 0.9% PCA for Opioid Naive	Nurse Loading Dose: Not Ordered PCA Dose: 0.2 mg Lockout: 10 Minutes MAX (Four hour dose limit): 3 mg intravenous, continuous
	For breakthrough pain: Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. RN may bolus *** every *** hours as needed. If pain persists, may increase PCA demand dose by *** mg ONCE. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber.
	Adjust doses for age, renal function or other factors.
<input type="checkbox"/> Nursing PCA Orders	
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

[] PCA Documentation	Routine, Every 12 hours At the beginning (or end of each shift), prior to clearing PCA pump data, the following must be documented: doses delivered, number of attempts, total amount of medication infused (in mg or mcg), and volume remaining in syringe (residual volume).
[] Patient education Pain pump	Routine, Once, Starting S For 1 Occurrences Patient/Family: Education for: Pain pump Provide patient education on appropriate use of PCA including no PCA by proxy. Only the patient may press the dosing button.
[] Pasero Opioid-induced Sedation Scale	Routine, Every 6 hours, Starting S Assess POSS while patient has an active PCA order. Contact provider if score 3 or 4.
[] Notify Physician	Routine, Until discontinued, Starting S, - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy
[] Stop the PCA pump and call ordering physician and/or CERT team for any of the following:	Routine, Until discontinued, Starting S, - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention
[] IV Fluids for provision of PCA Therapy (Single Response)	
() sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous
() dextrose 5% infusion	30 mL/hr, intravenous, continuous
() morPHINE PCA 30 mg/30 mL + Nursing PCA Orders	
[] morPHINE PCA 30 mg/30 mL (Single Response)	
() morPHINE PCA 30 mg/30 mL in sodium chloride 0.9% for Opioid Naive	Nurse Loading Dose: Not Ordered PCA Dose: 1 mg Lockout Interval: 10 Minutes MAX (Four hour dose limit): 20 mg intravenous, continuous For breakthrough pain: Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. RN may bolus *** every *** hours as needed. If pain persists, may increase PCA demand dose by *** mg ONCE. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. Adjust doses for age, renal function or other factors.
[] Nursing PCA Orders	
[] Vital signs - T/P/R/BP	Routine, Per unit protocol - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change
[] PCA Documentation	Routine, Every 12 hours At the beginning (or end of each shift), prior to clearing PCA pump data, the following must be documented: doses delivered, number of attempts, total amount of medication infused (in mg or mcg), and volume remaining in syringe (residual volume).
[] Patient education Pain pump	Routine, Once, Starting S For 1 Occurrences Patient/Family: Education for: Pain pump Provide patient education on appropriate use of PCA including no PCA by proxy. Only the patient may press the dosing button.

<input type="checkbox"/> Pasero Opioid-induced Sedation Scale	Routine, Every 6 hours, Starting S Assess POSS while patient has an active PCA order. Contact provider if score 3 or 4.
<input type="checkbox"/> Notify Physician	Routine, Until discontinued, Starting S, - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy
<input type="checkbox"/> Stop the PCA pump and call ordering physician and/or CERT team for any of the following:	Routine, Until discontinued, Starting S, - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

IV Fluids for provision of PCA Therapy (Single Response)

<input type="checkbox"/> sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous
<input type="checkbox"/> dextrose 5% infusion	30 mL/hr, intravenous, continuous

PCA Medications - HMSJ Only (Single Response)

fentaNYL PCA (SUBLIMAZE) 1500 mcg/30 mL + Nursing PCA Orders

fentaNYL PCA (SUBLIMAZE) 1500 mcg/30 mL (Single Response)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) 1500 mcg/30 mL PCA solution for Opioid Naive	Nurse Loading Dose: Not Ordered PCA Dose: 10 mcg Lockout: 10 Minutes Four Hour Dose Limit: 150 mcg intravenous, continuous For breakthrough pain: Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. RN may bolus *** every *** hours as needed. If pain persists, may increase PCA demand dose by *** mcg ONCE. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber.
--	--

Adjust doses for age, renal function or other factors.

Nursing PCA Orders

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change
---	--

PCA Documentation

Routine, Every 12 hours
At the beginning (or end of each shift), prior to clearing PCA pump data, the following must be documented: doses delivered, number of attempts, total amount of medication infused (in mg or mcg), and volume remaining in syringe (residual volume).

Patient education Pain pump

Routine, Once, Starting S For 1 Occurrences
Patient/Family:
Education for: Pain pump
Provide patient education on appropriate use of PCA including no PCA by proxy. Only the patient may press the dosing button.

Pasero Opioid-induced Sedation Scale

Routine, Every 6 hours, Starting S
Assess POSS while patient has an active PCA order. Contact provider if score 3 or 4.

[] Notify Physician	<p>Routine, Until discontinued, Starting S, - PCA pump infusion discontinued for any reason</p> <ul style="list-style-type: none"> - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy
[] Stop the PCA pump and call ordering physician and/or CERT team for any of the following:	<p>Routine, Until discontinued, Starting S, - Respiratory rate 10 per minute or less</p> <ul style="list-style-type: none"> - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention
[] IV Fluids for provision of PCA Therapy (Single Response)	
() sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous
() dextrose 5% infusion	30 mL/hr, intravenous, continuous
() hydromorPHONE PCA (DILAUDID) 30 mg/30 mL + Nursing PCA Orders	
[] hydromorPHONE PCA (DILAUDID) 30 mg/30 mL (Single Response)	
() hydromorPHONE (DILAUDID) 30 mg/30 mL in sodium chloride 0.9% PCA for Opioid Naive	<p>Nurse Loading Dose: Not Ordered
PCA Dose: 0.2 mg
Lockout Interval: 10 Minutes
MAX (Four hour dose limit): 3 mg intravenous, continuous</p> <p>For breakthrough pain: Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. RN may bolus *** every *** hours as needed. If pain persists, may increase PCA demand dose by *** mg ONCE. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber.</p> <p>Adjust doses for age, renal function or other factors.</p>
[] Nursing PCA Orders	
[] Vital signs - T/P/R/BP	<p>Routine, Per unit protocol</p> <ul style="list-style-type: none"> - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change
[] PCA Documentation	<p>Routine, Every 12 hours</p> <p>At the beginning (or end of each shift), prior to clearing PCA pump data, the following must be documented: doses delivered, number of attempts, total amount of medication infused (in mg or mcg), and volume remaining in syringe (residual volume).</p>
[] Patient education Pain pump	<p>Routine, Once, Starting S For 1 Occurrences</p> <p>Patient/Family: Education for: Pain pump</p> <p>Provide patient education on appropriate use of PCA including no PCA by proxy. Only the patient may press the dosing button.</p>
[] Pasero Opioid-induced Sedation Scale	<p>Routine, Every 6 hours, Starting S</p> <p>Assess POSS while patient has an active PCA order. Contact provider if score 3 or 4.</p>
[] Notify Physician	<p>Routine, Until discontinued, Starting S, - PCA pump infusion discontinued for any reason</p> <ul style="list-style-type: none"> - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

[] Stop the PCA pump and call ordering physician and/or CERT team for any of the following:	<p>Routine, Until discontinued, Starting S, - Respiratory rate 10 per minute or less</p> <ul style="list-style-type: none"> - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention
[] IV Fluids for provision of PCA Therapy (Single Response)	
() sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous
() dextrose 5% infusion	30 mL/hr, intravenous, continuous
() morPHINE PCA 30 mg/30 mL + Nursing PCA Orders	
[] morPHINE PCA 30 mg/30 mL (Single Response)	<p>Nurse Loading Dose: Not Ordered
PCA Dose: 1 mg
Lockout Interval: 10 Minutes
MAX (Four hour dose limit): 20 mg intravenous, continuous</p> <p>For breakthrough pain: Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. RN may bolus *** every *** hours as needed. If pain persists, may increase PCA demand dose by *** mg ONCE. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber.</p> <p>Adjust doses for age, renal function or other factors.</p>
[] Nursing PCA Orders	
[] Vital signs - T/P/R/BP	<p>Routine, Per unit protocol</p> <ul style="list-style-type: none"> - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change
[] PCA Documentation	<p>Routine, Every 12 hours</p> <p>At the beginning (or end of each shift), prior to clearing PCA pump data, the following must be documented: doses delivered, number of attempts, total amount of medication infused (in mg or mcg), and volume remaining in syringe (residual volume).</p>
[] Patient education Pain pump	<p>Routine, Once, Starting S For 1 Occurrences</p> <p>Patient/Family: Education for: Pain pump Provide patient education on appropriate use of PCA including no PCA by proxy. Only the patient may press the dosing button.</p>
[] Pasero Opioid-induced Sedation Scale	<p>Routine, Every 6 hours, Starting S</p> <p>Assess POSS while patient has an active PCA order. Contact provider if score 3 or 4.</p>
[] Notify Physician	<p>Routine, Until discontinued, Starting S, - PCA pump infusion discontinued for any reason</p> <ul style="list-style-type: none"> - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy
[] Stop the PCA pump and call ordering physician and/or CERT team for any of the following:	<p>Routine, Until discontinued, Starting S, - Respiratory rate 10 per minute or less</p> <ul style="list-style-type: none"> - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention
[] IV Fluids for provision of PCA Therapy (Single Response)	
() sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous
() dextrose 5% infusion	30 mL/hr, intravenous, continuous

Respiratory Depression and Somnolence

[X] naloxone (NARCAN) injection	0.2 mg, intravenous, once PRN, respiratory depression, as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3), Post-op Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.
---------------------------------	--

Antiemetics - HMSL, HMWB Only

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
[X] promethazine (PHENERGAN) injection	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMM, HMSJ, HMW, HMSTC Only

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
[X] promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSTJ Only

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel

<input checked="" type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Laxatives

<input type="checkbox"/> sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet	2 tablet, oral, nightly PRN, constipation, Post-op
<input type="checkbox"/> magnesium hydroxide suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR WORSE	30 mL, oral, every 6 hours PRN, constipation, Post-op Do not give if patient is on hemodialysis or is in chronic renal failure.
<input type="checkbox"/> bisacodyl (DULCOLAX) EC tablet	10 mg, oral, daily PRN, constipation, Post-op
<input type="checkbox"/> bisacodyl (DULCOLAX) suppository	10 mg, rectal, daily PRN, constipation, Post-op
<input type="checkbox"/> polyethylene glycol (MIRALAX) packet	17 g, oral, daily PRN, constipation, Post-op
<input type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily PRN, constipation, Post-op

Symptom Management

<input type="checkbox"/> pantoprazole (PROTONIX) EC tablet	40 mg, oral, daily at 0600, Post-op Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, headaches, fever, Temperature GREATER than 101 F., Post-op
<input type="checkbox"/> alum-mag hydroxide-simeth (MAALOX) 200-200-20 mg/5 mL suspension	30 mL, oral, every 6 hours PRN, indigestion, heartburn, Post-op
<input type="checkbox"/> magnesium hydroxide OR magnesium citrate (Single Response)	
() magnesium hydroxide suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR WORSE	30 mL, oral, every 6 hours PRN, constipation, Post-op Not Recommended for Chronic Kidney Disease Stage 3 Or Worse
() magnesium citrate solution - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR WORSE	300 mL, oral, once PRN, constipation, Post-op Not Recommended for Chronic Kidney Disease Stage 3 Or Worse

Itching: For Patients LESS than 70 years old (Single Response)

() diphenhydrAMINE (BENADRYL) tablet	25 mg, oral, every 6 hours PRN, itching, Post-op
() hydrOXYzine (ATARAX) tablet	10 mg, oral, every 6 hours PRN, itching, Post-op
() cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching, Post-op
() fexofenadine (ALLEGRA) tablet - For eGFR LESS than 80 mL/min, reduce frequency to once daily as needed	60 mg, oral, 2 times daily PRN, itching, Post-op

Itching: For Patients between 70-76 years old (Single Response)

() cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching, Post-op
--------------------------------	---

Itching: For Patients GREATER than 77 years old (Single Response)

() cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching, Post-op
--------------------------------	---

Insomnia: For Patients LESS than 70 years old (Single Response)

() zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep, Post-op
() ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep, Post-op

Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)

() ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep, Post-op
--------------------------------	---

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

VTE/DVT Risk Definitions

URL:

"\appt1\epicappprod\Restricted\OrderSets\VTEDEVTRISK DEFINITIONS.pdf"

URL:

"https://formweb.com/files/houstonmethodist/documents/COVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf"

[Anticoagulation Guide for COVID patients](#)

() Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)

() Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)

Moderate risk of VTE Routine, Once, PACU & Post-op

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op

Place sequential compression device (Single Response)

() Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

() Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

() Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)

Moderate risk of VTE Routine, Once, PACU & Post-op

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op

Place sequential compression device (Single Response)

() Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

() Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

() High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op

Place sequential compression device (Single Response)

() Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

() Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

() High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	LOW Risk of DVT (Selection Required)	
	Low Risk Definition Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/>	Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/>	Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/>	MODERATE Risk of DVT - Surgical (Selection Required)	
	Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/>	Moderate Risk (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	

<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
[] Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis - **"And" Linked Panel**
Order Sequential compression device

Contraindications exist for pharmacologic prophylaxis
Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous
Routine, Continuous, PACU & Post-op

Contraindications exist for pharmacologic prophylaxis **"And" Linked Panel**
AND mechanical prophylaxis

Contraindications exist for pharmacologic prophylaxis
Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Contraindications exist for mechanical prophylaxis
Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe
40 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op
Indication(s): VTE Prophylaxis

patients with CrCL LESS than 30 mL/min
30 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op
For Patients with CrCL LESS than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min
30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min
40 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection
2.5 mg, subcutaneous, daily, PACU & Post-op
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min
This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection
5,000 Units, subcutaneous, every 8 hours, PACU & Post-op

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)
5,000 Units, subcutaneous, every 12 hours, PACU & Post-op
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg
7,500 Units, subcutaneous, every 8 hours, PACU & Post-op
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet
oral, daily at 1700, PACU & Post-op
Indication:

Pharmacy consult to manage warfarin (COUMADIN)
STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis
Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous
Routine, Continuous, PACU & Post-op

HIGH Risk of DVT - Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once

High Risk Pharmacological Prophylaxis - Surgical Patient
(Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin for VTE Prophylaxis (Single Response)

enoxaparin (LOVENOX) 30 mg daily at 1700 30 mg, subcutaneous, daily at 1700
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) 30 mg every 12 hours 30 mg, subcutaneous, every 12 hours
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) 40 mg daily at 1700 40 mg, subcutaneous, daily at 1700
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) 40 mg every 12 hours 40 mg, subcutaneous, every 12 hours
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1
If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, Starting S+1
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet oral, daily at 1700, Starting S+1
Indication:

Pharmacy consult to manage warfarin (COUMADIN) STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):

Place/Maintain sequential compression device continuous Routine, Continuous

HIGH Risk of DVT - Non-Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op
Indication(s): VTE Prophylaxis

patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op
For Patients with CrCL LESS than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min 40 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, PACU & Post-op
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.
This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, PACU & Post-op

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, PACU & Post-op
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, PACU & Post-op
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet oral, daily at 1700, PACU & Post-op
Indication:

Pharmacy consult to manage warfarin (COUMADIN) STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Hip or Knee
(Arthroplasty) Surgical Patient (Single Response)
(Selection Required)

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

aspirin chewable tablet 162 mg, oral, daily, Starting S+1, PACU & Post-op

aspirin (ECOTRIN) enteric coated tablet 162 mg, oral, daily, Starting S+1, PACU & Post-op

Apixaban and Pharmacy Consult (Selection Required)

apixaban (ELIQUIS) tablet 2.5 mg, oral, 2 times daily, Starting S+1, PACU & Post-op
Indications: VTE prophylaxis

Pharmacy consult to monitor apixaban (ELIQUIS) therapy STAT, Until discontinued, Starting S
Indications: VTE prophylaxis

enoxaparin (LOVENOX) injection (Single Response)
(Selection Required)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) syringe 30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op
For Patients with CrCL LESS than 30 mL/min.
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
Indication(s): VTE Prophylaxis

enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min 40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
Indication(s): VTE Prophylaxis

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op
If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min
This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), PACU & Post-op Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

VTE/DVT Risk Definitions

URL:

"\\appt1\epicappprod\Restricted\OrderSets\VTEDEVTRISK DEFINITIONS.pdf"

[Anticoagulation Guide for COVID patients](#)

URL:

"https://formweb.com/files/houstonmethodist/documents/C OVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf"

<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)	
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op

<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/>	High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/>	High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	LOW Risk of DVT (Selection Required)	
	Low Risk Definition	
	Age less than 60 years and NO other VTE risk factors	
<input type="checkbox"/>	Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/>	Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/>	MODERATE Risk of DVT - Surgical (Selection Required)	
	Moderate Risk Definition	
	Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	
	One or more of the following medical conditions:	
	CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	
	Age 60 and above	
	Central line	
	History of DVT or family history of VTE	
	Anticipated length of stay GREATER than 48 hours	
	Less than fully and independently ambulatory	
	Estrogen therapy	
	Moderate or major surgery (not for cancer)	
	Major surgery within 3 months of admission	
<input type="checkbox"/>	Moderate Risk (Selection Required)	

<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
()	Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
()	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
()	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
()	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
()	patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
()	patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
()	patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
()	fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
()	heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
()	heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
()	HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
()	warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
()	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/>	Mechanical Prophylaxis (Single Response) (Selection Required)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition	
Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	
One or more of the following medical conditions:	
CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	
Age 60 and above	
Central line	
History of DVT or family history of VTE	
Anticipated length of stay GREATER than 48 hours	
Less than fully and independently ambulatory	
Estrogen therapy	
Moderate or major surgery (not for cancer)	
Major surgery within 3 months of admission	
<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin for VTE Prophylaxis (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg daily at 1700	30 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 30 mg every 12 hours	30 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg daily at 1700	40 mg, subcutaneous, daily at 1700 Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) 40 mg every 12 hours	40 mg, subcutaneous, every 12 hours Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, Starting S+1 For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S, PACU & Post-op For Patients with CrCL LESS than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op

<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1, PACU & Post-op Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1, PACU & Post-op For Patients with CrCL LESS than 30 mL/min. Indication(s): VTE Prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min. Indication(s): VTE Prophylaxis

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1, PACU & Post-op For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min Indication(s): VTE Prophylaxis
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op For patients with weight GREATER than 100 kg.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), PACU & Post-op Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

Labs

Labs POD #1

<input type="checkbox"/> CBC with platelet and differential	AM draw repeats For 2 Occurrences, Post-op
<input type="checkbox"/> Basic metabolic panel	AM draw repeats For 2 Occurrences, Post-op
<input type="checkbox"/> Prothrombin time with INR	AM draw repeats For 2 Occurrences, Post-op

Cardiology

Diagnostic Imaging

Imaging

<input type="checkbox"/> XR Tibia Fibula 2 Vw Left	Routine, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences, Post-op
<input type="checkbox"/> XR Tibia Fibula 2 Vw Right	Routine, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences, Post-op
<input type="checkbox"/> XR Ankle 3 Vw Bilateral	Routine, 1 time imaging, Starting S at 1:00 AM For 1 , Post-op
<input type="checkbox"/> XR Ankle 3+ Vw Left	Routine, 1 time imaging, Starting S at 1:00 AM For 1 , Post-op
<input type="checkbox"/> XR Ankle 3+ Vw Right	Routine, 1 time imaging, Starting S at 1:00 AM For 1 , Post-op

<input type="checkbox"/> XR Foot 3 Vw Bilateral	Routine, 1 time imaging, Starting S at 1:00 AM For 1 , Post-op
<input type="checkbox"/> XR Foot 3+ Vw Left	Routine, 1 time imaging, Starting S at 1:00 AM For 1 , Post-op
<input type="checkbox"/> XR Foot 3+ Vw Right	Routine, 1 time imaging, Starting S at 1:00 AM For 1 , Post-op

Other Diagnostic Studies

Respiratory

Rehab

Consults

Ancillary Consult

<input type="checkbox"/> Consult to Case Management	Consult Reason: Discharge Planning Post-op, And post discharge equipment needs. Plan discharge on POD #2-3.
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Discharge Placement Post-op, And post discharge equipment needs. Plan to discharge on POD #2-3.
<input checked="" type="checkbox"/> Consult PT eval and treat	Special Instructions: evaluate equipment needs at discharge Weight Bearing Status: Post-op
<input type="checkbox"/> Consult OT eval and treat	Special Instructions: Instruct on use of hip kit. Weight Bearing Status: Post-op
<input type="checkbox"/> Consult to Fracture Liaison Service	Clinical Indications: Post-op