

## Nursing

## Vitals

[X] Vital signs - T/P/R/BP (for Preservative Free Morphine Administration)	<p>Routine, Every hour For 24 Hours Nurse to reschedule vitals:</p> <p>For Obstetrical patients, follow Post Cesarean Section vital sign orders for the first two (2) hours.</p> <p>-Every 1 hour for 12 times (First 12 hours)</p> <p>-Then, every 2 hours for 6 times (Next 12 hours)</p> <p>-Followed by floor protocol (subsequent hours), PACU &amp; Post-op</p>
[X] Pulse oximetry	<p>Routine, Every hour For 24 Hours Current FIO2 or Room Air: Current FiO2 or Room Air; Nurse to reschedule vitals:</p> <p>-Every 1 hour for 12 times (First 12 hours)</p> <p>-Then, every 2 hours for 6 times (Next 12 hours)</p> <p>-Followed by floor protocol (subsequent hours), PACU &amp; Post-op</p>

## Level of Sedation

[X] Neurological assessment	<p>Routine, Once Assessment to Perform: Level of Consciousness Nurse to reschedule vitals:</p> <p>-Every 30 Minutes for 4 Times (First 2 Hours)</p> <p>-Then, every 1 Hour for 6 Times (Next 6 Hours)</p> <p>-Then, every 2 Hours for 6 Times (Next 12 Hours)</p> <p>-Then, every 4 Hours for 1 Time (Next 4 Hours), PACU &amp; Post-op</p>
-----------------------------	---

## Peripheral Vascular Assessment

[X] Peripheral vascular assessment	<p>Routine, Every 2 hours Until numbness/tingling is resolved., PACU &amp; Post-op</p>
------------------------------------	--

## Nursing

[ ] Straight cath	<p>Routine, Once For 1 Occurrences For URINARY RETENTION: If patient unable to void for 6 hours, one time, PACU &amp; Post-op</p>
[X] Foley catheter care	<p>Routine, Until discontinued, Starting S Orders: Maintain PACU &amp; Post-op</p>

## Notify Anesthesia

[X] Notify Anesthesia

Routine, Until discontinued, Starting S, -Nausea refractory to Promethazine

-Respiratory rate less than 10/min and SpO2 less than 92%

-Itching refractory to Nalbuphine

-Patient not arousable

-URINARY RETENTION: If patient unable to void for 6 hours, PACU & Post-op

## ERAS Pain Medications

### ERAS Multimodal Pain Medications

Click here for ERAS Guidelines from ACOG

URL:

"\appt1\epicappprod\Restricted\OrderSets\Perioperative pathways enhanced recovery after surgery.pdf"

Click here for ERAS Postoperative care guidelines

URL:

"\appt1\epicappprod\Restricted\OrderSets\Guidelines for postoperative care in cesarean delivery.pdf"

<input type="checkbox"/> ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours scheduled, For 4 Doses, PACU & Post-op x4 doses
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	1,000 mg, oral, every 6 hours scheduled, For 4 Doses, PACU & Post-op x4 doses, start 6 hours after last acetaminophen dose
<input type="checkbox"/> oxyCODone (ROXICODONE) IR tablet 5 mg	5 mg, oral, every 4 hours PRN, moderate pain (score 4-6), PACU & Post-op
<input type="checkbox"/> oxyCODone (ROXICODONE) IR tablet 10 mg	10 mg, oral, every 4 hours PRN, severe pain (score 7-10), PACU & Post-op

## Medication

### Multimodal Opioid-Sparing Pain Medications

When selecting pain medications within this section, please be sure to deselect duplicate medications from the pain control section of this order set.

ketorolac (TORADOL) IV (Single Response)  
Start NOW if not already given in O.R. or 6 hours after O.R. dose if given; Do not administer for Creatinine >1.1 or when anticoagulation status contraindicates administration.

ketorolac (TORADOL) 15 mg 15 mg, intravenous, every 6 hours, For 4 Doses

ketorolac (TORADOL) 30 mg 30 mg, intravenous, every 6 hours, For 4 Doses

ibuprofen (MOTRIN, ADVIL) tablet (Single Response)  
Start NOW if not ordering scheduled Ketorolac or 6 hours after last dose of ketorolac if given. Max dose 3200 mg/day

ibuprofen (ADVIL) tablet 800 mg 800 mg, oral, every 8 hours scheduled  
Start 6 hours after last Toradol dose administered.

ibuprofen (ADVIL) tablet 600 mg 600 mg, oral, every 6 hours scheduled  
Start 6 hours after last Toradol dose administered.

acetaminophen (TYLENOL) tablet (Single Response)  
Start NOW or 6 hrs after acetaminophen (OFIRMEV IV) O.R. dose if given  
Total Acetaminophen dose - which includes, IV, PO or combination i.e. Norco, APAP, etc. - should not exceed 4 gms/24 hours or 2 gms/24 hours for patients with cirrhosis or severe hepatic dysfunction

acetaminophen (TYLENOL) tablet 1000 mg 1,000 mg, oral, every 6 hours scheduled  
Start NOW or 6 hrs after Acetaminophen (OFIRMEV IV) O.R. dose if given

acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours scheduled  
Start NOW or 6 hrs after Acetaminophen (OFIRMEV IV) O.R. dose if given

**Antiemetics - HMWB Only**

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
[X] promethazine (PHENERGAN) injection	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

**Antiemetics - HHM Only**

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
[X] promethazine (PHENERGAN) 12.5 mg IV	6.25 mg, intravenous, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

**Antiemetics - HMSJ, HMW, HMSTC, HMTW Only**

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
[X] promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

**Antiemetics - HMSTJ Only**

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is able to tolerate oral medication.

[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IVPB or Oral or Rectal	<b>"Or" Linked Panel</b>
[X] promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	6.25 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Antiemetics - HMSL only

[X] ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
[X] promethazine (PHENERGAN) injection	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Itching - not HMSL

[X] nalbuphine (NUBAIN) injection	2 mg, intravenous, every 2 hour PRN, itching, PACU & Post-op Administer no more than 8 doses in 24 hours
-----------------------------------	---

#### Itching - HMSL Only

[X] nalbuphine (NUBAIN) injection	2 mg, intravenous, every 3 hours PRN, itching, PACU & Post-op Administer no more than 8 doses in 24 hours
-----------------------------------	--

#### Respiratory (Naloxone) - not HMSL

For patients with Respiratory Rate LESS than 8 per minute OR if patient is stuporous or unarousable.

[X] naloxone (NARCAN) injection	0.1 mg, intravenous, every 1 min PRN, respiratory depression, as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3)., PACU & Post-op Repeat Naloxone 0.1 mg once in 2 minutes if necessary (MAXIMUM 0.2 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times. Notify Anesthesia if administered.
---------------------------------	--

#### Respiratory (Naloxone) - HMSL only

For patients with Respiratory Rate LESS than 8 per minute OR if patient is stuporous or unarousable.

<input checked="" type="checkbox"/> naloxone (NARCAN) injection	0.1 mg, intravenous, every 1 min PRN, respiratory depression, as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3), PACU & Post-op Repeat Naloxone 0.1 mg once in 2 minutes if necessary (MAXIMUM 0.2 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times. Notify Anesthesia if administered.
---	---

**Moderate Pain (Pain Score 4-6) (Single Response)**

- ketorolac (TORADOL) IV - do not order if scheduled ketorolac is ordered (Single Response)

Do NOT use in patients with eGFR LESS than 30 mL/min AND/OR patients LESS than 17 years of age.

WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery.

<input type="checkbox"/> For patients ages GREATER than 64 OR weight LESS than 50 kg OR eGFR 30-59 mL/min - ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6) Do not order/administer PRN Ketorolac within the first 24 hours with scheduled Ketorolac. PRN Ketorolac may not be administered within 6 hours of last dose.
<input type="checkbox"/> For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6) Do not order/administer PRN Ketorolac within the first 24 hours with scheduled Ketorolac. PRN Ketorolac may not be administered within 6 hours of last dose.

- oxyCODone (ROXICODONE) (Single Response)

<input type="checkbox"/> oxyCODone (ROXICODONE) immediate release tablet	5 mg, oral, every 4 hours PRN, moderate pain (score 4-6), PACU & Post-op
--	--

**Severe Pain (Pain Score 7-10) (Single Response)**

<input type="checkbox"/> oxyCODone (ROXICODONE) immediate release tablet	10 mg, oral, every 4 hours PRN, severe pain (score 7-10), PACU & Post-op
--	--