Neonate PICC Line Insertion [5478]

Nursing	
X] No blood draws via central line	Routine, Continuous
X] No Blood or Blood Products via PICC/MID line	Routine, Until discontinued, Starting S
X] Notify Physician/Neonatal Nurse Practitioner if PICC/MID line becomes occluded or dressing is soiled or loose	Routine, Until discontinued, Starting S
X] Change dressing- PICC/MID Line	Routine, Per unit protocol Routine dressing changes will be performed only by the designated NICU Team Members
maging	
] XR Chest And Abdomen Child	Routine, 1 time imaging, Starting S at 1:00 AM For 1 Portable- Will call when ready.
] XR Chest 1 Vw Portable	Routine, 1 time imaging, Starting S at 1:00 AM For 1 Portable- Will call when ready.
Medications	
ohexol (OMNIPAQUE) and heparin	0.5 mL, intravenous, once, For 1 Doses
ohexol (OMNIPAQUE) and heparin	For neonatal PICC line placement verification, administer
ohexol (OMNIPAQUE) and heparin iohexol (OMNIPAQUE) 180 mg iodine/mL - For	For neonatal PICC line placement verification, administer
verification of neonatal PICC placement	For neonatal PICC line placement verification, administer contrast to the end of catheter. After X-ray, aspirate contrast.
ohexol (OMNIPAQUE) and heparin] iohexol (OMNIPAQUE) 180 mg iodine/mL - For verification of neonatal PICC placement] heparin (porcine) in 0.45% NaCl 1 unit/mL infusion] heparin (porcine) in 0.45% NaCl for Line Care - One Time Use	For neonatal PICC line placement verification, administer contrast to the end of catheter. After X-ray, aspirate contrast 0.5 mL/hr, intravenous, continuous intra-catheter, once, For 1 Doses
ohexol (OMNIPAQUE) and heparin] iohexol (OMNIPAQUE) 180 mg iodine/mL - For verification of neonatal PICC placement] heparin (porcine) in 0.45% NaCl 1 unit/mL infusion] heparin (porcine) in 0.45% NaCl for Line Care - One	For neonatal PICC line placement verification, administer contrast to the end of catheter. After X-ray, aspirate contrast 0.5 mL/hr, intravenous, continuous intra-catheter, once, For 1 Doses