Nursing

i tarong	
Vital Signs	
[X] Vital Signs	STAT, Per unit protocol
Nursing	
[X] ED bedside monitoring	STAT, Continuous
[X] NIH Stroke Scale	STAT, Once
[X] Neurological assessment	STAT, Per unit protocol
	Assessment to Perform: Glasgow Coma Scale, Level of
	Consciousness,Pupils
[X] Draw labs PRIOR to CT if it will not delay procedure	STAT, Once For 1 Occurrences
[X] Dysphagia screen	STAT, Once
	No oral intake until pass dysphagia screening
[X] No oral intake until pass dysphagia screening	STAT, Once For 1 Occurrences
Notify	
[X] Notify Physician	STAT, Until discontinued, Starting S, If patient presents with risk factors for sepsis, or altered mental status, or abnormal vital signs. Complete ED screening tool and notify ED physician for initiation of sepsis treatment.

IV Fluids

Peripheral IV Access

[X] Initiate and maintain IV	
[X] Insert peripheral IV	Routine, Once
[X] sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
[X] sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

IV Fluids

[] sodium chloride 0.9 % bolus 1000 mL	1,000 mL, intravenous, for 30 Minutes, once, For 1 Doses
[] sodium chloride 0.9 % bolus 500 mL	500 mL, intravenous, for 15 Minutes, once, For 1 Doses
[] sodium chloride 0.9 % infusion - 100 mL/hr	100 mL/hr, intravenous, continuous

Labs

Labs - STAT

[X] CBC and differential	STAT For 1 Occurrences
[X] Partial thromboplastin time	STAT For 1 Occurrences
[X] Prothrombin time with INR	STAT For 1 Occurrences
[X] Comprehensive metabolic panel	STAT For 1 Occurrences
[X] Bedside glucose	STAT, Once
	Perform prior to CT. May use EMS results if available.
[] Lactic acid level, SEPSIS - Now and repeat 2x every 3 hours	Now and repeat 2x every 3 hours For 3 Occurrences
[] Arterial blood gas	STAT For 1 Occurrences
[] Venous blood gas	STAT For 1 Occurrences
[] Hepatic function panel	STAT For 1 Occurrences
[] Lipid panel	STAT For 1 Occurrences
[] Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences
	Specimen Source: Urine
	Specimen Site:
[] Syphilis treponema screen with RPR confirmation	STAT For 1 Occurrences
(reverse algorithm)	Release to patient (Note: If manual release option is selected,
	result will auto release 10 days from finalization.):

Labs-STAT

[X] CBC and differential	STAT For 1 Occurrences
[X] Partial thromboplastin time	STAT For 1 Occurrences
[] Prothrombin time panel I-Stat	STAT For 1 Occurrences
[X] Comprehensive metabolic panel	STAT For 1 Occurrences
[X] Bedside glucose	STAT, Once
	Perform prior to CT. May use EMS results if available.
[] Lactic acid, I-Stat , SEPSIS	Now then every 3 hours For 3 Occurrences
[] Arterial blood gas	STAT For 1 Occurrences
[] Venous blood gas	STAT For 1 Occurrences
[] Hepatic function panel	STAT For 1 Occurrences
[] Lipid panel	STAT For 1 Occurrences
[] Urinalysis	STAT For 1 Occurrences
[] Urine culture	Once For 1 Occurrences, Urine
[] Syphilis treponema screen with RPR confirmation	STAT For 1 Occurrences
(reverse algorithm)	Release to patient (Note: If manual release option is selected,
	result will auto release 10 days from finalization.):

Labs-STAT

[X] CBC and differential	STAT For 1 Occurrences
[X] Partial thromboplastin time	STAT For 1 Occurrences
[] Prothrombin time panel I-Stat	STAT For 1 Occurrences
[X] Comprehensive metabolic panel	STAT For 1 Occurrences
[X] Bedside glucose	STAT, Once
	Perform prior to CT. May use EMS results if available.
[] Lactic acid level, SEPSIS - Now and repeat 2x eve hours	ry 3 Now and repeat 2x every 3 hours For 3 Occurrences
[] Arterial blood gas	STAT For 1 Occurrences
[] Venous blood gas	STAT For 1 Occurrences
[] Hepatic function panel	STAT For 1 Occurrences
[] Lipid panel	STAT For 1 Occurrences
[] Urine Culture and Urinalysis	"And" Linked Panel
[] Urine culture	Once For 1 Occurrences, Urine
[] Urinalysis	STAT For 1 Occurrences
[] Syphilis treponema screen with RPR confirmation (reverse algorithm)	STAT For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.):

Labs - Other

[] Troponin T	STAT For 1 Occurrences	
[] B natriuretic peptide	STAT For 1 Occurrences	
[] TSH	STAT For 1 Occurrences	
[] Ammonia level	STAT For 1 Occurrences	
[] Alcohol level, blood	STAT For 1 Occurrences	
[] Urine drugs of abuse screen	STAT For 1 Occurrences	

Labs-Other

[] Troponin, I-Stat	STAT For 1 Occurrences
[] B natriuretic pep, I-Stat	STAT For 1 Occurrences
[] TSH	STAT For 1 Occurrences
[] Ammonia level	STAT For 1 Occurrences
[] Alcohol level, blood	STAT For 1 Occurrences
[] Urine drugs of abuse screen	STAT For 1 Occurrences

Lab-Other

[] Troponin, I-Stat	STAT For 1 Occurrences	
[] B natriuretic peptide	STAT For 1 Occurrences	
[] TSH	STAT For 1 Occurrences	
[] Ammonia level	STAT For 1 Occurrences	
Drinted on 10/20/2022 at 0.56 AM from SUD		Dege 2 of 4

STAT For 1 Occurrences STAT For 1 Occurrences STAT For 1 Occurrences
STAT For 1 Occurrences
STAT For 1 Occurronces
STAT For 1 Occurrences
STAT For 1 Occurrences STAT For 1 Occurrences
STAT For 1 Occurrences
STAT For 1 Occurrences
Release to patient (Note: If manual release option is selected,
result will auto release 10 days from finalization.): STAT For 1 Occurrences
Release to patient (Note: If manual release option is selected,
result will auto release 10 days from finalization.):
Once For 1 Occurrences
Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.):
"And" Linked Panel
Once, Blood
Collect before antibiotics given. Blood cultures should be ordered x2, wit
each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line
should NEVER be used.
Once, Blood
Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both
sets from a peripheral site, please call the lab for assistance; an IV line
should NEVER be used.
STAT, Once
Clinical Indications: Other:
Other: CVA/TIA/AMS Interpreting Physician:
To be performed by ED Staff - Show immediately to physician
STAT, 1 time imaging, Starting S at 1:00 AM For 1
STAT, 1 time imaging, Starting S at 1:00 AM For 1
Perfusion Brain MRI
STAT, 1 time imaging, Starting S at 1:00 AM For 1 STAT, 1 time imaging, Starting S at 1:00 AM For 1
STAT, 1 time imaging, Starting S at 1:00 AM For 1
STAT, 1 time imaging, Starting S at 1:00 AM For 1
Occurrences
Occurrences STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences

[] CT Stroke Brain Wo Contrast	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences
	If meets stroke protocol criteria, do immediately
X-Ray	
[] Chest 1 Vw Portable	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences
[] Chest 2 Vw	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences
[] Cervical Spine Complete	STAT, 1 time imaging, Starting S at 1:00 AM For 1
Respiratory	
Respiratory	
[X] Oxygen therapy	STAT, Continuous Device: Nasal Cannula Rate in liters per minute: 2 Lpm Titrate to keep O2 Sat Above: Other (Specify)
	Specify titration to keep O2 Sat (%) Above: 94 Indications for O2 therapy: Respiratory distress Device 2:

Device 3:

Indications for O2 therapy: