

## Nursing

## Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol
---	----------------------------

## IV Fluids

## Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

## IV Fluids

<input type="checkbox"/> sodium chloride 0.9 % bolus 1000 mL	1,000 mL, intravenous, for 30 Minutes, once, For 1 Doses
<input type="checkbox"/> sodium chloride 0.9 % bolus 500 mL	500 mL, intravenous, for 15 Minutes, once, For 1 Doses
<input type="checkbox"/> sodium chloride 0.9 % infusion - 100 mL/hr	100 mL/hr, intravenous, continuous

## Medications

## Antiemetics

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet - 4 mg	4 mg, oral, once, For 1 Doses Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection - 4 mg	4 mg, intravenous, once, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV - 12.5 mg	12.5 mg, intravenous, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet - 12.5 mg	12.5 mg, oral, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository - 12.5 mg	12.5 mg, rectal, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

## Antiemetics

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet - 4 mg	4 mg, oral, once PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) injection - 4 mg	4 mg, intravenous, once PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) injection	12.5 mg, intravenous, once PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet - 12.5 mg	12.5 mg, oral, once PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository - 12.5 mg	12.5 mg, rectal, once PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

## Antiemetics

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet - 4 mg	4 mg, oral, once, For 1 Doses Give if patient is able to tolerate oral medication.
---	---

<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection - 4 mg	4 mg, intravenous, once, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB - 12.5 mg	12.5 mg, intravenous, for 30 Minutes, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet - 12.5 mg	12.5 mg, oral, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository - 12.5 mg	12.5 mg, rectal, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

## Labs

### Laboratory STAT

<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/> Reticulocyte count	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> Troponin T	STAT For 1 Occurrences
<input type="checkbox"/> Amylase	STAT For 1 Occurrences
<input type="checkbox"/> Lipase	STAT For 1 Occurrences
<input type="checkbox"/> Lactate dehydrogenase, LDH	STAT For 1 Occurrences
<input type="checkbox"/> Urine drugs of abuse screen	STAT For 1 Occurrences
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences Specimen Source: Urine Specimen Site:

### Laboratory-STAT

<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/> Reticulocyte count	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> Troponin, I-Stat	STAT For 1 Occurrences
<input type="checkbox"/> Amylase	STAT For 1 Occurrences
<input type="checkbox"/> Lipase	STAT For 1 Occurrences
<input type="checkbox"/> Lactate dehydrogenase, LDH	STAT For 1 Occurrences
<input type="checkbox"/> Urine drugs of abuse screen	STAT For 1 Occurrences
<input type="checkbox"/> Urine Culture and Urinalysis	<b>"And" Linked Panel</b>
<input type="checkbox"/> Urine culture	Once For 1 Occurrences, Urine
<input type="checkbox"/> Urinalysis	STAT For 1 Occurrences

### Laboratory - STAT

<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/> Reticulocyte count	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> Troponin, I-Stat	STAT For 1 Occurrences
<input type="checkbox"/> Amylase	STAT For 1 Occurrences
<input type="checkbox"/> Lipase	STAT For 1 Occurrences
<input type="checkbox"/> Lactate dehydrogenase, LDH	STAT For 1 Occurrences
<input type="checkbox"/> Urinalysis	STAT For 1 Occurrences
<input type="checkbox"/> Urine culture	STAT For 1 Occurrences, Urine

### Pregnancy Labs

<input type="checkbox"/> hCG QUALitative, serum	STAT For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.):
---	---

hCG QUALitative, urine STAT For 1 Occurrences  
Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.):

## Microbiology

Blood culture x 2 **"And" Linked Panel**

Blood Culture (Aerobic & Anaerobic) Once, Blood  
Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.

Blood Culture (Aerobic & Anaerobic) Once, Blood  
Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.

## Cardiology

### EKG

ED bedside monitoring STAT, Continuous

ECG 12 lead STAT, STAT For 1 Occurrences  
Clinical Indications: Other:  
Other: Per ACS Protocol  
Interpreting Physician:  
To be performed by ED Staff - Show immediately to MD

## Imaging

### Diagnostics X-Ray

Chest 1 Vw STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences

Chest 2 Vw STAT, 1 time imaging, Starting S at 1:00 AM For 1

Abdomen Acute Inc Chest STAT, 1 time imaging, Starting S at 1:00 AM For 1

## Respiratory

### Respiratory (Single Response)

( ) Oxygen therapy STAT, Continuous  
Device: Nasal Cannula  
Rate in liters per minute: 2 Lpm  
Rate in tenths of a liter per minute:  
O2 %:  
Device 2:  
Device 3:  
Titrate to keep O2 Sat Above: 92%  
Indications for O2 therapy:

## Consults

Additional Consults can be entered below in search field.

### Ancillary Consults

Note: Due to the current resource constraints, consultation orders received after 10 AM Monday - Friday will be seen the following business day.

Day of discharge consults will not be accepted.

Reason for Consult?

Consult Tracking: