

## Nursing

## Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol
---	----------------------------

## Nursing

<input type="checkbox"/> Complete consent for	STAT, Once For 1 Occurrences Procedure: Lumbar Puncture Diagnosis/Condition: Physician: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate?
<input type="checkbox"/> Lumbar puncture tray to bedside	STAT, Once For 1 Occurrences
<input type="checkbox"/> Straight cath	STAT, Once For 1 Occurrences For Urine

## IV Fluids

## IV Fluids

<input type="checkbox"/> dextrose 5%-0.225% sodium chloride infusion	intravenous, continuous
<input type="checkbox"/> dextrose 5%-0.45% sodium chloride infusion	intravenous, continuous
<input type="checkbox"/> sodium chloride 0.9 % bolus	1,000 mL, intravenous, for 60 Minutes, PRN If blood pressure drops by 20 mmHg from baseline and heart rate increases by 20 beats per minute from baseline; release conditional bolus order and notify physician immediately.

## Medications

## Medications

<input type="checkbox"/> acetaminophen (TYLENOL) 160 mg/5 mL (5 mL) solution	oral, once, For 1 Doses
<input type="checkbox"/> acetaminophen (TYLENOL) suppository	325 mg, rectal, once, For 1 Doses
<input type="checkbox"/> acetaminophen (TYLENOL) chewable tablet	160 mg, oral, once, For 1 Doses
<input type="checkbox"/> acetaminophen-codeine 120 mg-12 mg /5 mL (5 mL) solution	10 mL, oral, once, For 1 Doses The use of codeine-containing products is contraindicated in patients LESS THAN 12 years of age. Is this patient OVER 12 years of age? Y/N:
<input type="checkbox"/> ibuprofen (ADVIL,MOTRIN) 100 mg/5 mL suspension	oral, once, For 1 Doses
<input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	10 mL, oral, once, For 1 Doses
<input type="checkbox"/> diphenhydramINE (BENADRYL) injection	intravenous, once, For 1 Doses
<input type="checkbox"/> diphenhydramINE (BENADRYL) 12.5 mg/5 mL elixir	oral, once, For 1 Doses
<input type="checkbox"/> ondansetron (ZOFTRAN) IV	intravenous, once, For 1 Doses
<input type="checkbox"/> ondansetron (ZOFTRAN) oral solution	1 mg, oral, once, For 1 Doses
<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, once, For 1 Doses
<input type="checkbox"/> ranitidine (ZANTAC) 15 mg/mL syrup	oral, once, For 1 Doses Formulary-approved indication:
<input type="checkbox"/> albuterol (PROVENTIL) nebulizer solution	2.5 mg, nebulization, once, For 1 Doses Aerosol Delivery Device:
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection	40 mg, intravenous, once, For 1 Doses
<input type="checkbox"/> dexamethasone (DECADRON) injection	oral, once, For 1 Doses
<input type="checkbox"/> prednisolONE (PRELONE) 15 mg/5 mL syrup	oral, once, For 1 Doses
<input type="checkbox"/> prednisolONE (ORAPRED) 15 mg/5 mL solution	oral, once, For 1 Doses
<input type="checkbox"/> predniSONE (DELTASONE) tablet	5 mg, oral, once, For 1 Doses
<input type="checkbox"/> predniSONE (DELTASONE) tablet	10 mg, oral, once, For 1 Doses

## Labs

### Laboratory - Stat

<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/>	Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/>	Heterophile Ab screen	STAT For 1 Occurrences
<input type="checkbox"/>	Group A strep, rapid antigen	STAT For 1 Occurrences, Throat
<input type="checkbox"/>	RSV, rapid antigen	STAT For 1 Occurrences
<input type="checkbox"/>	Influenza antigen	STAT For 1 Occurrences, Nasopharyngeal
<input type="checkbox"/>	Lactic acid level	STAT For 1 Occurrences
<input type="checkbox"/>	Urinalysis, automated with microscopy	STAT For 1 Occurrences

### Laboratory

<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/>	Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/>	Heterophile Ab screen	STAT For 1 Occurrences
<input type="checkbox"/>	Group A strep, rapid antigen	STAT For 1 Occurrences, Throat
<input type="checkbox"/>	RSV, rapid antigen	STAT For 1 Occurrences
<input type="checkbox"/>	Influenza antigen	STAT For 1 Occurrences, Nasopharyngeal
<input type="checkbox"/>	Lactic acid, I-Stat	STAT For 1 Occurrences
<input type="checkbox"/>	Urinalysis	STAT For 1 Occurrences
<input type="checkbox"/>	Urine culture	STAT For 1 Occurrences, Urine

### Laboratory

<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/>	Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/>	Heterophile Ab screen	STAT For 1 Occurrences
<input type="checkbox"/>	Group A strep, rapid antigen	STAT For 1 Occurrences, Throat
<input type="checkbox"/>	RSV, rapid antigen	STAT For 1 Occurrences
<input type="checkbox"/>	Influenza antigen	STAT For 1 Occurrences, Nasopharyngeal
<input type="checkbox"/>	Lactic acid level	STAT For 1 Occurrences
<input type="checkbox"/>	Urine Culture and Urinalysis	<b>"And" Linked Panel</b>
<input type="checkbox"/>	Urine culture	Once For 1 Occurrences, Urine
<input type="checkbox"/>	Urinalysis	STAT For 1 Occurrences

### Microbiology

<input type="checkbox"/>	Blood culture x 2	<b>"And" Linked Panel</b>
<input type="checkbox"/>	Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
<input type="checkbox"/>	Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
<input type="checkbox"/>	Rotavirus antigen, stool	STAT For 1 Occurrences, Stool
<input type="checkbox"/>	Respiratory pathogen panel with COVID-19 RT-PCR	STAT For 1 Occurrences

## Imaging

### Diagnostic CT

<input type="checkbox"/>	CT Head Wo Contrast	STAT, 1 time imaging, Starting S at 1:00 AM For 1
--------------------------	---------------------	---

### Diagnostic X-Ray

<input type="checkbox"/>	Chest 1 Vw	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences
--------------------------	------------	---

[ ] Chest 2 Vw	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences
[ ] Pelvis And Hips Infant Child 2+ Vw	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Occurrences Please perform Frogview
[ ] Chest And Abdomen Child	STAT, 1 time imaging, Starting S at 1:00 AM For 1 Babygram

## Respiratory

### Respiratory Orders

[ ] Oxygen therapy	Routine, Continuous Device: Titrate to keep O2 Sat Above: Indications for O2 therapy:
[ ] Pulse oximetry	STAT, Daily Current FIO2 or Room Air:
[ ] Consult to Respiratory Therapy	Reason for Consult?