

Nursing

Nursing

<input checked="" type="checkbox"/> Contact isolation status	Open crib in continuous rooming-in OR contact isolation within the nursery. Isolation precautions observed by everyone who handles the baby.
<input checked="" type="checkbox"/> HSV Education	Routine, Once Patient/Family: Family Education for: Other (specify) Specify: Instruct caregivers to recognize signs and symptoms of HSV lesion and prevent newborn contact with potential HSV infection. Instruct on use of eye drops.

Medications

Medications

<input type="checkbox"/> acyclovir (ZOVIRAX) IV syringe	20 mg/kg, intravenous, Administer over: 60 Minutes, every 8 hours Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
<input type="checkbox"/> trifluridine (VIROPTIC) 1 % ophthalmic solution	1 drop, Both Eyes, 4 times daily Start AFTER cultures are obtained. Stop after 5 days if asymptomatic.

Laboratory

Laboratory - HMSJ Only

<input checked="" type="checkbox"/> Miscellaneous referral test - Herpes Culture	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Herpes Culture: Conjunctivae - To be done at 24 hours of life. Send Cultures to TCH Virology Lab.
<input checked="" type="checkbox"/> Miscellaneous referral test - Herpes Culture	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Herpes Culture: Nasopharynx - To be done at 24 hours of life. Send Cultures to TCH Virology Lab.
<input checked="" type="checkbox"/> Miscellaneous referral test - Herpes Culture	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Herpes Culture: Oropharynx - To be done at 24 hours of life. Send Cultures to TCH Virology Lab.
<input checked="" type="checkbox"/> Miscellaneous referral test - Herpes Culture	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Herpes Culture: Rectal swab - To be done at 24 hours of life. Send Cultures to TCH Virology Lab.

Laboratory - All Facilities (Not HMSJ)

<input checked="" type="checkbox"/> Herpes simplex virus by PCR	Once Specimen Source: Other Conjunctivae - To be done at 24 hours of life.
<input checked="" type="checkbox"/> Herpes simplex virus by PCR	Once Specimen Source: Other Nasopharynx - To be done at 24 hours of life.
<input checked="" type="checkbox"/> Herpes simplex virus by PCR	Once Specimen Source: Swab Rectal swab - To be done at 24 hours of life.

