PERIPHERAL Line Antithymocyte (THYMOGLOBULIN) [4510] Nursing Nursing [X] Notify Physician for vitals: Routine, Until discontinued, Starting S Temperature greater than: Temperature less than: Systolic BP greater than: Systolic BP less than: 90 Diastolic BP greater than: Diastolic BP less than: 60 MAP less than: Heart rate greater than (BPM): 110 Heart rate less than (BPM): Respiratory rate greater than: Respiratory rate less than: SpO2 less than: [X] Notify Physician for severe chills Routine, Until discontinued, Starting S Medications **Premedications** [X] acetaminophen (TYLENOL) tablet 650 mg, oral, once, For 1 Doses Give 30 minutes PRIOR to Antithymocyte Globulin infusion. "Or" Linked Panel [X] Premedication: diphenhydramine (BENADRYL) tablet OR injection [X] diphenhydrAMINE (BENADRYL) tablet 25 mg, oral, once, For 1 Doses Give 30 minutes PRIOR to Antithymocyte Globulin infusion. 25 mg, intravenous, once, For 1 Doses [X] diphenhydrAMINE (BENADRYL) injection Give 30 minutes PRIOR to Antithymocyte Globulin infusion. [X] Premedications: Steroids (Single Response) () methylPREDNISolone sodium succinate 40 mg, intravenous, once, For 1 Doses (Solu-MEDROL) injection Give 30 minutes PRIOR to Antithymocyte Globulin infusion. () hydrocortisone sodium succinate intravenous, once, For 1 Doses (Solu-CORTEF) injection Give 30 minutes PRIOR to Antithymocyte Globulin infusion. **Antithymocyte Globulin** [X] antithymocyte globulin (rabbit) (THYMOGLOBULIN) 1.5 1.5 mg/kg, intravenous, Administer over: 6 Hours, once, mg/kg, HEParin (porcine) 2 Units/mL, hydrocortisone Starting H+30 Minutes, For 1 Doses sodium succinate (PF) (Solu-CORTEF) 0.05 mg/mL in sodium chloride 0.9% 250 mL IVPB **Infusion Reaction Management** [X] hydrocortisone sodium succinate (Solu-CORTEF) 100 mg, intravenous, once PRN, chills injection [X] acetaminophen (TYLENOL) tablet 325 mg, oral, every 6 hours PRN, fever, for Temperature GREATER than 100 F Labs

AM draw For 1 Occurrences

[X] CBC with platelet and differential

Laboratory in AM