

Retinopathy of Prematurity Exam [714]

Nursing

Nursing

- | | |
|---|---|
| <input type="checkbox"/> Provide equipment / supplies at bedside- Retinopathy | Routine, Once For 1 Occurrences
Supplies: Other (specify)
Other: Have the sterile lid speculum and lens loop at the infant's bedside. |
| <input checked="" type="checkbox"/> Apply phototherapy mask to eyes for 4 hours post ROP exam | Routine, Continuous
Phototherapy: Double
Apply phototherapy mask to eyes 4 hours post dilation. |

Medications

Medications

- | | |
|--|---|
| <input checked="" type="checkbox"/> cyclopentolate-phenylephrine (CYCLOMYDRYL) 0.2-1 % ophthalmic solution | 1 drop, Both Eyes, PRN, instill every 5 minutes for 3 doses
To be administered 45 minutes PRIOR to eye exam. |
| <input checked="" type="checkbox"/> sucrose 24 % (TOOT SWEET) oral solution | 0.2 mL, oral, once PRN, mild pain (score 1-3), pain |
| <input checked="" type="checkbox"/> proparacaine (ALCAINE) 0.5 % ophthalmic solution | 1 drop, Both Eyes, PRN, for eye exam
To be administered 30 seconds PRIOR to eye exam. |
| <input checked="" type="checkbox"/> artificial tears(hypromellose) (SYSTANE/GENTEAL) 0.3 % ophthalmic gel | 1 Application, Both Eyes, PRN, dry eyes, for eye exam
To be administered for each eye exam. |