

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="radio"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="radio"/>	Outpatient observation services under general supervision	Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="radio"/>	Outpatient in a bed - extended recovery	Admitting Physician: Bed request comments: PACU & Post-op
<input type="radio"/>	Admit to Inpatient	Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

@CERMSG(674511:):@

Code Status (Single Response)

DNR and Modified Code orders should be placed by the responsible physician.

- | | |
|--|--|
| <input type="checkbox"/> Full code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Did the patient/surrogate require the use of an interpreter?
Did the patient/surrogate require the use of an interpreter?
Does patient have decision-making capacity?
Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	
<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Did the patient/surrogate require the use of an interpreter? Did the patient/surrogate require the use of an interpreter? Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions ((For use when a patient is NOT in a cardiopulmonary arrest))	I understand that if the patient is NOT in a cardiopulmonary arrest, the selected treatments will NOT be provided. I understand that all other unselected medically indicated treatments will be provided. Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Post-op
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol, Post-op
--	-------------------------------------

Activity

<input type="checkbox"/> Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges Post-op
<input checked="" type="checkbox"/> Up in chair	Routine, Until discontinued, Starting S Specify: Up in chair Additional modifier: This evening, Post-op
<input checked="" type="checkbox"/> Ambulate with assistance	Routine, 4 times daily, Starting S+1 Specify: with assistance Post-op

Nursing Care

<input type="checkbox"/> Abdominal binder	Routine, Once Waking hours only? Nurse to schedule? Special Instructions: Post-op
<input checked="" type="checkbox"/> Encourage deep breathing and coughing	Routine, Every 2 hours while awake Until ambulatory, Post-op
<input checked="" type="checkbox"/> Incentive spirometry	Routine, Every 2 hours while awake Place at bedside. Encourage patient to use., Post-op
<input type="checkbox"/> Intake and output	Routine, Every shift For 24 Hours, Post-op
<input type="checkbox"/> K-pad to bedside	Routine, Until discontinued, Starting S Apply as needed to area of pain, Post-op
<input type="checkbox"/> Saline lock IV	Routine, Continuous, Post-op
<input type="checkbox"/> No other analgesia until PCA is discontinued	Routine, Until discontinued, Starting S, Post-op

Nursing POD 1

<input type="checkbox"/> Remove dressing	Routine, Until discontinued, Starting S+1 Remove abdominal dressing or vaginal pack if present, Post-op
<input type="checkbox"/> Nursing wound care	Routine, Daily, Starting S+1 Location: Site: Irrigate wound? Apply: Dressing Type: Clean incision with water, Post-op
<input type="checkbox"/> Saline lock IV	Routine, Continuous, Starting S+1, Post-op
<input type="checkbox"/> Discontinue IV	Routine, Once, Starting S+1 On POD 1 if patient is afebrile and tolerating diet, Post-op
<input type="checkbox"/> Discontinue PCA on Post-Op day # 1	Routine, Until discontinued, Starting S+1 Prior to discontinuing foley, Post-op
<input checked="" type="checkbox"/> Remove Foley catheter	Routine, Once, Starting S+1 D/C Foley in AM if urine is clear. DO NOT DC FOLEY IF ANTERIOR REPAIR OR BLADDER SURGERY., Post-op
<input type="checkbox"/> Post-op voiding trial	Routine, Once, Starting S+1, Post-op

Notify

<input checked="" type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 101 Temperature less than: Systolic BP greater than: 170 Systolic BP less than: 90 Diastolic BP greater than: 110 Diastolic BP less than: 60 MAP less than: Heart rate greater than (BPM): 120 Heart rate less than (BPM): 60 Respiratory rate greater than: 28 Respiratory rate less than: 10 SpO2 less than: And for urine output less than 30 milliliters per hour
--	--

Diet

<input checked="" type="checkbox"/> NPO except ice chips	Diet effective now, Starting S NPO: Except Ice chips Pre-Operative fasting options: Until no longer nauseated, An NPO order without explicit exceptions means nothing can be given orally to the patient., Post-op
--	---

<input type="checkbox"/> Diet - Clear Liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input checked="" type="checkbox"/> Diet -	Diet effective midnight, Starting S+1 at 12:01 AM Diet(s): Advance Diet as Tolerated? Yes Target Diet: Regular Advance target diet criteria: IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Advance diet as tolerated 12 hours PostOP, Post-op

IV Fluids

IV Fluids

<input checked="" type="checkbox"/> lactated Ringer's infusion	125 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5 % and lactated Ringer's infusion	125 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> sodium chloride 0.9 % infusion	125 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5%-0.9% sodium chloride infusion	125 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5 % and sodium chloride 0.9 % with potassium chloride 20 mEq/L infusion	125 mL/hr, intravenous, at 100 mL/hr, continuous, Post-op

Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

Medications

Antibiotics: cefazolin (ANCEF) for patients LESS than or EQUAL to 120 kg (Single Response)

<input type="checkbox"/> cefazolin (ANCEF) IV	2 g, intravenous, once, For 1 Doses, Post-op Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
---	--

Antibiotics: cefazolin (ANCEF) for patients GREATER than 120 kg (Single Response)

<input type="checkbox"/> cefazolin (ANCEF) IV	3 g, intravenous, once, For 1 Doses, Post-op Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy:
---	--

Antibiotics: if Penicillin or Beta-Lactam Allergic

If patient is Penicillin or Beta-Lactam Allergic: Choose ONE option from Section 1 and ONE option from Section 2. TWO agents MUST be selected for Core Measure compliance.

<input type="checkbox"/> Section 1 (Single Response)	
<input type="checkbox"/> metronidazole (FLAGYL) IV	500 mg, intravenous, once, For 1 Doses, Post-op For penicillin or beta-lactam allergic patients. Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy:
<input type="checkbox"/> clindamycin (CLEOCIN) IV - Recommended ONLY for patients with high risk for penicillin anaphylaxis that are culture isolate sensitive to Clindamycin.	900 mg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Post-op For penicillin or beta-lactam allergic patients. Reason for Therapy:

[] Section 2 (Single Response)

() levofloxacin (LEVAQUIN) IV	500 mg, intravenous, once, For 1 Doses, Post-op For penicillin or beta-lactam allergic patients. Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy:
() gentamicin (GARAMYCIN) IV	5 mg/kg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Post-op For penicillin or beta-lactam allergic patients. Reason for Therapy:

Mild Pain (Pain Score 1-3) (Single Response)

() acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op
() ibuprofen (ADVIL, MOTRIN) tablet - Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury	600 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op
() tramADol (ULTRAM) tablet - Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury	25 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Not recommended in patients with eGFR LESS than 30 mL/min OR in acutekidney injury. Not to exceed 400 mg/day. Allowance for Patient Preference:

NALOXONE FOR OBGYN SURGERY POST OP OPIOID PAIN MEDICATIONS

[X] naloxone (NARCAN) 0.4 mg/mL injection	intravenous, PRN, respiratory depression, opioid reversal, L&D Pre-Delivery
---	---

Moderate Pain (Pain Score 4-6) (Single Response)

() acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	2 tablet, oral, every 3 hours PRN, moderate pain (score 4-6), Post-op The use of codeine-containing products is contraindicated in patients LESS THAN 12 years of age. Is this patient OVER 12 years of age? Y/N: Allowance for Patient Preference:
() HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 4 hours PRN, moderate pain (score 4-6), Post-op Allowance for Patient Preference:
() oxyCODONE-acetaminophen (PERCOCET) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Allowance for Patient Preference:
() ketorolac (TORADOL) tablet - Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury	10 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Not recommended in patients with eGFR LESS than 30 mL/min OR in acutekidney injury
() ketorolac (TORADOL) IV (Single Response)	Do NOT use in patients with eGFR LESS than 30 mL/min AND/OR patients LESS than 17 years of age. WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery.
() For patients ages GREATER than 64 OR weight LESS than 50 kg OR eGFR 30-59 mL/min - ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6)
() For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6)

<input type="checkbox"/> traMADol (ULTRAM) tablet	50 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury. Not to exceed 400 mg/day. Allowance for Patient Preference:
---	---

Severe Pain (Pain Score 7-10) (Single Response)

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	2 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Monitor and record pain scores and respiratory status. Allowance for Patient Preference:
<input type="checkbox"/> oxyCODONE-acetaminophen (PERCOCET) 5-325 mg per tablet	2 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Monitor and record pain scores and respiratory status. Allowance for Patient Preference:
<input type="checkbox"/> traMADol (ULTRAM) tablet	100 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury. Not to exceed 400 mg/day. Allowance for Patient Preference:
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.8 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op If patient is NPO or cannot tolerate Oral medication, administer the ordered injection.
<input type="checkbox"/> morPHINE injection	4 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op And Notify MD. If patient is NPO or cannot tolerate Oral medication, administer the ordered injection. Allowance for Patient Preference:

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) injection	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.

<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, Administer over: 30 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Itching: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet	25 mg, oral, every 6 hours PRN, itching
<input type="checkbox"/> hydrOXYzine (ATARAX) tablet	10 mg, oral, every 6 hours PRN, itching
<input checked="" type="checkbox"/> cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet - For eGFR LESS than 80 mL/min, reduce frequency to once daily as needed	60 mg, oral, 2 times daily PRN, itching

Bowel Care PRN (Single Response)

<input type="checkbox"/> sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet	2 tablet, oral, nightly PRN, constipation, Post-op
<input type="checkbox"/> magnesium hydroxide suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR WORSE	30 mL, oral, every 12 hours PRN, constipation, Post-op Do not give if patient is on hemodialysis or is in chronic renal failure.
<input type="checkbox"/> bisacodyl (DULCOLAX) EC tablet	10 mg, oral, daily PRN, constipation, Post-op
<input type="checkbox"/> bisacodyl (DULCOLAX) suppository	10 mg, rectal, daily PRN, constipation, Post-op

Bowel Care Scheduled

<input checked="" type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily, Post-op
--	--------------------------------------

Flatulence

<input type="checkbox"/> simethicone (MYLICON) chewable tablet	160 mg, oral, 4 times daily PRN, flatulence, Post-op
--	--

Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
---	--------------------------------

Insomnia: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep
<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

() MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

[] Moderate Risk (Selection Required)

[] Moderate risk of VTE Routine, Once, PACU & Post-op

[] Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

() Contraindications exist for pharmacologic prophylaxis "And" Linked Panel BUT order Sequential compression device

[] Contraindications exist for pharmacologic prophylaxis Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):

[] Place/Maintain sequential compression device continuous Routine, Continuous

() Contraindications exist for pharmacologic prophylaxis "And" Linked Panel AND mechanical prophylaxis

[] Contraindications exist for pharmacologic prophylaxis Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):

[] Contraindications exist for mechanical prophylaxis Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):

() enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight Dose

LESS THAN 100kg enoxaparin 40mg daily

100 to 139kg enoxaparin 30mg every 12 hours

GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours

() For CrCl LESS than 30mL/min - enoxaparin (LOVENOX)

subcutaneous Daily at 1700

[] enoxaparin (LOVENOX) injection 30 mg, subcutaneous, daily at 1700, Starting S+1
Indication(s):

() For CrCl GREATER than or EQUAL TO 30 mL/min -

enoxaparin (LOVENOX) subcutaneous

[] enoxaparin (LOVENOX) injection subcutaneous, Starting S+1
Indication(s):

() fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

() heparin (porcine) injection

5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)

5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM

Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

() heparin (porcine) injection - For Patients with weight GREATER than 100 kg

7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

For patients with weight GREATER than 100 kg.

() warfarin (COUMADIN) tablet

oral, daily at 1700, Starting S+1

Indication:

() Pharmacy consult to manage warfarin (COUMADIN)

STAT, Until discontinued, Starting S

Indication:

[] Mechanical Prophylaxis (Single Response) (Selection Required)

() Contraindications exist for mechanical prophylaxis

Routine, Once

No mechanical VTE prophylaxis due to the following contraindication(s):

() Place/Maintain sequential compression device continuous

Routine, Continuous

() MODERATE Risk of DVT - Non-Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

[] Moderate Risk (Selection Required)

[] Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis - **"And" Linked Panel**
Order Sequential compression device

Contraindications exist for pharmacologic prophylaxis
Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

Place/Maintain sequential compression device continuous
Routine, Continuous

Contraindications exist for pharmacologic prophylaxis **"And" Linked Panel**
AND mechanical prophylaxis

Contraindications exist for pharmacologic prophylaxis
Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

Contraindications exist for mechanical prophylaxis
Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight Dose

LESS THAN 100kg enoxaparin 40mg daily

100 to 139kg enoxaparin 30mg every 12 hours

GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours

For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700

enoxaparin (LOVENOX) injection
30 mg, subcutaneous, daily at 1700
Indication(s):

For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous

enoxaparin (LOVENOX) injection
subcutaneous
Indication(s):

fondaparinux (ARIXTRA) injection
2.5 mg, subcutaneous, daily
If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min
This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection
5,000 Units, subcutaneous, every 8 hours

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)
5,000 Units, subcutaneous, every 12 hours
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

HEParin (porcine) injection - For Patients with weight GREATER than 100 kg
7,500 Units, subcutaneous, every 8 hours
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet
oral, daily at 1700
Indication:

Pharmacy consult to manage warfarin (COUMADIN)
STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis
Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):

Place/Maintain sequential compression device continuous
Routine, Continuous

HIGH Risk of DVT - Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

High risk of VTE Routine, Once, PACU & Post-op

High Risk Pharmacological Prophylaxis - Surgical Patient
(Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)
(Selection Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight Dose

LESS THAN 100kg enoxaparin 40mg daily

100 to 139kg enoxaparin 30mg every 12 hours

GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours

For CrCl LESS than 30mL/min - enoxaparin (LOVENOX)
subcutaneous Daily at 1700

enoxaparin (LOVENOX) injection 30 mg, subcutaneous, daily at 1700, Starting S+1
Indication(s):

For CrCl GREATER than or EQUAL TO 30 mL/min -
enoxaparin (LOVENOX) subcutaneous

enoxaparin (LOVENOX) injection subcutaneous, Starting S+1
Indication(s):

fondaparinux (ARIXTRA) injection 2.5 mg, subcutaneous, daily, Starting S+1
If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

heparin (porcine) injection 5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs) 5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM
Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

heparin (porcine) injection - For Patients with weight GREATER than 100 kg 7,500 Units, subcutaneous, every 8 hours, Starting S+1
For patients with weight GREATER than 100 kg.

warfarin (COUMADIN) tablet oral, daily at 1700, Starting S+1
Indication:

Pharmacy consult to manage warfarin (COUMADIN) STAT, Until discontinued, Starting S
Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition	
Both pharmacologic AND mechanical prophylaxis must be addressed.	
One or more of the following medical conditions:	
Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)	
Severe fracture of hip, pelvis or leg	
Acute spinal cord injury with paresis	
Multiple major traumas	
Abdominal or pelvic surgery for CANCER	
Acute ischemic stroke	
History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
Patient renal status: @CRCL@	
For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:	
Weight Dose	
LESS THAN 100kg enoxaparin 40mg daily	
100 to 139kg enoxaparin 30mg every 12 hours	
GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours	
<input type="checkbox"/> For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700	
<input type="checkbox"/> enoxaparin (LOVENOX) injection	30 mg, subcutaneous, daily at 1700 Indication(s):
<input type="checkbox"/> For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous	
<input type="checkbox"/> enoxaparin (LOVENOX) injection	subcutaneous Indication(s):
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> heparin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
Patient renal status: @CRCL@	
For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight: Weight Dose LESS THAN 100kg enoxaparin 40mg daily 100 to 139kg enoxaparin 30mg every 12 hours GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours	
<input type="checkbox"/> For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700	
<input type="checkbox"/> enoxaparin (LOVENOX) injection	30 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s):
<input type="checkbox"/> For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous	
<input type="checkbox"/> enoxaparin (LOVENOX) injection	subcutaneous, Starting S+1 Indication(s):
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	
2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):	

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> heparin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

<input type="checkbox"/>	Moderate Risk (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/>	Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	"And" Linked Panel
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
	Patient renal status: @CRCL@	
	For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight: Weight Dose LESS THAN 100kg enoxaparin 40mg daily 100 to 139kg enoxaparin 30mg every 12 hours GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours	
<input type="checkbox"/>	For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700	
<input type="checkbox"/>	enoxaparin (LOVENOX) injection	30 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s):
<input type="checkbox"/>	For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous	
<input type="checkbox"/>	enoxaparin (LOVENOX) injection	subcutaneous, Starting S+1 Indication(s):
<input type="checkbox"/>	fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/>	heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/>	heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/>	heparin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
<input type="checkbox"/>	warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/>	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/>	Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous

() MODERATE Risk of DVT - Non-Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

[] Moderate Risk (Selection Required)

[] Moderate risk of VTE Routine, Once, PACU & Post-op

[] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

() Contraindications exist for pharmacologic prophylaxis - **"And" Linked Panel**
Order Sequential compression device

[] Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

[] Place/Maintain sequential compression device continuous Routine, Continuous

() Contraindications exist for pharmacologic prophylaxis **"And" Linked Panel**
AND mechanical prophylaxis

[] Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):

[] Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):

() enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight Dose

LESS THAN 100kg enoxaparin 40mg daily

100 to 139kg enoxaparin 30mg every 12 hours

GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours

() For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700

[] enoxaparin (LOVENOX) injection 30 mg, subcutaneous, daily at 1700
Indication(s):

() For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous

[] enoxaparin (LOVENOX) injection subcutaneous
Indication(s):

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> HEParin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
Patient renal status: @CRCL@	
For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight: Weight Dose LESS THAN 100kg enoxaparin 40mg daily 100 to 139kg enoxaparin 30mg every 12 hours GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours	
<input type="checkbox"/> For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700	
<input type="checkbox"/> enoxaparin (LOVENOX) injection	30 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s):
<input type="checkbox"/> For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous	

<input type="checkbox"/> enoxaparin (LOVENOX) injection	subcutaneous, Starting S+1 Indication(s):
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> heparin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, Starting S+1 For patients with weight GREATER than 100 kg.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
Patient renal status: @CRCL@	
For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight: Weight Dose LESS THAN 100kg enoxaparin 40mg daily 100 to 139kg enoxaparin 30mg every 12 hours GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours	
<input type="checkbox"/> For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700	

<input type="checkbox"/> enoxaparin (LOVENOX) injection	30 mg, subcutaneous, daily at 1700 Indication(s):
() For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous	
<input type="checkbox"/> enoxaparin (LOVENOX) injection	subcutaneous Indication(s):
() fondaparinux (ARIXTRA) injection	
	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	
	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	
	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() heparin (porcine) injection - For Patients with weight GREATER than 100 kg	
	7,500 Units, subcutaneous, every 8 hours For patients with weight GREATER than 100 kg.
() warfarin (COUMADIN) tablet	
	oral, daily at 1700, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	
	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
() Contraindications exist for mechanical prophylaxis	
	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	
() HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
() Contraindications exist for pharmacologic prophylaxis	
	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
() aspirin chewable tablet	
	162 mg, oral, daily, Starting S+1
() aspirin (ECOTRIN) enteric coated tablet	
	162 mg, oral, daily, Starting S+1
() Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
() enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight Dose

LESS THAN 100kg enoxaparin 40mg daily

100 to 139kg enoxaparin 30mg every 12 hours

GREATER THAN or EQUAL to 140kg enoxaparin 40mg every 12 hours

() For CrCl LESS than 30mL/min - enoxaparin (LOVENOX) subcutaneous Daily at 1700	
[] enoxaparin (LOVENOX) injection	30 mg, subcutaneous, daily at 1700, Starting S+1 Indication(s):
() For CrCl GREATER than or EQUAL TO 30 mL/min - enoxaparin (LOVENOX) subcutaneous	
[] enoxaparin (LOVENOX) injection	subcutaneous, Starting S+1 Indication(s):
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() heparin (porcine) injection - For Patients with weight GREATER than 100 kg	7,500 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM For patients with weight GREATER than 100 kg.
() Rivaroxaban and Pharmacy Consult (Selection Required)	
[] rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
[] Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
() warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

Rh Negative Mother

Nursing

[X] Rhogam Workup: If Mother is Rh Negative, complete Rhogam workup and administer Rh immune globulin 50 mcg (or dose determined by lab antibody results) IM within 72 hours of delivery.	Routine, Until discontinued, Starting S, Post-op
---	--

Labs

[] Fetal Screen	Conditional Frequency For 1 Occurrences Conditional- One activation- If Rh Negative Mom and Rh Positive infant, Post-op
------------------	--

<input type="checkbox"/> Rhogam Type and Screen	Once, Post-op
---	---------------

Medication

<input type="checkbox"/> rho(D) immune globulin (HYPERRHO/RHOAM) injection	300 mcg, intramuscular, PRN, Rhogam Workup: If Mother is Rh Negative, complete Rhogam workup and administer Rh immune globulin 50 mcg (or dose determined by lab antibody results) IM within 72 hours of delivery., For 1 Doses, Post-op
--	--

Labs Tomorrow

Hematology

<input type="checkbox"/> CBC with differential	AM draw, Starting S+1 For 1 Occurrences, Post-op
<input type="checkbox"/> Hemoglobin and hematocrit	Once, Post-op

Chemistry

<input type="checkbox"/> Basic metabolic panel	Once, Post-op
--	---------------

Consults

For Physician Consult orders use sidebar

Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason: Post-op
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Consult PT Eval and Treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: Post-op
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic: Post-op
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult? Post-op
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult? Post-op