

General

Phase of Care

This order must be checked to allow group discontinuation of pre-delivery orders by nursing during post partum phase of care.

Discontinue pre-delivery orders Routine, Once, L&D Pre-Delivery

Admission Orders (Single Response) (Selection Required)

Do not use Outpatient Observation Services Under General Supervision order for patients who are receiving Outpatient TRIAGE services.

Admit to L&D
 Admitting Physician:
 Diagnosis: Pregnancy
 Bed request comments:
 L&D Pre-Delivery

Code Status

Full code Code Status decision reached by: Patient by means of Oral Directive
 L&D Pre-Delivery

DNR (Do Not Resuscitate) (Selection Required)

DNR (Do Not Resuscitate)
 Did the patient/surrogate require the use of an interpreter?
 Did the patient/surrogate require the use of an interpreter?
 Does patient have decision-making capacity?
 L&D Pre-Delivery

Consult to Palliative Care Service

Consult to Palliative Care Service
 Priority:
 Reason for Consult?
 Order?
 Name of referring provider:
 Enter call back number:

Consult to Social Work
 Reason for Consult:
 L&D Pre-Delivery

Modified Code
 Did the patient/surrogate require the use of an interpreter?
 Did the patient/surrogate require the use of an interpreter?
 Does patient have decision-making capacity?
 Modified Code restrictions:
 L&D Pre-Delivery

Treatment Restrictions ((For use when a patient is NOT in a cardiopulmonary arrest))
 I understand that if the patient is NOT in a cardiopulmonary arrest, the selected treatments will NOT be provided. I understand that all other unselected medically indicated treatments will be provided.
 Treatment Restriction decision reached by:
 Specify Treatment Restrictions:
 L&D Pre-Delivery

Isolation

Airborne isolation status

Airborne isolation status Details
 Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics. Once, L&D Pre-Delivery

Contact isolation status Details

Droplet isolation status Details

Enteric isolation status Details

Precautions

<input type="checkbox"/>	Aspiration precautions	L&D Pre-Delivery
<input type="checkbox"/>	Fall precautions	Increased observation level needed: On Admission and every 8 hours, L&D Pre-Delivery
<input type="checkbox"/>	Latex precautions	L&D Pre-Delivery
<input type="checkbox"/>	Seizure precautions	Increased observation level needed: L&D Pre-Delivery

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	L&D Pre-Delivery
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	L&D Pre-Delivery
<input type="checkbox"/>	Acute Renal Failure	L&D Pre-Delivery
<input type="checkbox"/>	Acute Respiratory Failure	L&D Pre-Delivery
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	L&D Pre-Delivery
<input type="checkbox"/>	Anemia	L&D Pre-Delivery
<input type="checkbox"/>	Bacteremia	L&D Pre-Delivery
<input type="checkbox"/>	Bipolar disorder, unspecified	L&D Pre-Delivery
<input type="checkbox"/>	Cardiac Arrest	L&D Pre-Delivery
<input type="checkbox"/>	Cardiac Dysrhythmia	L&D Pre-Delivery
<input type="checkbox"/>	Cardiogenic Shock	L&D Pre-Delivery
<input type="checkbox"/>	Decubitus Ulcer	L&D Pre-Delivery
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	L&D Pre-Delivery
<input type="checkbox"/>	Disorder of Liver	L&D Pre-Delivery
<input type="checkbox"/>	Electrolyte and Fluid Disorder	L&D Pre-Delivery
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	L&D Pre-Delivery
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	L&D Pre-Delivery
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	L&D Pre-Delivery
<input type="checkbox"/>	Other Alteration of Consciousness	L&D Pre-Delivery
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	L&D Pre-Delivery
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	L&D Pre-Delivery
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	L&D Pre-Delivery
<input type="checkbox"/>	Protein-calorie Malnutrition	L&D Pre-Delivery
<input type="checkbox"/>	Psychosis, unspecified psychosis type	L&D Pre-Delivery
<input type="checkbox"/>	Schizophrenia Disorder	L&D Pre-Delivery
<input type="checkbox"/>	Sepsis	L&D Pre-Delivery
<input type="checkbox"/>	Septic Shock	L&D Pre-Delivery
<input type="checkbox"/>	Septicemia	L&D Pre-Delivery
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	L&D Pre-Delivery
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	L&D Pre-Delivery
<input type="checkbox"/>	Present on Admission-History of preterm premature rupture of membranes	L&D Pre-Delivery

Nursing

Vital Signs

<input checked="" type="checkbox"/>	Vital signs - T/P/R/BP (per unit protocol)	Routine, Per unit protocol, L&D Pre-Delivery
-------------------------------------	--	--

Activity

<input type="checkbox"/>	Strict bed rest	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input checked="" type="checkbox"/>	Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges L&D Pre-Delivery
<input type="checkbox"/>	May use birthing ball	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/>	Bed rest with bathroom privileges for BM	Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges For BM, L&D Pre-Delivery
<input type="checkbox"/>	Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated L&D Pre-Delivery

May use Whirlpool tub if membranes intact Routine, Until discontinued, Starting S, L&D Pre-Delivery

Nursing Care

Apply external fetal monitor

Monitor fetal heart tones continuous Routine, Continuous
Type: Continuous
L&D Pre-Delivery

Monitor fetal heart tones intermittent Routine, Continuous
Type: Intermittent
L&D Pre-Delivery

Fetal nonstress test Routine, Once, L&D Pre-Delivery

Tocometry Routine, Until discontinued, Starting S
Type: Continuous
L&D Pre-Delivery

Apply internal fetal monitor (FSE)

Fetal scalp monitor Routine, Continuous
Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured.
May place if GBS positive: Yes
L&D Pre-Delivery

POC Amnisure Once, L&D Pre-Delivery

Sterile vaginal exam Routine, Until discontinued, Starting S
Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery

For urinary retention: Assist patient to void on bedpan in upright position prior to straight cath x 1 followed by placement of indwelling urinary catheter if necessary Routine, Until discontinued, Starting S, L&D Pre-Delivery

Straight cath Routine, Conditional Frequency For 1 Occurrences
If patient unable to void on bedpan, preferably in upright position, straight cath x 1. If patient unable to void after 1st straight cath, insert indwelling urinary catheter., L&D Pre-Delivery

Insert and maintain Foley

Insert Foley catheter Routine, Conditional Frequency For 1 Occurrences
Type:
Size:
Urinometer needed:
If regional block and patient unable to void insert indwelling urinary catheter., L&D Pre-Delivery

Foley Catheter Care Routine, Conditional Frequency For 1 Occurrences
Orders: Maintain
L&D Pre-Delivery

Patient may have epidural Routine, Once
Notify Anesthesia immediately upon patient's request and begin pre-epidural hydration, L&D Pre-Delivery

Place/Maintain sequential compression device continuous Routine, Continuous, L&D Pre-Delivery

Place antiembolic stockings Routine, Until discontinued, Starting S, L&D Pre-Delivery

Nursing Care

Apply external fetal monitor

Monitor fetal heart tones continuous Routine, Continuous
Type: Continuous
L&D Pre-Delivery

Monitor fetal heart tones intermittent Routine, Continuous
Type: Intermittent
L&D Pre-Delivery

Fetal nonstress test Routine, Once, L&D Pre-Delivery

Tocometry Routine, Until discontinued, Starting S
Type: Continuous
L&D Pre-Delivery

<input type="checkbox"/>	Apply internal fetal monitor (FSE)	
<input type="checkbox"/>	Fetal scalp monitor	Routine, Continuous Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured. May place if GBS positive: Yes L&D Pre-Delivery
<input type="checkbox"/>	Amnisure	Once, Amniotic fluid, L&D Pre-Delivery
<input checked="" type="checkbox"/>	Sterile vaginal exam	Routine, Until discontinued, Starting S Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery
<input type="checkbox"/>	For urinary retention: Assist patient to void on bedpan in upright position prior to straight cath x 1 followed by placement of indwelling urinary catheter if necessary	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/>	Straight cath	Routine, Conditional Frequency For 1 Occurrences If patient unable to void on bedpan, preferably in upright position, straight cath x 1. If patient unable to void after 1st straight cath, insert indwelling urinary catheter., L&D Pre-Delivery
<input checked="" type="checkbox"/>	Insert and maintain Foley	
<input checked="" type="checkbox"/>	Insert Foley catheter	Routine, Conditional Frequency For 1 Occurrences Type: Size: Urnometer needed: If regional block and patient unable to void insert indwelling urinary catheter., L&D Pre-Delivery
<input checked="" type="checkbox"/>	Foley Catheter Care	Routine, Conditional Frequency For 1 Occurrences Orders: Maintain L&D Pre-Delivery
<input type="checkbox"/>	Patient may have epidural	Routine, Once Notify Anesthesia immediately upon patient's request and begin pre-epidural hydration, L&D Pre-Delivery
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/>	Place antiembolic stockings	Routine, Until discontinued, Starting S, L&D Pre-Delivery

Nursing Care

<input checked="" type="checkbox"/>	Apply external fetal monitor	
<input checked="" type="checkbox"/>	Monitor fetal heart tones continuous	Routine, Continuous Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Monitor fetal heart tones intermittent	Routine, Continuous Type: Intermittent L&D Pre-Delivery
<input type="checkbox"/>	Fetal nonstress test	Routine, Once, L&D Pre-Delivery
<input checked="" type="checkbox"/>	Tocometry	Routine, Until discontinued, Starting S Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Apply internal fetal monitor (FSE)	
<input type="checkbox"/>	Fetal scalp monitor	Routine, Continuous Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured. May place if GBS positive: Yes L&D Pre-Delivery
<input type="checkbox"/>	POC Amnisure	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/>	Sterile vaginal exam	Routine, Until discontinued, Starting S Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery
<input type="checkbox"/>	For urinary retention: Assist patient to void on bedpan in upright position prior to straight cath x 1 followed by placement of indwelling urinary catheter if necessary	Routine, Until discontinued, Starting S, L&D Pre-Delivery

<input type="checkbox"/> Straight cath	Routine, Conditional Frequency For 1 Occurrences If patient unable to void on bedpan, preferably in upright position, straight cath x 1. If patient unable to void after 1st straight cath, insert indwelling urinary catheter., L&D Pre-Delivery
<input checked="" type="checkbox"/> Insert and maintain Foley	
<input checked="" type="checkbox"/> Insert Foley catheter	Routine, Conditional Frequency For 1 Occurrences Type: Size: Urinometer needed: If regional block and patient unable to void insert indwelling urinary catheter., L&D Pre-Delivery
<input checked="" type="checkbox"/> Foley Catheter Care	Routine, Conditional Frequency For 1 Occurrences Orders: Maintain L&D Pre-Delivery
<input type="checkbox"/> Patient may have epidural	Routine, Once Notify Anesthesia immediately upon patient's request and begin pre-epidural hydration, L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/> Place antiembolic stockings	Routine, Until discontinued, Starting S, L&D Pre-Delivery
Notify	
<input checked="" type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 100.3 Temperature less than: Systolic BP greater than: 160 Systolic BP less than: 80 Diastolic BP greater than: 110 Diastolic BP less than: 50 MAP less than: Heart rate greater than (BPM): Heart rate less than (BPM): Respiratory rate greater than: 24 Respiratory rate less than: 12 SpO2 less than:
<input checked="" type="checkbox"/> Notify Anesthesiologist immediately if patient requests an epidural and begin pre-epidural hydration	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input checked="" type="checkbox"/> Notify Nursery and Neonatologist to attend delivery if indicated	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> Notify (General)	Routine, Until discontinued, Starting S, L&D Pre-Delivery
Bowel Care	
<input type="checkbox"/> Tap water enema	Routine, Once On Admission, L&D Pre-Delivery
<input type="checkbox"/> mineral oil enema	1 enema, rectal, once PRN, constipation, L&D Pre-Delivery Once on admission
Consent	
<input checked="" type="checkbox"/> Complete Consent Form	Routine, Once Consent For: Delivery Procedure: Vaginal delivery of fetus and placenta with possible cesarean section, possible episiotomy, and possible use of vacuum/forceps. Diagnosis/Condition: Physician: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? L&D Pre-Delivery

Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: An NPO order without explicit exceptions means nothing can be given orally to the patient., L&D Pre-Delivery
<input checked="" type="checkbox"/> NPO with ice chips	Diet effective now, Starting S NPO: Except Ice chips Pre-Operative fasting options: 1/2 cup per hour, An NPO order without explicit exceptions means nothing can be given orally to the patient., L&D Pre-Delivery
<input type="checkbox"/> Diet - Clear liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: L&D Pre-Delivery

IV Fluids

IV Fluids

<input checked="" type="checkbox"/> lactated Ringer's bolus	1,000 mL, intravenous, Administer over: 30 Minutes, once PRN, If patient requests epidural - for epidural prehydration, L&D Pre-Delivery Notify Anesthesiologist immediately if patient requests Epidural and begin pre-epidural hydration.
<input type="checkbox"/> dextrose 5 % and lactated Ringer's infusion	125 mL/hr, intravenous, continuous, L&D Pre-Delivery
<input checked="" type="checkbox"/> lactated ringer's infusion	125 mL/hr, intravenous, continuous, L&D Pre-Delivery

Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled, L&D Pre-Delivery if IV is saline locked
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care, L&D Pre-Delivery

Medications

Nitrous Oxide Orders

<input type="checkbox"/> Nitrous Oxide Administration Orders	
<input type="checkbox"/> Activity	
<input type="checkbox"/> Fall precautions - while using nitrous oxide	Increased observation level needed: L&D Pre-Delivery
<input type="checkbox"/> Consent	
<input type="checkbox"/> Complete Consent Form	Routine, Once Consent For: Use of Nitrous Oxide Procedure: Use of Nitrous Oxide for pain control Diagnosis/Condition: Physician: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? L&D Pre-Delivery
<input type="checkbox"/> Vital Signs	
<input type="checkbox"/> Continuous Fetal monitoring	Routine, Once, L&D Pre-Delivery

<input type="checkbox"/> Vital Signs - Nitrous Oxide use	Routine, Per unit protocol For Until specified Nitrous Oxide Use - Baseline Vital signs within One Hour prior to administration, Vital signs 15 minutes after start time, Vital signs hourly throughout duration of use, Pain scale documented every 15 minutes during Nitrous Oxide use, L&D Pre-Delivery
<input type="checkbox"/> Continuous Oxygen saturation monitoring	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Patient Education	
<input type="checkbox"/> RN to provide patient education prior to the initiation of Nitrous Oxide.	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> Nitrous Oxide	
<input type="checkbox"/> nitrous oxide gas	inhalation, continuous PRN, labor pain analgesia, L&D Pre-Delivery

Antibiotics (Single Response)

Does your patient have a penicillin allergy?

<input type="checkbox"/> No	
<input type="checkbox"/> Antibiotics: if GBS+ or Unknown (Single Response)	
<input type="checkbox"/> penicillin G IVPB Loading and Maintenance Dose - Prophylaxis Regimen for GBS	"Followed by" Linked Panel
<input type="checkbox"/> Loading Dose - penicillin G (POTASSIUM) IV	5 Million Units, intravenous, Administer over: 30 Minutes, once, For 1 Doses, L&D Pre-Delivery If GBS positive Indication: Medical Prophylaxis
<input type="checkbox"/> Maintenance Dose - penicillin G (POTASSIUM) IV	2.5 Million Units, intravenous, Administer over: 30 Minutes, every 4 hours, Starting H+4 Hours, L&D Pre-Delivery If GBS positive Indication: Medical Prophylaxis
<input type="checkbox"/> ampicillin IVPB Loading and Maintenance Dose - Alternative Regimen for GBS	
"Followed by" Linked Panel	
<input type="checkbox"/> Loading Dose - ampicillin IV	2 g, intravenous, Administer over: 30 Minutes, once, For 1 Doses, L&D Pre-Delivery If GBS positive Reason for Therapy: Medical Prophylaxis
<input type="checkbox"/> Maintenance Dose - ampicillin IV	1 g, intravenous, Administer over: 30 Minutes, every 4 hours, Starting H+4 Hours, L&D Pre-Delivery If GBS positive Reason for Therapy: Medical Prophylaxis
<input type="checkbox"/> Prophylaxis Regimen for C-section	
<input type="checkbox"/> Patients LESS than or equal to 120 kg	
<input type="checkbox"/> ceFAZolin (ANCEF) IV - Give within 60 minutes prior to C-Section	2 g, intravenous, once PRN, Prior to C-Section, Pre-op Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> azithromycin (ZITHROMAX) IV - Give within 60 minutes prior to C-Section	500 mg, intravenous, Administer over: 60 Minutes, once PRN, Prior to C-Section, Pre-op Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> Patients GREATER than 120 kg	
<input type="checkbox"/> ceFAZolin (ANCEF) IV - Give within 60 minutes prior to C-Section	3 g, intravenous, once PRN, Prior to C-Section, Pre-op Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:

[] azithromycin (ZITHROMAX) IV - Give within 60 minutes prior to C-Section	500 mg, intravenous, Administer over: 60 Minutes, once PRN, Prior to C-Section, Pre-op Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
() Yes	
[] Penicillin Allergic and GBS + (Single Response)	
() ceFAZolin (ANCEF) IV Loading and Maintenance Doses - if GBS Positive	"Followed by" Linked Panel
Recommended for patients NOT high risk for anaphylaxis	
[] Loading Dose - cefazolin (ANCEF) IV	2 g, intravenous, once, For 1 Doses, L&D Pre-Delivery If GBS positive Reason for Therapy: Medical Prophylaxis Medical Prophylaxis: The maximum duration of therapy that can be entered at this time is 56 days. Order can be renewed if required at that time.
[] Maintenance Dose - cefazolin (ANCEF) IV	1 g, intravenous, every 8 hours, L&D Pre-Delivery If GBS positive. Through delivery then discontinue. Reason for Therapy: Medical Prophylaxis Medical Prophylaxis: The maximum duration of therapy that can be entered at this time is 56 days. Order can be renewed if required at that time.
() clindamycin (CLEOCIN) IV Loading and Maintenance Doses - if GBS Positive	"Followed by" Linked Panel
Recommended ONLY for patients with high risk for penicillin anaphylaxis that are culture isolate sensitive to Clindamycin.	
[] Loading Dose - clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, L&D Pre-Delivery If GBS positive Reason for Therapy: Medical Prophylaxis
[] Maintenance Dose - clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, every 8 hours, L&D Pre-Delivery If GBS positive. Through delivery then discontinue. Reason for Therapy: Medical Prophylaxis
() vancomycin (VANCOCIN) IVPB - if GBS Positive (Selection Required)	
Adjust doses for renal function if CrCl LESS THAN 60 mL/min	
[] vancomycin (VANCOCIN) IV	20 mg/kg, intravenous, every 8 hours, L&D Pre-Delivery If GBS positive. Ensure through delivery. Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy: Medical Prophylaxis Medical Prophylaxis: The maximum duration of therapy that can be entered at this time is 56 days. Order can be renewed if required at that time.
[] Prophylaxis Regimen for C-section	"And" Linked Panel
[] azithromycin (ZITHROMAX)	500 mg, intravenous, Administer over: 60 Minutes, once, For 1 Doses, Pre-op Nurse to send medication(s) to operating room - To be administered by Anesthesiologist. To be given 1 hour prior to skin incision. Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:

<input type="checkbox"/> clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Pre-op Nurse to send medication(s) to operating room - To be administered by Anesthesiologist. To be given 1 hour prior to skin incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> gentamicin (GARAMICIN) IVPB	5 mg/kg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Pre-op Nurse to send medication(s) to operating room - To be administered by Anesthesiologist. To be given 1 hour prior to skin incision. Reason for Therapy: Surgical Prophylaxis

PRN Antihypertensives

<input type="checkbox"/> For blood pressure GREATER than or EQUAL to 160 mmHg - labetalol (NORMODYNE) tablet	200 mg, oral, 2 times daily PRN, high blood pressure, For Systolic BP GREATER than 160mmHg and Diastolic BP GREATER than 110mmHg., L&D Pre-Delivery DO NOT give incremental dosing as this will compromise fetal circulation. BP & HR HOLD parameters for this order: Hold Parameters requested BP & HR HOLD for: HOLD for: Contact Physician if: For Systolic BP GREATER than 160mmHG and Diastolic BP GREATER than 110mmHg
<input type="checkbox"/> hydrALAZINE (APRESOLINE) tablet	5 mg, oral, every 6 hours PRN, high blood pressure, For blood pressure GREATER than 160mmHG, L&D Pre-Delivery DO NOT give incremental dosing as this will compromise fetal circulation. BP HOLD parameters for this order: Contact Physician if:

PRN Severe Hypertension

<input type="checkbox"/> NIFEdipine (PROCARDIA) capsule	10 mg, oral, once PRN, high blood pressure, for severe BP elevations of 15 min or more. Recheck BP in 15 min., For 1 Doses, L&D Pre-Delivery BP & HR HOLD parameters for this order: Contact Physician if: For Systolic BP GREATER than 160mmHG and Diastolic BP GREATER than 110mmHg
<input type="checkbox"/> labetalol (NORMODYNE, TRANDATE) injection	20 mg, intravenous, once PRN, high blood pressure, for severe blood pressure elevation (systolic BP GREATER than or EQUAL to 110 mm Hg) persisting for 15 minutes or more., For 1 Doses, L&D Pre-Delivery Give IV Push over 2 minutes. Repeat BP measurements in 10 minutes and record results. BP & HR HOLD parameters for this order: Contact Physician if: For Systolic BP GREATER than 160mmHG and Diastolic BP GREATER than 110mmHg

Non-Reassuring FHR with Tachysystole

<input checked="" type="checkbox"/> terbutaline (BRETHINE) injection	0.25 mg, subcutaneous, once PRN, tachysystole with non-reassuring fetal heart rate, L&D Pre-Delivery
--	--

Cervical Ripening (Single Response)

<input type="checkbox"/> Vaginal - misoprostol (CYTOTEC) tablet for vaginal use	25 mcg, vaginal, every 4 hours, L&D Pre-Delivery
<input type="checkbox"/> dinoprostone (CERVIDIL) vaginal insert	10 mg, vaginal, Administer over: 12 Hours, once, For 1 Doses, L&D Pre-Delivery Remove Dinoprostone (Cervidil) 12 hours after placement (or if non-reassuring FHR tracing, tachysystole, onset of active labor). After insertion of Dinoprostone (Cervidil), position patient supine with lateral tilt for 2 hours

Induction/Augmentation (Single Response)

() misoprostol (CYTOTEC) tablet for vaginal use	25 mcg, vaginal, every 4 hours, L&D Pre-Delivery
() Oxytocin Induction (Single Response)	
() Low Dose	<p>2-30 milli-units/min, intravenous, titrated, L&D Pre-Delivery Begin 30 minutes after the removal of dinoprostone (CERVIDIL) insert OR begin 4 hours after the last dose of misoprostol (CYTOTEC).</p> <p>In the absence of FHR abnormalities, start at 2 mu/min and increase by 2 mu/min every 20 minutes to a maximum of 30 milliunits/minute or until adequate uterine activity is achieved. Call the physician if 30 milliunits is reached without adequate contractions.</p> <p>Adequate uterine activity is defined as: uterine contractions that are 2-3 minutes apart, contraction duration of 40-90 seconds and moderate intensity by palpation or 50-60 mmHg above baseline with IUPC. Contractions are not to exceed 5 contractions in 10 minutes averaged over a 30-minute window, last 2 minutes or more, occur within 1 minute of each other, or result in insufficient resting tone between contractions.</p>
() Medium Dose	<p>4-30 milli-units/min, intravenous, titrated, L&D Pre-Delivery Begin 30 minutes after the removal of dinoprostone (CERVIDIL) insert OR begin 4 hours after the last dose of misoprostol (CYTOTEC).</p> <p>In the absence of FHR abnormalities, start at 4 mu/min and increase by 4 mu/min every 30 minutes to a maximum of 30 milliunits/ minute or until adequate uterine activity is achieved. Call the physician if 30 milliunits is reached without adequate contractions.</p> <p>Adequate uterine activity is defined as: uterine contractions that are 2-3 minutes apart, contraction duration of 40-90 seconds and moderate intensity by palpation or 50-60 mmHg above baseline with IUPC. Contractions are not to exceed 5 contractions in 10 minutes averaged over a 30-minute window, last 2 minutes or more, occur within 1 minute of each other, or result in insufficient resting tone between contractions.</p>
() High Dose	<p>6-30 milli-units/min, intravenous, titrated, L&D Pre-Delivery Begin 30 minutes after the removal of dinoprostone (CERVIDIL) insert OR begin 4 hours after the last dose of misoprostol (CYTOTEC).</p> <p>In the absence of FHR abnormalities, start at 6 mu/min and increase by 6 mu/min every 30 minutes to a maximum of 30 milliunits/minute or until adequate uterine activity is achieved. Call the physician if 30 milliunits is reached without adequate contractions.</p> <p>Adequate uterine activity is defined as: uterine contractions that are 2-3 minutes apart, contraction duration of 40-90 seconds and moderate intensity by palpation or 50-60 mmHg above baseline with IUPC. Contractions are not to exceed 5 contractions in 10 minutes averaged over a 30-minute window, last 2 minutes or more, occur within 1 minute of each other, or result in insufficient resting tone between contractions.</p>

PostPartum Oxytocin

[X] oxytocin (PITOCIN) Bolus and Maintenance Infusion	"Followed by" Linked Panel
[X] oxytocin 30 unit/500 mL bolus from bag	10 Units, intravenous, Administer over: 30 Minutes, once, For 1 Doses, L&D Pre-Delivery
[X] oxytocin (PITOCIN) infusion	<p>5.7 Units/hr, intravenous, continuous, Starting H+30 Minutes, L&D Pre-Delivery Run at 95 mL/hr for 3.5 hours. Total 20 units over 3.5 hours. (Infuse at rate of 95 mL/hr over 3.5 hours) Increase to 999 mL/hr for uterine tone and bleeding per physician instructions.</p>

Bleeding Medications PostPartum (Single Response)

() oxytocin (PITOCIN) infusion and methylergonovine (METHERGINE)	"And" Linked Panel
--	---------------------------

methylergonovine (METHERGINE) is contraindicated if BP GREATER than 140/90 mmHg

- | | |
|--|---|
| <input type="checkbox"/> oxytocin (PITOCIN) infusion | 5.7 Units/hr, intravenous, continuous PRN, PostPartum Vaginal Bleeding, Postpartum
If uterine atony or if excessive bleeding persists, infuse oxytocin at 999mL/hr. |
| <input type="checkbox"/> methylergonovine (METHERGINE) injection
- Contraindicated if BP GREATER than 140/90 mmHg | 200 mcg, intramuscular, once PRN, as needed for vaginal bleeding not controlled by oxytocin, Postpartum
Use if inadequate response to oxytocin. Notify Physician if further treatment needed. Contraindicated if BP GREATER than 140/90 mmHg |

() oxytocin (PITOCIN) infusion AND carboprost (HEMABATE) injection And diphenoxylate-atropine (LOMOTIL) oral dose **"And" Linked Panel**

- | | |
|--|--|
| <input type="checkbox"/> oxytocin (PITOCIN) infusion | 5.7 Units/hr, intravenous, continuous PRN, PostPartum Vaginal Bleeding, Postpartum
If uterine atony or if excessive bleeding persists, infuse oxytocin at 999mL/hr. |
| <input type="checkbox"/> carboprost (HEMABATE) injection | 250 mcg, intramuscular, once PRN, for Vaginal Bleeding uncontrolled by oxytocin., Postpartum |
| <input type="checkbox"/> diphenoxylate-atropine (LOMOTIL)
2.5-0.025 mg per tablet | 1 tablet, oral, once PRN, diarrhea, Postpartum |

() oxytocin (PITOCIN) infusion and misoprostol (CYTOTEC) **"And" Linked Panel**

- | | |
|---|---|
| <input type="checkbox"/> oxytocin (PITOCIN) infusion | 5.7 Units/hr, intravenous, continuous PRN, Postpartum Bleeding, Postpartum
If uterine atony or if excessive bleeding persists, infuse oxytocin at 999mL/hr |
| <input type="checkbox"/> misoprostol (CYTOTEC) tablet | 1,000 mcg, rectal, once PRN, as needed for vaginal bleeding not controlled by oxytocin, Postpartum
Use if inadequate response to oxytocin. Notify Physician if further treatment needed. |

Fetal Demise

- | | |
|---|------------------------|
| <input type="checkbox"/> misoprostol (CYTOTEC) tablet | oral, L&D Pre-Delivery |
|---|------------------------|

NALOXONE FOR LABOR ADMISSION OPIOID PAIN MEDICATIONS

- | | |
|---|---|
| <input checked="" type="checkbox"/> naloxone (NARCAN) 0.4 mg/mL injection | intravenous, PRN, respiratory depression, opioid reversal, L&D Pre-Delivery |
|---|---|

Moderate Pain (Pain Score 4-6) (Single Response)

- | | |
|------------------------------------|---|
| () fentaNYL (SUBLIMAZE) injection | 50 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), L&D Pre-Delivery
Allowance for Patient Preference: |
| () morPHINE injection | 2 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery
Allowance for Patient Preference: |
| () nalbuphine (NUBAIN) injection | 5 mg, intravenous, every 4 hours PRN, moderate pain (score 4-6), L&D Pre-Delivery
Allowance for Patient Preference: |

Moderate Pain (Pain Score 4-6) (Single Response)

- | | |
|------------------------------------|---|
| () fentaNYL (SUBLIMAZE) injection | 50 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), L&D Pre-Delivery
Allowance for Patient Preference: |
| () morPHINE injection | 2 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery
Allowance for Patient Preference: |
| () nalbuphine (NUBAIN) injection | 5 mg, intravenous, every 4 hours PRN, moderate pain (score 4-6), L&D Pre-Delivery
Allowance for Patient Preference: |

Moderate Pain (Pain Score 4-6) (Single Response)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	50 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), L&D Pre-Delivery Allowance for Patient Preference:
<input type="checkbox"/> morPHINE injection	2 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery Allowance for Patient Preference:
<input type="checkbox"/> nalbuphine (NUBAIN) injection	5 mg, intravenous, every 4 hours PRN, moderate pain (score 4-6), L&D Pre-Delivery Allowance for Patient Preference:

Severe Pain (Pain Score 7-10) (Single Response)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	100 mcg, intravenous, every 2 hour PRN, severe pain (score 7-10), L&D Pre-Delivery Allowance for Patient Preference:
<input type="checkbox"/> morPHINE injection	4 mg, intravenous, once PRN, severe pain (score 7-10), L&D Pre-Delivery Allowance for Patient Preference:
<input type="checkbox"/> nalbuphine (NUBAIN) injection	10 mg, intravenous, every 4 hours PRN, severe pain (score 7-10), L&D Pre-Delivery Allowance for Patient Preference:

Severe Pain (Pain Score 7-10) (Single Response)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	100 mcg, intravenous, every 2 hour PRN, severe pain (score 7-10), L&D Pre-Delivery Allowance for Patient Preference:
<input type="checkbox"/> morPHINE injection	4 mg, intravenous, once PRN, severe pain (score 7-10), L&D Pre-Delivery Allowance for Patient Preference:
<input type="checkbox"/> nalbuphine (NUBAIN) injection	10 mg, intravenous, every 4 hours PRN, severe pain (score 7-10), L&D Pre-Delivery Allowance for Patient Preference:

Local Anesthetics - NOT HMTW

<input checked="" type="checkbox"/> lidocaine (XYLOCAINE) 10 mg/mL (1 %) injection	intradermal, PRN, As needed for perineal repair, L&D Pre-Delivery
--	---

Local Anesthetics - HMTW Only

<input checked="" type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1 %) injection	intradermal, PRN, As needed for perineal repair, L&D Pre-Delivery
---	---

Antiemetics - HMH, HMSJ, HMW, HMSTC, HMTW Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSL, HMWB Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input checked="" type="checkbox"/> promethazine (PHENERGAN) injection	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSTJ Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, Administer over: 30 Minutes, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

VTE/DVT Risk Definitions

URL:

"\\epic-nas.et0922.epichosted.com\static\OrderSets\VTE Risk Assessment Tool v7_MAK FINAL.pdf"

<input type="checkbox"/> VERY LOW Risk of DVT (Selection Required)	"And" Linked Panel
<input type="checkbox"/> Very low risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Ambulate	Routine, 3 times daily Specify: Early ambulation, L&D Pre-Delivery
<input type="checkbox"/> Avoid dehydration	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	

Anticipated admission LESS than or EQUAL to 72 hours.

Does not meet moderate or high risk criteria:

Moderate Risk High Risk

Anticipated or actual LOS admission GREATER than or EQUAL to 72 hours High risk thrombophilia with no prior VTE

Prior idiopathic, or estrogen related VTE

Low risk thrombophilia AND family history of VTE OR single prior VTE

Receiving outpatient prophylactic LMWH or UFH

<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, SCDs are recommended while in bed and until fully ambulatory L&D Pre-Delivery
--	--

<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous While in bed AND until fully ambulatory. Encourage early ambulation. Avoid dehydration., L&D Pre-Delivery

MODERATE Risk of DVT (Selection Required)

Anticipated or actual LOS admission GREATER than 72 hours; does not meet High risk criteria. CONSIDER prophylactic LMWH/UFH (consult Anesthesia for delivery considerations)

High Risk

High risk thrombophilia with no prior VTE

Prior idiopathic, or estrogen related VTE

Low risk thrombophilia AND family history of VTE OR single prior VTE

Receiving outpatient prophylactic LMWH or UFH

<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous SCD throughout hospitalization. Encourage early ambulation. Avoid dehydration., L&D Pre-Delivery

HIGH Risk of DVT (Selection Required)

High risk thrombophilia with no prior VTE

Prior idiopathic or estrogen related VTE

Low risk thrombophilia AND (family history of VTE OR single prior VTE)

Receiving outpatient prophylactic LMWH or UFH

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous While in bed AND until fully ambulatory. Encourage early ambulation. Avoid dehydration., L&D Pre-Delivery

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)

<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery

Place/Maintain sequential compression device continuous Routine, Continuous, L&D Pre-Delivery

DVT Risk and Prophylaxis Tool (Single Response)

VTE/DVT Risk Definitions

URL:

"\\epic-nas.et0922.epichosted.com\\static\\OrderSets\\VTE Risk Assessment Tool v7_MAK FINAL.pdf"

VERY LOW Risk of DVT (Selection Required)

"And" Linked Panel

Very low risk of VTE Routine, Once, L&D Pre-Delivery

Ambulate Routine, 3 times daily
Specify:
Early ambulation, L&D Pre-Delivery

Avoid dehydration Routine, Until discontinued, Starting S, L&D Pre-Delivery

LOW Risk of DVT (Selection Required)

Anticipated admission LESS than or EQUAL to 72 hours.

Does not meet moderate or high risk criteria:

Moderate Risk High Risk

Anticipated or actual LOS admission GREATER than or EQUAL to 72 hours High risk thrombophilia with no prior VTE

Prior idiopathic, or estrogen related VTE

Low risk thrombophilia AND family history of VTE OR single prior VTE

Receiving outpatient prophylactic LMWH or UFH

Low risk of VTE Routine, Once
Low risk: Due to low risk, SCDs are recommended while in bed and until fully ambulatory
L&D Pre-Delivery

Place sequential compression device (Single Response)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
L&D Pre-Delivery

Place/Maintain sequential compression device continuous Routine, Continuous
While in bed AND until fully ambulatory. Encourage early ambulation.
Avoid dehydration., L&D Pre-Delivery

MODERATE Risk of DVT (Selection Required)

Anticipated or actual LOS admission GREATER than 72 hours; does not meet High risk criteria. CONSIDER prophylactic LMWH/UFH (consult Anesthesia for delivery considerations)

High Risk

High risk thrombophilia with no prior VTE

Prior idiopathic, or estrogen related VTE

Low risk thrombophilia AND family history of VTE OR single prior VTE

Receiving outpatient prophylactic LMWH or UFH

Moderate Risk (Selection Required)

Moderate risk of VTE Routine, Once, L&D Pre-Delivery

Mechanical Prophylaxis (Single Response) (Selection Required)

Contraindications exist for mechanical prophylaxis Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
L&D Pre-Delivery

Place/Maintain sequential compression device continuous Routine, Continuous
SCD throughout hospitalization. Encourage early ambulation. Avoid dehydration., L&D Pre-Delivery

HIGH Risk of DVT (Selection Required)

High risk thrombophilia with no prior VTE

Prior idiopathic or estrogen related VTE

Low risk thrombophilia AND (family history of VTE OR single prior VTE)

Receiving outpatient prophylactic LMWH or UFH

High Risk (Selection Required)

<input type="checkbox"/>	High risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous While in bed AND until fully ambulatory. Encourage early ambulation. Avoid dehydration., L&D Pre-Delivery
<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)	
<input type="checkbox"/>	Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/>	Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/>	Moderate risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery
<input type="checkbox"/>	High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/>	High risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/>	Place sequential compression device (Single Response)	
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery

<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: L&D Pre-Delivery
<input type="checkbox"/> Place sequential compression device (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): L&D Pre-Delivery
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, L&D Pre-Delivery

Labs

COVID-19 Qualitative PCR

<input type="checkbox"/> COVID-19 qualitative RT-PCR - Nasal Swab	STAT For 1 Occurrences Specimen Source: Nasal Swab Is this for pre-procedure or non-PUI assessment? Yes Please select a reason for ordering, if applicable. Laboring patient Please select a reason for ordering, if applicable. Laboring patient L&D Pre-Delivery
---	---

Labs HMM

<input type="checkbox"/> Rubella antibody, IgG	Once, L&D Pre-Delivery
<input type="checkbox"/> Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/> Urine drugs of abuse screen	Once, L&D Pre-Delivery
<input type="checkbox"/> Bedside glucose	Routine, Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> OB Panel	
<input checked="" type="checkbox"/> CBC with differential	Once, L&D Pre-Delivery
<input type="checkbox"/> Basic metabolic panel	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> Hepatitis B surface antigen	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input type="checkbox"/> Urine dipstick	Once, L&D Pre-Delivery

<input checked="" type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Fetal Demise Panel	
<input type="checkbox"/> Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/> Antithrombin III level	Once, L&D Pre-Delivery
<input type="checkbox"/> Cardiolipin antibodies	Once, L&D Pre-Delivery
<input type="checkbox"/> Factor V leiden by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/> Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/> Homocysteine, plasma	Once, L&D Pre-Delivery
<input type="checkbox"/> Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/> Lupus anticoagulant panel	Once, L&D Pre-Delivery
<input type="checkbox"/> Parvovirus B19 antibody, IgG and IgM	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin mutation, factor II, by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/> TSH	Once, L&D Pre-Delivery

Labs HMW

<input type="checkbox"/> Blood gas, arterial, cord	Once, L&D Pre-Delivery
<input type="checkbox"/> Blood gas, venous, cord	Once, L&D Pre-Delivery
<input type="checkbox"/> Rubella antibody, IgG	Once, L&D Pre-Delivery
<input type="checkbox"/> Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/> Urine drugs of abuse screen	Once, L&D Pre-Delivery
<input type="checkbox"/> Bedside glucose	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> OB Panel	
<input type="checkbox"/> Bedside glucose	Routine, Every hour, L&D Pre-Delivery
<input type="checkbox"/> CBC with platelet and differential	Once, L&D Pre-Delivery

<input type="checkbox"/>	Basic metabolic panel	Once, L&D Pre-Delivery
<input type="checkbox"/>	Hepatitis B surface antigen	Once, L&D Pre-Delivery
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/>	Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/>	Fetal Demise Panel	
<input type="checkbox"/>	Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/>	Antithrombin III level	Once, L&D Pre-Delivery
<input type="checkbox"/>	Cardiolipin antibodies	Once, L&D Pre-Delivery
<input type="checkbox"/>	Factor V leiden by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/>	Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/>	Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/>	Homocysteine, plasma	Once, L&D Pre-Delivery
<input type="checkbox"/>	Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/>	Lupus anticoagulant panel	Once, L&D Pre-Delivery
<input type="checkbox"/>	Parvovirus B19 antibody, IgG and IgM	Once, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin mutation, factor II, by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/>	Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/>	TSH	Once, L&D Pre-Delivery
Labs HMSL		
<input type="checkbox"/>	Blood gas, arterial, cord	Once, L&D Pre-Delivery
<input type="checkbox"/>	Blood gas, venous, cord	Once, L&D Pre-Delivery
<input type="checkbox"/>	Rubella antibody, IgG	Once, L&D Pre-Delivery

<input type="checkbox"/> Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/> Urine drugs of abuse screen	Once, L&D Pre-Delivery
<input type="checkbox"/> Bedside glucose	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> OB Panel	
<input type="checkbox"/> Bedside glucose	Routine, Every hour, L&D Pre-Delivery
<input type="checkbox"/> CBC with platelet and differential	Once, L&D Pre-Delivery
<input type="checkbox"/> Basic metabolic panel	Once, L&D Pre-Delivery
<input type="checkbox"/> Hepatitis B surface antigen	Once, L&D Pre-Delivery
<input type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Fetal Demise Panel	
<input type="checkbox"/> Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/> Antithrombin III level	Once, L&D Pre-Delivery
<input type="checkbox"/> Cardiolipin antibodies	Once, L&D Pre-Delivery
<input type="checkbox"/> Factor V leiden by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/> Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/> Homocysteine, plasma	Once, L&D Pre-Delivery
<input type="checkbox"/> Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/> Lupus anticoagulant panel	Once, L&D Pre-Delivery
<input type="checkbox"/> Parvovirus B19 antibody, IgG and IgM	Once, L&D Pre-Delivery

<input type="checkbox"/> Prothrombin mutation, factor II, by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/> TSH	Once, L&D Pre-Delivery

Labs HMB

<input type="checkbox"/> Cord blood gas, arterial	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Cord blood gas, venous	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Rubella antibody, IgG	Once, L&D Pre-Delivery
<input type="checkbox"/> Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/> Urine drugs of abuse screen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Bedside glucose	Routine, Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> OB Panel	
<input checked="" type="checkbox"/> CBC with differential	Once, L&D Pre-Delivery
<input type="checkbox"/> Basic metabolic panel	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> Hepatitis B surface antigen	Once, L&D Pre-Delivery
<input type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Fetal Demise Panel	
<input type="checkbox"/> Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/> Antithrombin III level	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Cardiolipin antibodies	Once, L&D Pre-Delivery

<input type="checkbox"/> Factor V leiden by PCR	STAT For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/> Homocysteine, plasma	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/> Lupus anticoagulant panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Parvovirus B19 antibody, IgG and IgM	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin mutation, factor II, by PCR	STAT For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> TSH	STAT For 1 Occurrences, L&D Pre-Delivery

Labs HMCL

<input type="checkbox"/> Blood gas, arterial, cord	Once, L&D Pre-Delivery
<input type="checkbox"/> Blood gas, venous, cord	Once, L&D Pre-Delivery
<input type="checkbox"/> Rubella antibody, IgG	Once, L&D Pre-Delivery
<input type="checkbox"/> Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/> Urine drugs of abuse screen	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> OB Panel	
<input checked="" type="checkbox"/> CBC with differential	Once, L&D Pre-Delivery
<input type="checkbox"/> Basic metabolic panel	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Hepatitis B surface antigen	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input type="checkbox"/> Urine dipstick	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery

<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Fetal Demise Panel	
<input type="checkbox"/>	Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/>	Antithrombin III level	Once, L&D Pre-Delivery
<input type="checkbox"/>	Cardiolipin antibodies	Once, L&D Pre-Delivery
<input type="checkbox"/>	Factor V leiden by PCR	Once
		Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/>	Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/>	Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/>	Homocysteine, plasma	Once, L&D Pre-Delivery
<input type="checkbox"/>	Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/>	Lupus anticoagulant panel	Once, L&D Pre-Delivery
<input type="checkbox"/>	Parvovirus B19 antibody, IgG and IgM	Once, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin mutation, factor II, by PCR	Once
		Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/>	Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/>	TSH	Once, L&D Pre-Delivery

Labs HMTW

<input type="checkbox"/>	Blood gas, arterial, cord	Once, L&D Pre-Delivery
<input type="checkbox"/>	Blood gas, venous, cord	Once, L&D Pre-Delivery
<input type="checkbox"/>	Rubella antibody, IgG	Once, L&D Pre-Delivery
<input type="checkbox"/>	Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/>	Urine drugs of abuse screen	Once, L&D Pre-Delivery
<input type="checkbox"/>	Bedside glucose	Routine, Once, L&D Pre-Delivery
<input checked="" type="checkbox"/>	OB Panel	
<input checked="" type="checkbox"/>	CBC with differential	Once, L&D Pre-Delivery
<input type="checkbox"/>	Basic metabolic panel	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/>	Hepatitis B surface antigen	Once, L&D Pre-Delivery
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once
		Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once
		Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery

<input checked="" type="checkbox"/> Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Fetal Demise Panel	
<input type="checkbox"/> Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/> Antithrombin III level	Once, L&D Pre-Delivery
<input type="checkbox"/> Cardiolipin antibodies	Once, L&D Pre-Delivery
<input type="checkbox"/> Factor V leiden by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/> Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/> Homocysteine, plasma	Once, L&D Pre-Delivery
<input type="checkbox"/> Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/> Lupus anticoagulant panel	Once, L&D Pre-Delivery
<input type="checkbox"/> Parvovirus B19 antibody, IgG and IgM	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin mutation, factor II, by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/> TSH	Once, L&D Pre-Delivery

Labs HMWB

<input type="checkbox"/> Rubella antibody, IgG	Once, L&D Pre-Delivery
<input type="checkbox"/> Surgical pathology request	Collection Date: 1/4/2024 Collection Time: Surgical Specimen: Gestational Age: Specimen Site: Number of specimens: Malignancy: Irradiation: Specimen Status: Chemotherapy: Hormonal Therapy: Clinical History: Pre-Operative Diagnosis: Procedure Type: L&D Pre-Delivery
<input type="checkbox"/> Urine drugs of abuse screen	Once, L&D Pre-Delivery
<input type="checkbox"/> Bedside glucose	Routine, Once, L&D Pre-Delivery
<input checked="" type="checkbox"/> OB Panel	
<input checked="" type="checkbox"/> CBC with differential	Once For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Basic metabolic panel	Once For 1 Occurrences, L&D Pre-Delivery

<input checked="" type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Hepatitis B surface antigen	Once For 1 Occurrences, L&D Pre-Delivery
<input checked="" type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input checked="" type="checkbox"/> Type and screen, obstetrical patient	Once, L&D Pre-Delivery
<input type="checkbox"/> Urine dipstick	Once For 1 Occurrences, L&D Pre-Delivery
<input checked="" type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once For 1 Occurrences Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> LDH	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, Post-op
<input type="checkbox"/> Fetal Demise Panel	
<input type="checkbox"/> Antibody screen (gel)	Once, L&D Pre-Delivery
<input type="checkbox"/> Antithrombin III level	Once, L&D Pre-Delivery
<input type="checkbox"/> Cardiolipin antibodies	Once, L&D Pre-Delivery
<input type="checkbox"/> Factor V leiden by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/> Hemoglobin A1c	Once, L&D Pre-Delivery
<input type="checkbox"/> Homocysteine, plasma	Once, L&D Pre-Delivery
<input type="checkbox"/> Kleihauer-Betke	Once, L&D Pre-Delivery
<input type="checkbox"/> Lupus anticoagulant panel	Once, L&D Pre-Delivery
<input type="checkbox"/> Parvovirus B19 antibody, IgG and IgM	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin mutation, factor II, by PCR	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/> TSH	Once, L&D Pre-Delivery

24 Hour urine

<input type="checkbox"/> 24 Hour urine	
<input type="checkbox"/> Creatinine clearance, urine, 24 hour	Once, L&D Pre-Delivery
<input type="checkbox"/> Protein, urine, 24 hour	Once, L&D Pre-Delivery

Magnesium and D-dimer

<input type="checkbox"/> D-dimer	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Magnesium level	STAT For 1 Occurrences, L&D Pre-Delivery

Respiratory

Oxygen

Oxygen therapy

Routine, As needed
Device: Non-rebreather mask
Rate in liters per minute:
Titrate to keep O2 Sat Above: Other (Specify)
Specify titration to keep O2 Sat (%) Above: 94
Indications for O2 therapy: Fetal indication
Continue O2 for 30 minutes per event., L&D Pre-Delivery

Consults

For Physician Consult orders use sidebar

Physician Consults

Consult Anesthesiology

Reason for Consult?
Patient/Clinical information communicated?
Patient/clinical information communicated?
L&D Pre-Delivery

Consult Maternal and Fetal Medicine

Reason for Consult?
Patient/Clinical information communicated?
Patient/clinical information communicated?
L&D Pre-Delivery

Consult Neonatology

Reason for Consult?
Patient/Clinical information communicated?
Patient/clinical information communicated?
L&D Pre-Delivery

Ancillary consults

Consult to PT eval and treat

Reasons for referral to Physical Therapy (mark all applicable):
Are there any restrictions for positioning or mobility?
Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal):
Weight Bearing Status:
L&D Pre-Delivery

Consult to Social Work

Reason for Consult:
L&D Pre-Delivery

Consult to Spiritual Care

Reason for consult?
L&D Pre-Delivery