

Enhanced Recovery After Surgery (ERAS) Orders

[Guidelines for Antenatal and Preoperative care in Cesarean Delivery: Enhanced Recovery After Surgery Society Recommendations](#)

URL:
"https://www.clinicalkey.com/#!/content/playContent/1-s2.0-S0002937818307634?returnurl=https:%2F%2Flinkinghub.elsevier.com%2Fretrieve%2Fpii%2FS0002937818307634%3Fshowall%3Dtrue&referrer="

[Practice Guidelines for Preoperative Fasting and the Use of Pharmacologic Agents](#)

URL:
"https://insights.ovid.com/crossref?an=00000542-20170300-00014"

Cesarean Section ERAS Diet and Nutrition PreOP

Click here for ACOG Committee Opinion

URL:
"\\epic-nas.et0922.epichosted.com\static\OrderSets\Perioperative pathways enhanced recovery after surgery.pdf"

Click here for ERAS Guidelines

URL:
"\\epic-nas.et0922.epichosted.com\static\OrderSets\Guidelines for Antenatal and Preoperative care in Cesarean Delivery.pdf"

Click here for Preoperative Fasting Guidelines

URL:
"\\epic-nas.et0922.epichosted.com\static\OrderSets\Practice Guidelines for Preoperative Fasting.pdf"

NPO except clear liquids, start 8 hours prior to surgery

Diet effective now, Starting S
NPO:
Pre-Operative fasting options:
An NPO order without explicit exceptions means nothing can be given orally to the patient., Pre-op

Clear liquids can be continued until 2 hours prior to surgery

Diet effective now, Starting S
Diet(s): Clear Liquids
Other Options:
Advance Diet as Tolerated?
IDDSI Liquid Consistency:
Fluid Restriction:
Foods to Avoid:
Pre-op

Provide 1 bottle Ensure Pre-Surgery Clear to all non-diabetics at PAT visit. Instruct Patient to complete by 2 hours prior to surgery.

Routine, Once For 1 Occurrences, Pre-op

acetaminophen (TYLENOL) oral

acetaminophen (TYLENOL) tablet

1,000 mg, oral, once, For 1 Doses, Pre-op to be given 1 hour before start of surgery

acetaminophen ER (TYLENOL) 8 hr tablet

1,300 mg, oral, once, For 1 Doses, Pre-op to be given 1 hour before start of surgery

General

Pre Anesthesia Testing Orders

The ambulatory orders in this section are specifically for Pre Anesthesia Testing. For additional PAT orders, please use 'Future Status' and 'Pre-Admission Testing Phase of Care'

Other Diagnostic Studies

ECG Pre/Post Op

Routine, Ancillary Performed, Status: Future, Expires: S+366, Pre-Admission Testing

Pv carotid duplex

Status: Future, Expires: S+366, Routine, Clinic Performed, Pre-Admission Testing

Us vein mapping lower extremity

Status: Future, Expires: S+366, Routine, Clinic Performed, Pre-Admission Testing

MRSA PCR

Routine, Status: Future, Expires: S+366, Clinic Collect, Nares, Pre-Admission Testing

Us duplex venous lower extremity Status: Future, Expires: S+366, Routine, Clinic Performed, Pre-Admission Testing

Respiratory

Spirometry pre & post w/ bronchodilator, diffusion, lung volumes Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry, diffusion, lung volumes Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry pre & post w/ bronchodilator Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Encounter type?

Body Plethysmographic lung volumes Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Encounter type?

Spirometry Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Encounter type?

OP Diffusion Capacity Combination Panel (Single Response)

Spirometry, diffusion Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry, diffusion, lung volumes Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry, diffusion, MIPS/MEPS Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry, diffusion, lung volumes, MIPS/MEPS Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry pre & post w/ bronchodilator, diffusion Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry pre & post w/ bronchodilator, diffusion, lung volumes Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry pre & post w/ bronchodilator, diffusion, MIPS/MEPS Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Spirometry pre & post w/ bronchodilator, diffusion, lung volumes, MIPS/MEPS Routine, Status: Future, Expires: S+366, Pre-Admission Testing
Location:
Encounter type?

Laboratory: Preoperative Testing Labs

COVID-19 qualitative RT-PCR - Nasal Swab Routine, Status: Future, Expires: S+366, Clinic Collect
Specimen Source: Nasal Swab
Is this for pre-procedure or non-PUI assessment? Yes
Pre-Admission Testing

CBC with platelet and differential Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Comprehensive metabolic panel Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Basic metabolic panel Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Prothrombin time with INR Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Partial thromboplastin time Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Hepatic function panel Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Platelet function analysis Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Hemoglobin A1c Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

<input type="checkbox"/>	Type and screen	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	hCG qualitative, serum screen	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/>	POC pregnancy, urine	Routine, Point Of Care, Pre-Admission Testing
<input type="checkbox"/>	Urinalysis, automated with microscopy	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/> Laboratory: Additional Labs		
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Routine, Status: Future, Expires: S+366, Clinic Collect Specimen Source: Urine Specimen Site: Clean catch Pre-Admission Testing
<input type="checkbox"/>	CBC hemogram	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/>	Acute viral hepatitis panel (HAV, HBV, HCV)	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Thromboelastograph	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Vitamin D 25 hydroxy level	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Methicillin-resistant staphylococcus aureus (MRSA), NAA	Routine, Status: Future, Expires: S+366, Clinic Collect, Nares, Pre-Admission Testing
<input type="checkbox"/>	T3	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	T4	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Thyroid stimulating hormone	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Prostate specific antigen	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/> Laboratory: OB Additional Labs HMB		
<input type="checkbox"/>	COVID-19 qualitative RT-PCR	Routine, Status: Future, Expires: S+366, Clinic Collect Specimen Source: Is this for pre-procedure or non-PUI assessment? Pre-Admission Testing
<input type="checkbox"/>	CBC with platelet and differential	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Hepatitis B surface antigen	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/>	Type and screen, obstetrical patient	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Rubella Ab IgG	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Routine, Status: Future, Expires: S+366, Clinic Collect Specimen Source: Urine Specimen Site: Clean catch Pre-Admission Testing
<input type="checkbox"/>	Urine drugs of abuse screen	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Basic metabolic panel	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with platelet and differential	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Comprehensive metabolic panel	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Prothrombin time with INR	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Partial thromboplastin time, activated	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Fibrinogen	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Uric acid level	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	LDH	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Protein, urine, random	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Laboratory: Additional for Bariatric patients	
<input type="checkbox"/>	Lipid panel	Routine, Status: Future, Expires: S+366, Clinic Collect Has the patient been fasting for 8 hours or more? Pre-Admission Testing
<input type="checkbox"/>	hCG qualitative, serum screen	Routine, Status: Future, Expires: S+366, Clinic Collect Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-Admission Testing
<input type="checkbox"/>	Total iron binding capacity	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	T4, free	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Thyroid stimulating hormone	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Hemoglobin A1c	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Parathyroid hormone	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	CBC with platelet and differential	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Prothrombin time with INR	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Partial thromboplastin time, activated	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Vitamin A level, plasma or serum	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Vitamin B12 level	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/>	Vitamin D 25 hydroxy level	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

<input type="checkbox"/> Copper level, serum	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/> Folate level	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/> Vitamin B1 (thiamine)	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing
<input type="checkbox"/> Zinc level, serum	Routine, Status: Future, Expires: S+366, Clinic Collect, Pre-Admission Testing

Case Request (Single Response)

<input type="checkbox"/> DELIVERY, CESAREAN	Panel 1 DELIVERY, CESAREAN, Scheduling/ADT, Scheduling/ADT
<input type="checkbox"/> C-SECTION + BILATERAL TUBAL LIGATION	Panel 1 C-SECTION + BILATERAL TUBAL LIGATION, Scheduling/ADT, Scheduling/ADT
<input type="checkbox"/> Case request operating room	Scheduling/ADT, Scheduling/ADT

Inpatient only procedure (Single Response) (Selection Required)

<input type="checkbox"/> Admit to L&D	Admitting Physician: Bed request comments: Pre-op
---------------------------------------	---

Precautions

<input type="checkbox"/> Aspiration precautions	Pre-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Pre-op
<input type="checkbox"/> Latex precautions	Pre-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Pre-op

Nursing

Vital signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol, Pre-op
--	------------------------------------

Activity

<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S, Pre-op
<input checked="" type="checkbox"/> Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: Pre-op
<input type="checkbox"/> Ambulate with assistance	Routine, 3 times daily Specify: with assistance Pre-op
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated Pre-op

Nursing care

<input checked="" type="checkbox"/> Monitor fetal heart tones	Routine, Once Type: Intermittent With non stress test. Obtain 30 min fetal heart tracing. If FHR non-reassuring, continue monitoring and notify physician., Pre-op
<input checked="" type="checkbox"/> Fetal nonstress test	Routine, Once, Pre-op
<input checked="" type="checkbox"/> Insert and maintain Foley	
<input checked="" type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: May insert in OR or after regional anesthesia effective, Pre-op

<input checked="" type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain Pre-op
<input checked="" type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, Pre-op
<input checked="" type="checkbox"/> Place antiembolic stockings	Routine, Until discontinued, Starting S, Pre-op
Diet	
<input checked="" type="checkbox"/> NPO	Diet effective now, Starting S NPO: Except Sips of clear liquids Pre-Operative fasting options: Clear Liquid intake is acceptable up to two hours before surgery, An NPO order without explicit exceptions means nothing can be given orally to the patient., Pre-op
<input type="checkbox"/> Diet - Clear Liquids	Diet effective now, Starting S Diet(s): Clear Liquids Other Options: Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Clear Liquid intake is acceptable up to two hours before surgery, L&D Pre-Delivery
Consent	
<input type="checkbox"/> Complete consent for Primary Cesarean Section	Routine, Once Procedure: Primary Cesarean Section Diagnosis/Condition: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Consent for Primary Cesarean Section, Pre-op
<input type="checkbox"/> Complete consent for Primary Cesarean Section with Bilateral Tubal Ligation	Routine, Once Procedure: Primary Cesarean Section with Bilateral Tubal Ligation Diagnosis/Condition: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Consent for Primary Cesarean Section with Bilateral Tubal Ligation, Pre-op
<input type="checkbox"/> Complete consent for Primary Cesarean Section with Bilateral Salpingectomy	Routine, Once Procedure: Primary Cesarean Section with Bilateral Salpingectomy Diagnosis/Condition: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Consent for Primary Cesarean Section with Bilateral Salpingectomy, Pre-op
<input type="checkbox"/> Complete consent for Repeat Cesarean Section	Routine, Once Procedure: Repeat Cesarean Section Diagnosis/Condition: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Consent for Repeat Cesarean Section, Pre-op

<input type="checkbox"/> Complete consent for Repeat Cesarean Section with Bilateral Tubal Ligation	Routine, Once Procedure: Repeat Cesarean Section with Bilateral Tubal Ligation Diagnosis/Condition: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Consent for Repeat Cesarean Section with Bilateral Tubal Ligation, Pre-op
<input type="checkbox"/> Complete consent for Repeat Cesarean Section with Bilateral Salpingectomy	Routine, Once Procedure: Repeat Cesarean Section with Bilateral Salpingectomy Diagnosis/Condition: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Consent for Repeat Cesarean Section with Bilateral Salpingectomy, Pre-op

IV Fluids

IV Fluids

<input checked="" type="checkbox"/> lactated ringer's bolus and infusion	"And" Linked Panel
<input checked="" type="checkbox"/> lactated Ringer's bolus	1,000 mL, intravenous, Administer over: 30 Minutes, once PRN, If patient requests epidural - for epidural prehydration, Pre-op Notify Anesthesiologist immediately if patient requests Epidural and begin pre-epidural hydration
<input checked="" type="checkbox"/> lactated Ringer's infusion	125 mL/hr, intravenous, continuous, Pre-op

Insert and Maintain IV

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once, Pre-op
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled, Pre-op
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care, Pre-op

Local Anesthetic with Venipuncture

<input type="checkbox"/> buffered lidocaine 1% injection	0.15 mL, intravenous, once, For 1 Doses, Pre-op
--	---

Local Anesthetic with Venipuncture

<input type="checkbox"/> lidocaine PF 1% (XYLOCAINE) injection	0.15 mL, injection, once, For 1 Doses, Pre-op
--	---

Medications

PreOp Antibiotics: For Patients LESS than or EQUAL to 120 kg **Give 60 minutes PRIOR to C-Section (Single Response)**

<input checked="" type="checkbox"/> Pre-op Antibiotics	
<input checked="" type="checkbox"/> ceFAZolin (ANCEF) IV - Give within 60 minutes prior to C-Section	2 g, intravenous, once, For 1 Doses, Pre-op Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> azithromycin (ZITHROMAX) IV - Give within 60 minutes prior to C-Section	500 mg, intravenous, Administer over: 60 Minutes, once, For 1 Doses, Pre-op Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:

() If Penicillin Allergic: clindamycin (CLEOCIN) IV and gentamicin (GARAMYCIN) IV		"And" Linked Panel
<input type="checkbox"/> clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Pre-op	Nurse to send medication(s) to operating room - To be administered by Anesthesiologist.To be given 1 hour prior to skin incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> If Penicillin Allergic - gentamicin (GARAMYCIN) IVPB - Give within 60 minutes prior to C-Section	5 mg/kg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Pre-op	Nurse to send medication(s) to operating room - To be administered by Anesthesiologist.To be given 1 hour prior to skin incision. Reason for Therapy: Surgical Prophylaxis

PreOp Antibiotics: For Patients GREATER than 120 kg **Give 60 minutes PRIOR to C-Section (Single Response)**

(X) Pre-op Antibiotics		
<input checked="" type="checkbox"/> ceFAZolin (ANCEF) IV - Give within 60 minutes prior to C-Section	3 g, intravenous, once, For 1 Doses, Pre-op	Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> azithromycin (ZITHROMAX) IV - Give within 60 minutes prior to C-Section	500 mg, intravenous, Administer over: 60 Minutes, once, For 1 Doses, Pre-op	Give within 60 minutes prior to C-Section Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied: Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:

() If Penicillin Allergic: clindamycin (CLEOCIN) IV and gentamicin (GARAMYCIN) IV		"And" Linked Panel
<input type="checkbox"/> clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Pre-op	Nurse to send medication(s) to operating room - To be administered by Anesthesiologist.To be given 1 hour prior to skin incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> If Penicillin Allergic - gentamicin (GARAMYCIN) IVPB - Give within 60 minutes prior to C-Section	5 mg/kg, intravenous, Administer over: 30 Minutes, once, For 1 Doses, Pre-op	Nurse to send medication(s) to operating room - To be administered by Anesthesiologist.To be given 1 hour prior to skin incision. Reason for Therapy: Surgical Prophylaxis

Antibiotics (Single Response)

Does your patient have a penicillin allergy?

() No (Single Response)
Please select the appropriate regimen

() Prophylaxis Regimen for GBS - penicillin G IVPB Loading and Maintenance Dose		"Followed by" Linked Panel
<input type="checkbox"/> Loading Dose - penicillin G (POTASSIUM) IV	5 Million Units, intravenous, Administer over: 30 Minutes, once, For 1 Doses, L&D Pre-Delivery	If GBS positive Indication: Medical Prophylaxis
<input type="checkbox"/> Maintenance Dose - penicillin G (POTASSIUM) IV	2.5 Million Units, intravenous, Administer over: 30 Minutes, every 4 hours, Starting H+4 Hours, L&D Pre-Delivery	If GBS positive Indication: Medical Prophylaxis
() Alternative Regimen for GBS - ampicillin IVPB Loading and Maintenance Dose		"Followed by" Linked Panel

<input type="checkbox"/> Loading Dose - ampicillin IV	2 g, intravenous, Administer over: 30 Minutes, once, For 1 Doses, L&D Pre-Delivery If GBS positive Reason for Therapy: Medical Prophylaxis
<input type="checkbox"/> Maintenance Dose - ampicillin IV	1 g, intravenous, Administer over: 30 Minutes, every 6 hours, Starting H+6 Hours, L&D Pre-Delivery If GBS positive Reason for Therapy: Medical Prophylaxis

Yes (Single Response)

Is your patient LOW Risk or HIGH Risk for Anaphylaxis?

LOW Risk

"Followed by" Linked Panel

Recommended for patients NOT high risk for anaphylaxis

<input type="checkbox"/> Loading Dose - cefazolin (ANCEF) IV	2,000 mg, intravenous, once, For 1 Doses Reason for Therapy: Medical Prophylaxis Medical Prophylaxis: The maximum duration of therapy that can be entered at this time is 56 days. Order can be renewed if required at that time.
<input type="checkbox"/> Maintenance Dose - cefazolin (ANCEF) IV	1,000 mg, intravenous, every 8 hours Through delivery then discontinue. Reason for Therapy: Medical Prophylaxis Medical Prophylaxis: The maximum duration of therapy that can be entered at this time is 56 days. Order can be renewed if required at that time.

HIGH Risk (Single Response)

clindamycin (CLEOCIN) IV Loading and Maintenance Doses

"Followed by" Linked Panel

Recommended ONLY for patients with high risk for penicillin anaphylaxis that are culture isolate sensitive to Clindamycin.

<input type="checkbox"/> Loading Dose - clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, once, For 1 Doses Reason for Therapy: Medical Prophylaxis
<input type="checkbox"/> Maintenance Dose - clindamycin (CLEOCIN) IV	900 mg, intravenous, Administer over: 30 Minutes, every 8 hours Through delivery then discontinue. Reason for Therapy: Medical Prophylaxis
<input type="checkbox"/> If clindamycin-resistant GBS - vancomycin (VANCOCIN) 20 mg/kg IV + Pharmacy Consult to Dose (Selection Required)	
<input type="checkbox"/> vancomycin (VANCOCIN) IV	20 mg/kg, intravenous, every 8 hours, L&D Pre-Delivery Reason for Therapy: Reason for Therapy: Medical Prophylaxis
<input type="checkbox"/> Pharmacy consult to manage vancomycin	STAT, Until discontinued, Starting S Reason for Therapy: Duration of Therapy (Days):

Pre-Anesthesia Medications

<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, once PRN, Decrease gastric acidity, Pre-op Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:
<input type="checkbox"/> sodium citrate-citric acid (BICITRA) solution	30 mL, oral, once PRN, Decrease gastric acidity, Pre-op
<input type="checkbox"/> metoclopramide (REGLAN) injection	10 mg, intravenous, once PRN, nausea, Decrease gastric acidity, Pre-op Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:

Labs

COVID-19 Qualitative PCR

<input type="checkbox"/> COVID-19 qualitative RT-PCR - Nasal Swab	STAT For 1 Occurrences Specimen Source: Nasal Swab Is this for pre-procedure or non-PUI assessment? Yes Pre-op
---	---

Labs HMWB

<input type="checkbox"/> Bedside glucose	Routine, Once For 1 Occurrences, Pre-op
<input type="checkbox"/> OB Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/> Hepatitis B surface antigen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	STAT For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/> Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/> Urine dipstick	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences Specimen Source: Urine Specimen Site: Pre-op
<input type="checkbox"/> No prenatal records	
<input type="checkbox"/> Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, Pre-op

Labs HMW

<input type="checkbox"/> Bedside glucose	Routine, Once, Pre-op
<input type="checkbox"/> OB Panel	
<input type="checkbox"/> CBC with differential	Once, Pre-op
<input type="checkbox"/> Basic metabolic panel	Once, Pre-op
<input type="checkbox"/> Hepatitis B surface antigen	Once, Pre-op
<input type="checkbox"/> HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/> Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/> Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: Pre-op
<input type="checkbox"/> No prenatal records	

<input type="checkbox"/>	Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Pre-op

Labs HMSL

<input type="checkbox"/>	Bedside glucose	Routine, Once, Pre-op
<input type="checkbox"/>	OB Panel	
<input type="checkbox"/>	CBC with differential	Once, Pre-op
<input type="checkbox"/>	Basic metabolic panel	Once, Pre-op
<input type="checkbox"/>	Hepatitis B surface antigen	Once, Pre-op
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: Pre-op

<input type="checkbox"/>	No prenatal records	
<input type="checkbox"/>	Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Pre-op

Labs HMB

<input type="checkbox"/>	Bedside glucose	Routine, Once, Pre-op
<input type="checkbox"/>	OB Panel	
<input type="checkbox"/>	CBC with differential	Once, Pre-op
<input type="checkbox"/>	Basic metabolic panel	Once, Pre-op
<input type="checkbox"/>	Hepatitis B surface antigen	Once, Pre-op
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op

<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: Pre-op
<input type="checkbox"/>	No prenatal records	
<input type="checkbox"/>	Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Pre-op

Labs HMCL

<input type="checkbox"/>	Bedside glucose	Routine, Once, Pre-op
<input type="checkbox"/>	OB Panel	
<input type="checkbox"/>	CBC with differential	Once, Pre-op
<input type="checkbox"/>	Basic metabolic panel	Once, Pre-op
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Hepatitis B surface antigen	Once, Pre-op
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/>	Urine dipstick	Once, Pre-op
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: Pre-op
<input type="checkbox"/>	No prenatal records	
<input type="checkbox"/>	Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Pre-op

Labs HMTW

<input type="checkbox"/>	Bedside glucose	Routine, Once, Pre-op
<input type="checkbox"/>	OB Panel	
<input type="checkbox"/>	CBC with differential	Once, Pre-op
<input type="checkbox"/>	Basic metabolic panel	Once, Pre-op
<input type="checkbox"/>	Hepatitis B surface antigen	Once, Pre-op
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: Pre-op
<input type="checkbox"/>	No prenatal records	
<input type="checkbox"/>	Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Pre-op

Labs HMM

<input type="checkbox"/>	Bedside glucose	Routine, Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	OB Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Basic metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Hepatitis B surface antigen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	HIV 1/2 antigen/antibody, fourth generation, with reflexes	Once For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Syphilis treponema screen with RPR confirmation (reverse algorithm)	Once Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): Pre-op
<input type="checkbox"/>	Type and screen, obstetrical patient	Once, Pre-op
<input type="checkbox"/>	Urine dipstick	Once, Pre-op
<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences Specimen Source: Urine Specimen Site: Pre-op
<input type="checkbox"/>	No prenatal records	
<input type="checkbox"/>	Rubella antibody, IgG	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine drugs of abuse screen	Once, Pre-op
<input type="checkbox"/>	Pre-Eclamptic Lab Panel	
<input type="checkbox"/>	CBC with differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op

<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Uric acid	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	LDH	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Urine Protein and Creatinine	
<input type="checkbox"/>	Creatinine level, urine, random	Once For 1 Occurrences, Pre-op
<input type="checkbox"/>	Protein, urine, random	Once For 1 Occurrences, Pre-op

24 Hour urine

<input type="checkbox"/>	24 Hour Urine	
<input type="checkbox"/>	Creatinine clearance, urine, 24 hour	Once, Pre-op
<input type="checkbox"/>	Protein, urine, 24 hour	Once, Pre-op

Consults

For Physician Consult orders use sidebar

Physician Consults

<input type="checkbox"/>	Consult Anesthesiology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? L&D Pre-Delivery
<input type="checkbox"/>	Consult Maternal and Fetal Medicine	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? L&D Pre-Delivery
<input type="checkbox"/>	Consult Neonatology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? L&D Pre-Delivery

Ancillary consults

<input type="checkbox"/>	Consult to PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: L&D Pre-Delivery
<input type="checkbox"/>	Consult to Social Work	Reason for Consult: L&D Pre-Delivery
<input type="checkbox"/>	Consult to Spiritual Care	Reason for consult? L&D Pre-Delivery