

Adult Anesthesia Pre-Op [1459]

[Click Here for Anesthesia Preoperative Testing Protocol](#)

URL:

"\\epic-nas.et0922.epichosted.com\\static\\OrderSets\\HM
Anesthesia Protocol 090619.pdf"

Nursing

Nursing

<input type="checkbox"/> Access Implanted Vascular Access Device	Routine, Once, Pre-op
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once, Pre-op

Labs and Testing

Labs and Testing

<input checked="" type="checkbox"/> Labs and Testing	Routine, Until discontinued, Starting S To be performed per Anesthesia Peri-Operative Testing Protocol, Pre-op
<input checked="" type="checkbox"/> Lactic acid, I-Stat	Once, Starting S For 1 Occurrences FOR ALL PATIENTS DIAGNOSED WITH RENAL FAILURE, Pre-op
<input checked="" type="checkbox"/> Bedside glucose	Routine, Once, Starting S For 1 Occurrences FOR ALL PATIENTS DIAGNOSED WITH DIABETES, Pre-op
<input checked="" type="checkbox"/> POC pregnancy, urine	Once, Starting S For 1 Occurrences Release to patient (Note: If manual release option is selected, result will auto release 10 days from finalization.): FOR ALL FEMALE PATIENTS OF CHILD BEARING AGE UNDER 55 OR HAVING MENSES, Pre-op

IV Fluids

Maintenance IV Fluids (Single Response)

<input type="checkbox"/> lactated Ringer's infusion	30 mL/hr, intravenous, continuous, Pre-op
<input type="checkbox"/> For patients on dialysis OR coming for dialysis access - sodium chloride 0.9 % infusion	30 mL/hr, intravenous, continuous, Pre-op
<input checked="" type="checkbox"/> For patients diagnosed with Renal Failure and/or CHF - sodium chloride 0.9 % infusion	500 mL, intravenous, at 30 mL/hr, PRN, If Patient diagnosed with Renal Failure and/or Congestive Heart Failure, Pre-op

Medications

Regional Block Medications

<input type="checkbox"/> BUPivacaine PF (MARCAINE) injection 0.25% or 0.5% (Single Response)	
<input type="checkbox"/> BUPivacaine (PF) (MARCAINE) 0.25 % (2.5 mg/mL) solution	150 mg, infiltration, once PRN, regional anesthesia block placement, Pre-op Infiltration dosing for regional anesthesia block placement
<input type="checkbox"/> BUPivacaine (PF) (MARCAINE) 0.5 % (5 mg/mL) injection	100 mg, infiltration, once PRN, regional anesthesia block placement, Pre-op Infiltration dosing for regional anesthesia block placement
<input type="checkbox"/> BUPivacaine liposome (EXPAREL) 1.3 % (13.3 mg/mL) injection	266 mg, infiltration, once PRN, regional anesthesia block placement, Pre-op Infiltration dosing for regional anesthesia block placement
<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25-100 mcg, intravenous, once PRN, other, for sedation during regional anesthesia block placement, For 1 Doses, Pre-op Dosing for regional anesthesia block placement Allowance for Patient Preference:
<input type="checkbox"/> MIDAZolam (PF) (VERSED) injection	1-2 mg, intravenous, once PRN, sedation, during regional anesthesia block placement, For 1 Doses, Pre-op Dosing for regional anesthesia block placement Indication(s): Sedation

<input type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1 %) injection	5 mL, infiltration, once PRN, regional anesthesia block placement, Pre-op Infiltration dosing for regional anesthesia block placement
<input type="checkbox"/> dexAMETHasone (DECADRON) injection	4 mg, Other, once PRN, regional anesthesia block placement, Pre-op Infiltration dosing for regional anesthesia block placement

Medications - MIDAZolam

<input type="checkbox"/> MIDAZolam (VERSED) injection	2 mg, intravenous, once, For 1 Doses, Pre-op Indication(s):
-------------------------------------------------------	----------------------------------------------------------------

GI Drugs: For Patients LESS than 65 years old

<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, once, For 1 Doses, Pre-op Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:
<input type="checkbox"/> ondansetron (ZOFRAN) IV	4 mg, intravenous, once, For 1 Doses, Pre-op
<input type="checkbox"/> metoclopramide (REGLAN) injection	10 mg, intravenous, once, For 1 Doses, Pre-op Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:
<input type="checkbox"/> scopolamine (TRANSDERM-SCOP) 1.5 mg - For Patients LESS than 65 years old	1 patch, transdermal, Administer over: 72 Hours, every 72 hours, Pre-op

GI Drugs: For Patients GREATER than or EQUAL to 65 years old

<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, once, For 1 Doses, Pre-op Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:
<input type="checkbox"/> ondansetron (ZOFRAN) IV	4 mg, intravenous, once, For 1 Doses, Pre-op
<input type="checkbox"/> metoclopramide (REGLAN) injection	10 mg, intravenous, once, For 1 Doses, Pre-op Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:

Pre-Operative Analgesics - HMM Only

<input checked="" type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate(1 mL) injection	0.5 mL, intradermal, once, For 1 Doses, Pre-op May use up to 0.5 milliliters for comfort. For PreOp Use Only
<input type="checkbox"/> traMADol (ULTRAM) tablet	150 mg, oral, once, For 1 Doses, Pre-op Allowance for Patient Preference:
<input type="checkbox"/> celecoxib (CeleBREX) capsule	400 mg, oral, once, For 1 Doses, Pre-op
<input type="checkbox"/> pregabalin (LYRICA) or gabapentin (NEURONTIN) oral doses - For Patients GREATER than or EQUAL to 65 years old (Single Response)	
<input type="checkbox"/> pregabalin (LYRICA) capsule	75 mg, oral, once, For 1 Doses, Pre-op
<input type="checkbox"/> gabapentin (NEURONTIN) capsule	600 mg, oral, once, For 1 Doses, Pre-op
<input type="checkbox"/> pregabalin (LYRICA) or gabapentin (NEURONTIN) oral doses - For Patients LESS than 65 years old (Single Response)	
<input type="checkbox"/> pregabalin (LYRICA) capsule	150 mg, oral, once, For 1 Doses, Pre-op
<input type="checkbox"/> gabapentin (NEURONTIN) capsule	900 mg, oral, once, For 1 Doses, Pre-op
<input type="checkbox"/> Opioid Analgesic Options: morPHINE or fentaNYL or HYDROmorphone (Single Response)	
<input type="checkbox"/> morPHINE injection	3 mg, intravenous, every 15 min PRN, moderate pain (score 4-6), severe pain (score 7-10), For 3 Doses, Pre-op Allowance for Patient Preference:
<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 10 min PRN, moderate pain (score 4-6), severe pain (score 7-10), For 2 Doses, Pre-op Allowance for Patient Preference:

HYDROmorphone (DILAUDID) injection 0.2 mg, intravenous, every 5 min PRN, moderate pain (score 4-6), severe pain (score 7-10), For 5 Doses, Pre-op

Pre-Operative Analgesics - NOT HMH

lidocaine PF (XYLOCAINE) 10 mg/mL (1 %) injection 0.5 mL, intradermal, once, For 1 Doses, Pre-op
May use up to 0.5 milliliters for comfort.
For PreOp Use Only

traMADol (ULTRAM) tablet 150 mg, oral, once, For 1 Doses, Pre-op
Allowance for Patient Preference:

celecoxib (CeleBREX) capsule 400 mg, oral, once, For 1 Doses, Pre-op

pregabalin (LYRICA) or gabapentin (NEURONTIN) oral doses - For Patients GREATER than or EQUAL to 65 years old (Single Response)

pregabalin (LYRICA) capsule 75 mg, oral, once, For 1 Doses, Pre-op

gabapentin (NEURONTIN) capsule 600 mg, oral, once, For 1 Doses, Pre-op

pregabalin (LYRICA) or gabapentin (NEURONTIN) oral doses - For Patients LESS than 65 years old (Single Response)

pregabalin (LYRICA) capsule 150 mg, oral, once, For 1 Doses, Pre-op

gabapentin (NEURONTIN) capsule 900 mg, oral, once, For 1 Doses, Pre-op

Opioid Analgesic Options: morPHINE or fentaNYL or HYDROmorphone (Single Response)

morPHINE injection 3 mg, intravenous, every 15 min PRN, moderate pain (score 4-6), severe pain (score 7-10), For 3 Doses, Pre-op
Allowance for Patient Preference:

fentaNYL (SUBLIMAZE) injection 25 mcg, intravenous, every 10 min PRN, moderate pain (score 4-6), severe pain (score 7-10), For 2 Doses, Pre-op
Allowance for Patient Preference:

HYDROmorphone (DILAUDID) injection 0.2 mg, intravenous, every 5 min PRN, moderate pain (score 4-6), severe pain (score 7-10), For 5 Doses, Pre-op

naloxone (NARCAN) - Respiratory Depression

For patients with Respiratory Rate LESS than 8 per minute OR if patient is stuporous or unarousable.

naloxone (NARCAN) injection 0.1 mg, intravenous, every 1 min PRN, respiratory depression, as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3)., Pre-op
Repeat Naloxone 0.1 mg once in 2 minutes if necessary (MAXIMUM 0.2 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times. Notify Anesthesia if administered.