

Alcohol Withdrawal Standing Order (CIWA) [1161]

General

Nursing

Vital Signs

<input checked="" type="checkbox"/> Vital Signs-Per unit Protocol	Routine, Per unit protocol CIWA-Ar Alcohol Withdrawal Assessment
---	---

Nursing

<input type="checkbox"/> Use the CIWA-Ar Alcohol Withdrawal Assessment Tool to assess the patient's need for symptom based treatment.	Routine, Until discontinued, Starting S
<input type="checkbox"/> Document alcohol withdrawal score and treatment and reassess on flowsheet.	Routine, Until discontinued, Starting S
<input type="checkbox"/> Assess the need for seizure and aspiration precautions.	Routine, Until discontinued, Starting S
<input type="checkbox"/> Discontinue IV	Routine, Once Discontinue maintenance IVF.

IV Fluids

Symptom Based Treatment

Mild Symptoms (Score less than 8)

<input type="checkbox"/> Mild symptoms: For CIWA score less than 8 - No treatment, reassess patient in 4 hours and document and score on MAR.	Routine, Until discontinued, Starting S
---	---

Moderate Symptoms (Score 8-15)

<input type="checkbox"/> Moderate symptoms: for CIWA score 8-15 -- Document score and reassess patient in 4 hours.	Routine, Until discontinued, Starting S
--	---

LORazepam (ATIVAN) Oral or IV

"Or" Linked Panel

<input type="checkbox"/> LORazepam (ATIVAN) tablet	1 mg, oral, every 4 hours PRN, withdrawal, for CIWA score 8-15 Give the tablet if the patient can tolerate oral medication. Indication(s): Withdrawal
<input type="checkbox"/> LORazepam (ATIVAN) injection	1 mg, intravenous, every 4 hours PRN, withdrawal, for CIWA score 8-15 Give if unable to take oral OR symptoms inadequately controlled on oral medication. Indication(s): Withdrawal

Severe Symptoms (Score GREATER than 15)

<input type="checkbox"/> Severe symptoms: for CIWA score greater than 15 - Document score and reassess patient in 4 hours.	Routine, Until discontinued, Starting S
<input type="checkbox"/> Notify Physician if CIWA greater than 20.	Routine, Until discontinued, Starting S, If CIWA greater than 20.

LORazepam (ATIVAN) Oral or IV

"Or" Linked Panel

<input type="checkbox"/> LORazepam (ATIVAN) tablet	2 mg, oral, every 4 hours PRN, agitation, for CIWA score GREATER than 15 Give the tablet if the patient can tolerate oral medication. Indication(s): Withdrawal
<input type="checkbox"/> LORazepam (ATIVAN) injection	2 mg, intravenous, every 4 hours PRN, agitation, for CIWA score GREATER than 15 Give if unable to take oral OR symptoms inadequately controlled on oral medication. Indication(s): Withdrawal

Medications

IV Access

<input type="checkbox"/> MVI 10 mL/Folic Acid 1 mg/Thiamine 100 mg intravenous infusion ("Banana Bag")	100 mL/hr, intravenous, once, For 1 Doses Per Med Staff Policy, R.Ph. will automatically switch IV to equivalent PO dose when above approved criteria are satisfied:
<input type="checkbox"/> Prevention of Wernicke's Encephalopathy: thiamine (B-1) IVPB	200 mg, intravenous, 3 times daily
<input type="checkbox"/> Treatment of Wernicke's Encephalopathy: thiamine (B-1) IVPB	500 mg, intravenous, 3 times daily

If NO IV Access

<input type="checkbox"/> thiamine (B1) IV Push Once followed by Oral Daily Starting Tomorrow	"Followed by" Linked Panel
<input type="checkbox"/> thiamine (B-1) injection	100 mg, intravenous, once, For 1 Doses
<input type="checkbox"/> thiamine tablet	100 mg, oral, daily, Starting S+1
<input type="checkbox"/> folic acid (FOLVITE) tablet	1 mg, oral, daily
<input type="checkbox"/> multivitamin tablet with minerals	1 tablet, oral, daily
<input type="checkbox"/> chlordiazepoxide (LIBRIUM) capsule	"Or" Linked Panel
<input type="checkbox"/> For CIWA score 9-15 - chlordiazePOXIDE (LIBRIUM) capsule	5 mg, oral, 3 times daily PRN, withdrawal For CIWA scores 9-15. Indication(s): Withdrawal
<input type="checkbox"/> For CIWA score GREATER than 15 - chlordiazePOXIDE (LIBRIUM) capsule	10 mg, oral, 3 times daily PRN, withdrawal For CIWA scores GREATER than 15. Indication(s): Withdrawal
<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting

Labs

Labs Today

<input type="checkbox"/> CBC with differential	Once
<input type="checkbox"/> Alcohol level, blood	Once
<input type="checkbox"/> Basic metabolic panel	Once
<input type="checkbox"/> Hepatic function panel	Once
<input type="checkbox"/> hCG, serum, quantitative	Once
	Release to patient (Note: If manual release option is selected, result will auto release 5 days from finalization.):
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site:
<input type="checkbox"/> Urine drugs of abuse screen	Once

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Consults

<input type="checkbox"/> Consult to social work - Patient requesting rehab placement	Reason for Consult: Other Specify Specify: Patient requesting rehab placement.
<input type="checkbox"/> Consult to social work - Alcohol and Drug resources	Reason for Consult: Other Specify Specify: Alcohol and Drug resources. Alcohol and Drug resources.

Additional Orders

